

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/026361

International filing date: 13 August 2004 (13.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/495,172
Filing date: 14 August 2003 (14.08.2003)

Date of receipt at the International Bureau: 19 February 2007 (19.02.2007)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1576686

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

February 12, 2007

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/495,172

FILING DATE: *August 14, 2003*

RELATED PCT APPLICATION NUMBER: PCT/US04/26361

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS *US60/495,172*



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

16623 U.S. PTO
08/14/03

PTO/SB/16 (05-03)

Approved for use through 4/30/2003. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

Express Mail Label N . EV342494129US

INVENTOR(S)

Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
Helen	Francis -Lang	San Francisco, California
Christopher G.	Winter	Palo Alto, California

☒ Additional Inventors are being named on the 1 separately numbered sheets attached hereto

TITLE OF THE INVENTION (500 characters max)

MBCATs as Modifiers of the Beta-Catenin Pathway and Methods of Use

CORRESPONDENCE ADDRESS

Direct all correspondence to:

☒ Customer Number

23500

Place Customer Number
Bar Code Label here

OR

Type Customer Number here

☐ Firm or
Individual Name

Address

Address

City

State

ZIP

Country

Telephone

Fax

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification Number of Pages

104

☐ CD(s), Number

☐ Drawing(s) Number of Sheets

☒ Other (specify)

return receipt postcard

☐ Application Data Sheet. See 37 CFR 1.76

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

☐ Applicant claims small entity status. See 37 CFR 1.27.

☐ A check or money order is enclosed to cover the filing fees

☒ The Director is hereby authorized to charge filing

fees or credit any overpayment to Deposit Account Number:

50-1108

FILING FEE
AMOUNT (\$)

160

☐ Payment by credit card. Form PTO-2038 is attached.

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are: _____.

Respectfully submitted,

SIGNATURE

TYPED or PRINTED NAME

Laleh Shayesteh

TELEPHONE

650.837.8223

[Page 1 of 2]

Date

8/14/2003

REGISTRATION NO.

47,937

(if appropriate)

Docket Number:

EX03-053P

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

19704 U.S. PTO
60/495172
08/14/03

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (05-03)

Approved for use through 4/30/2003. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Docket Number		EX03-053P
INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle [if any])	Family or Surname	Residence (City and either State or Foreign Country)
Richard Benn Abegania	Ventura	Daly City, California
Lynn Margaret	Bjerke	San Francisco, California
Kim	Lickteig	San Francisco, California
Joanne I.	Adamkewicz	San Francisco, California
HaiGuang	Zhang	El Sobrante, California
Bing	Hai	Redwood City, California

[Page 2 of 2]

Number 1 of 1

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

MBCATs AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

BACKGROUND OF THE INVENTION

The *Drosophila Melanogaster* Armadillo/beta-catenin protein is implicated in multiple cellular functions. The protein functions in cell signaling via the Wingless (Wg)/Wnt signaling pathway. It also functions as a cell adhesion protein at the cell membrane in a complex with E-cadherin and alpha-catenin (Cox et al. (1996) J. Cell Biol. 134: 133-148; Godt and Tepass (1998) Nature 395: 387-391; White et al. (1998) J Cell Biol. 140:183-195). These two roles of beta -catenin can be separated from each other (Orsulic and Peifer (1996) J. Cell Biol. 134: 1283-1300; Sanson et al. (1996) Nature 383: 627-630).

In Wingless cell signaling, beta -catenin levels are tightly regulated by a complex containing APC, Axin, and GSK3 beta /SGG/ZW3 (Peifer et al. (1994) Development 120: 369-380).

The Wingless/ beta -catenin signaling pathway is frequently mutated in human cancers, particularly those of the colon. Mutations in the tumor suppressor gene APC, as well as point mutations in beta -catenin itself lead to the stabilization of the beta -catenin protein and inappropriate activation of this pathway.

The ability to manipulate the genomes of model organisms such as *Drosophila* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Mechler BM et al., 1985 EMBO J 4:1551-1557; Gateff E. 1982 Adv. Cancer Res. 37: 33-74; Watson KL., et al., 1994 J Cell Sci. 18: 19-33; Miklos GL, and Rubin GM. 1996 Cell 86:521-529; Wassarman DA, et al., 1995 Curr Opin Gen Dev 5: 44-50; and Booth DR. 1999 Cancer

Metastasis Rev. 18: 261-284). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a “genetic entry point”) that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a “modifier” involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as beta-catenin, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

SUMMARY OF THE INVENTION

We have discovered genes that modify the beta-catenin pathway in *Drosophila*, and identified their human orthologs, hereinafter referred to as modifier of beta catenin (MBCAT). The invention provides methods for utilizing these beta-catenin modifier genes and polypeptides to identify MBCAT-modulating agents that are candidate therapeutic agents that can be used in the treatment of disorders associated with defective or impaired beta-catenin function and/or MBCAT function. Preferred MBCAT-modulating agents specifically bind to MBCAT polypeptides and restore beta-catenin function. Other preferred MBCAT-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress MBCAT gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

MBCAT modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with an MBCAT polypeptide or nucleic acid. In one embodiment, candidate MBCAT modulating agents are tested with an assay system comprising a MBCAT polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate beta-catenin modulating agents. The assay system may be cell-based or

cell-free. MBCAT-modulating agents include MBCAT related proteins (e.g. dominant negative mutants, and biotherapeutics); MBCAT -specific antibodies; MBCAT -specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind to or interact with MBCAT or compete with MBCAT binding partner (e.g. by binding to an MBCAT binding partner). In one specific embodiment, a small molecule modulator is identified using a binding assay. In specific embodiments, the screening assay system is selected from an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

In another embodiment, candidate beta-catenin pathway modulating agents are further tested using a second assay system that detects changes in the beta-catenin pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the beta-catenin pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

The invention further provides methods for modulating the MBCAT function and/or the beta-catenin pathway in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a MBCAT polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated the beta-catenin pathway.

DETAILED DESCRIPTION OF THE INVENTION

In a screen to identify enhancers and suppressors of the Wg signaling pathway, we generated activated beta -catenin models in *Drosophila* based on human tumor data (Polakis (2000) Genes and Development 14: 1837-1851). We identified modifiers of the Wg pathway and identified their orthologs. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, MBCAT genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a

defective beta-catenin signaling pathway, such as cancer. Table 1 (Example II) lists the modifiers and their orthologs.

In vitro and in vivo methods of assessing MBCAT function are provided herein. Modulation of the MBCAT or their respective binding partners is useful for understanding the association of the beta-catenin pathway and its members in normal and disease conditions and for developing diagnostics and therapeutic modalities for beta-catenin related pathologies. MBCAT-modulating agents that act by inhibiting or enhancing MBCAT expression, directly or indirectly, for example, by affecting an MBCAT function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. MBCAT modulating agents are useful in diagnosis, therapy and pharmaceutical development.

Nucleic acids and polypeptides of the invention

Sequences related to MBCAT nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) or RefSeq number), shown in Table 1 and in the appended sequence listing.

The term “MBCAT polypeptide” refers to a full-length MBCAT protein or a functionally active fragment or derivative thereof. A “functionally active” MBCAT fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type MBCAT protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of MBCAT proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one embodiment, a functionally active MBCAT polypeptide is a MBCAT derivative capable of rescuing defective endogenous MBCAT activity, such as in cell based or animal assays; the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of an MBCAT, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2). Methods

for obtaining MBCAT polypeptides are also further described below. In some embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of an MBCAT. In further preferred embodiments, the fragment comprises the entire functionally active domain.

The term "MBCAT nucleic acid" refers to a DNA or RNA molecule that encodes a MBCAT polypeptide. Preferably, the MBCAT polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with human MBCAT. Methods of identifying orthologs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, *Proc Natl Acad Sci* (1998) 95:5849-5856; Huynen MA *et al.*, *Genome Research* (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD *et al.*, 1994, *Nucleic Acids Res* 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Drosophila*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a

specified portion of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; database: European Bioinformatics Institute; Smith and Waterman, 1981, *J. of Molec.Biol.*, 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring

matrix developed by Dayhoff (Dayhoff: Atlas of Protein Sequences and Structure, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 Nucl. Acids Res. 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of an MBCAT. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of an MBCAT under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100

$\mu\text{g/ml}$ salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 $\mu\text{g/ml}$ denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

Isolation, Production, Expression, and Mis-expression of MBCAT

Nucleic Acids and Polypeptides

MBCAT nucleic acids and polypeptides, are useful for identifying and testing agents that modulate MBCAT function and for other applications related to the involvement of MBCAT in the beta-catenin pathway. MBCAT nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes.

Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins).

Overexpression of an MBCAT protein for assays used to assess MBCAT function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) *Protein Expression: A Practical Approach*, Oxford University Press Inc., New York 1999; Stanbury PF et al., *Principles of Fermentation Technology*, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) *Protein Purification Protocols*, Humana

Press, New Jersey, 1996; Coligan JE et al, Current Protocols in Protein Science (eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant MBCAT is expressed in a cell line known to have defective beta-catenin function. The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding an MBCAT polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native MBCAT gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the MBCAT gene product, the expression vector can comprise a promoter operably linked to an MBCAT gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the MBCAT gene product based on the physical or functional properties of the MBCAT protein in *in vitro* assay systems (*e.g.* immunoassays).

The MBCAT protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the MBCAT gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native MBCAT proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of MBCAT or other genes associated with the beta-catenin pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

Genetically modified animals

Animal models that have been genetically modified to alter MBCAT expression may be used in *in vivo* assays to test for activity of a candidate beta-catenin modulating agent, or to further assess the role of MBCAT in a beta-catenin pathway process such as apoptosis or cell proliferation. Preferably, the altered MBCAT expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal MBCAT expression. The genetically modified animal may additionally have altered beta-catenin expression (*e.g.* beta-catenin knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, *i.e.* mosaic animals (see, for example, techniques described by Jakobovits, 1994, *Curr. Biol.* 4:761-763.) or stably integrated into its germ line DNA (*i.e.*, in the genomic sequence of most or all of its cells). Heterologous nucleic acid is

introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster *et al.*, Proc. Nat. Acad. Sci. USA 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Pat. No. 4,873,191 by Wagner *et al.*, and Hogan, B., Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, Science (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) Nature 402:370-371; for transgenic Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830; for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a “knock-out” animal having a heterozygous or homozygous alteration in the sequence of an endogenous MBCAT gene that results in a decrease of MBCAT function, preferably such that MBCAT expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse MBCAT gene is used to construct a homologous

recombination vector suitable for altering an endogenous MBCAT gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, *Science* (1989) 244:1288-1292; Joyner *et al.*, *Nature* (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, *Science* (1989) 244:1281-1288; Simms *et al.*, *Bio/Technology* (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) *Scan J Immunol* 40:257-264; Declerck PJ *et al.*, (1995) *J Biol Chem.* 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the MBCAT gene, e.g., by introduction of additional copies of MBCAT, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the MBCAT gene. Such regulatory sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, *PNAS* (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) *Science* 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell (Sun X *et al* (2000) *Nat Genet* 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the beta-catenin pathway, as animal models of disease and disorders implicating defective beta-catenin function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered MBCAT function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered MBCAT expression that receive candidate therapeutic agent.

In addition to the above-described genetically modified animals having altered MBCAT function, animal models having defective beta-catenin function (and otherwise normal MBCAT function), can be used in the methods of the present invention. For example, a beta-catenin knockout mouse can be used to assess, *in vivo*, the activity of a candidate beta-catenin modulating agent identified in one of the *in vitro* assays described below. Preferably, the candidate beta-catenin modulating agent when administered to a model system with cells defective in beta-catenin function, produces a detectable phenotypic change in the model system indicating that the beta-catenin function is restored, i.e., the cells exhibit normal cell cycle progression.

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate the function of MBCAT and/or the beta-catenin pathway. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the beta-catenin pathway, as well as in further analysis of the MBCAT protein and its contribution to the beta-catenin pathway. Accordingly, the invention also provides methods for modulating the beta-catenin pathway comprising the step of specifically modulating MBCAT activity by administering a MBCAT-interacting or -modulating agent.

As used herein, an “MBCAT-modulating agent” is any agent that modulates MBCAT function, for example, an agent that interacts with MBCAT to inhibit or enhance MBCAT activity or otherwise affect normal MBCAT function. MBCAT function can be affected at any level, including transcription, protein expression, protein localization, and

cellular or extra-cellular activity. In a preferred embodiment, the MBCAT - modulating agent specifically modulates the function of the MBCAT. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the MBCAT polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the MBCAT. These phrases also encompass modulating agents that alter the interaction of the MBCAT with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of an MBCAT, or to a protein/binding partner complex, and altering MBCAT function). In a further preferred embodiment, the MBCAT- modulating agent is a modulator of the beta-catenin pathway (e.g. it restores and/or upregulates beta-catenin function) and thus is also a beta-catenin-modulating agent.

Preferred MBCAT-modulating agents include small molecule compounds; MBCAT-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

Small molecule modulators

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight less than 10,000, preferably less than 5,000, more preferably less than 1,000, and most preferably less than 500 daltons. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the MBCAT protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can

also be identified by screening compound libraries for MBCAT–modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the beta-catenin pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

Protein Modulators

Specific MBCAT-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the beta-catenin pathway and related disorders, as well as in validation assays for other MBCAT-modulating agents. In a preferred embodiment, MBCAT-interacting proteins affect normal MBCAT function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, MBCAT-interacting proteins are useful in detecting and providing information about the function of MBCAT proteins, as is relevant to beta-catenin related disorders, such as cancer (e.g., for diagnostic means).

An MBCAT-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with an MBCAT, such as a member of the MBCAT pathway that modulates MBCAT expression, localization, and/or activity. MBCAT-modulators include dominant negative forms of MBCAT-interacting proteins and of MBCAT proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous MBCAT-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. &

Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., *Gene* (2000) 250:1-14; Drees BL *Curr Opin Chem Biol* (1999) 3:64-70; Vidal M and Legrain P *Nucleic Acids Res* (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, *Nature* (2000) 405:837-846; Yates JR 3rd, *Trends Genet* (2000) 16:5-8).

An MBCAT-interacting protein may be an exogenous protein, such as an MBCAT-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory; Harlow and Lane (1999) *Using antibodies: a laboratory manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). MBCAT antibodies are further discussed below.

In preferred embodiments, an MBCAT-interacting protein specifically binds an MBCAT protein. In alternative preferred embodiments, an MBCAT-modulating agent binds an MBCAT substrate, binding partner, or cofactor.

Antibodies

In another embodiment, the protein modulator is an MBCAT specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify MBCAT modulators. The antibodies can also be used in dissecting the portions of the MBCAT pathway responsible for various cellular responses and in the general processing and maturation of the MBCAT.

Antibodies that specifically bind MBCAT polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of MBCAT polypeptide, and more preferably, to human MBCAT. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab').sub.2 fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Epitopes of MBCAT which are particularly antigenic can be selected, for example, by routine screening of MBCAT polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), *Proc. Natl. Acad. Sci. U.S.A.* 78:3824-28; Hopp and Wood, (1983) *Mol. Immunol.* 20:483-89;

Sutcliffe et al., (1983) *Science* 219:660-66) to the amino acid sequence of an MBCAT. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1} , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be generated against crude cell extracts of MBCAT or substantially purified fragments thereof. If MBCAT fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of an MBCAT protein. In a particular embodiment, MBCAT-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

The presence of MBCAT-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding MBCAT polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

Chimeric antibodies specific to MBCAT polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al., *Proc. Natl. Acad. Sci.* (1984) 81:6851-6855; Neuberger et al., *Nature* (1984) 312:604-608; Takeda et al., *Nature* (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. *Blood* 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 *Nature* 323: 323-327). Humanized antibodies contain ~10% murine sequences and ~90% human sequences, and thus further

reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

MBCAT-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to

about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml. Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

Specific biotherapeutics

In a preferred embodiment, an MBCAT-interacting protein may have biotherapeutic applications. Biotherapeutic agents formulated in pharmaceutically acceptable carriers and dosages may be used to activate or inhibit signal transduction pathways. This modulation may be accomplished by binding a ligand, thus inhibiting the activity of the pathway; or by binding a receptor, either to inhibit activation of, or to activate, the receptor. Alternatively, the biotherapeutic may itself be a ligand capable of activating or inhibiting a receptor. Biotherapeutic agents and methods of producing them are described in detail in U.S. Pat. No. 6,146,628.

When the MBCAT is a ligand, it may be used as a biotherapeutic agent to activate or inhibit its natural receptor. Alternatively, antibodies against MBCAT, as described in the previous section, may be used as biotherapeutic agents.

When the MBCAT is a receptor, its ligand(s), antibodies to the ligand(s) or the MBCAT itself may be used as biotherapeutics to modulate the activity of MBCAT in the beta-catenin pathway.

Nucleic Acid Modulators

Other preferred MBCAT-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit MBCAT

activity. Preferred nucleic acid modulators interfere with the function of the MBCAT nucleic acid such as DNA replication, transcription, translocation of the MBCAT RNA to the site of protein translation, translation of protein from the MBCAT RNA, splicing of the MBCAT RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the MBCAT RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to an MBCAT mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. MBCAT-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiarnidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Alternative preferred MBCAT nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are

known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al*., Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, an MBCAT-specific nucleic acid modulator is used in an assay to further elucidate the role of the MBCAT in the beta-catenin pathway, and/or its relationship to other members of the pathway. In another aspect of the invention, an MBCAT-specific antisense oligomer is used as a therapeutic agent for treatment of beta-catenin-related disease states.

Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of MBCAT activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the MBCAT nucleic acid or protein. In general, secondary assays further assess the activity of a MBCAT

modulating agent identified by a primary assay and may confirm that the modulating agent affects MBCAT in a manner relevant to the beta-catenin pathway. In some cases, MBCAT modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising an MBCAT polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. kinase activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates MBCAT activity, and hence the beta-catenin pathway. The MBCAT polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding), transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive,

colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of MBCAT and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when MBCAT-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the MBCAT protein may be assayed by various known methods such as substrate processing (e.g. ability of the candidate MBCAT-specific binding agents to function as negative effectors in MBCAT-expressing cells), binding equilibrium constants (usually at least about 10^7 M^{-1} , preferably at least about 10^8 M^{-1} , more preferably at least about 10^9 M^{-1}), and immunogenicity (e.g. ability to elicit MBCAT specific antibody in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a MBCAT polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The MBCAT polypeptide can be full length or a fragment thereof that retains functional MBCAT activity. The MBCAT polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The MBCAT polypeptide is preferably human MBCAT, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of MBCAT interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has MBCAT –specific binding activity, and can be used to assess normal MBCAT gene function.

Suitable assay formats that may be adapted to screen for MBCAT modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput

and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, *Curr Opin Chem Biol* (1998) 2:597-603; Sundberg SA, *Curr Opin Biotechnol* 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.*, Selvin PR, *Nat Struct Biol* (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, *Curr Opin Chem Biol* (2000) 4:445-451).

A variety of suitable assay systems may be used to identify candidate MBCAT and beta-catenin pathway modulators (*e.g.* U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); and U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

Protein kinases, key signal transduction proteins that may be either membrane-associated or intracellular, catalyze the transfer of gamma phosphate from adenosine triphosphate (ATP) to a serine, threonine or tyrosine residue in a protein substrate. Radioassays, which monitor the transfer from [γ - ^{32}P or ^{33}P]ATP, are frequently used to assay kinase activity. For instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from [γ - ^{33}P] ATP to a biotinylated peptide substrate. The substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, *J Biomol Screen* (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand. Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, *Dev Biol Stand* (1999) 97:121-133). Another example of antibody based assays for protein kinase

activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., *Anal Biochem* 1996 Jul 1;238(2):159-64).

Transporter proteins carry a range of substrates, including nutrients, ions, amino acids, and drugs, across cell membranes. Assays for modulators of transporters may use labeled substrates. For instance, exemplary high throughput screens to identify compounds that interact with different peptide and anion transporters both use fluorescently labeled substrates; the assay for peptide transport additionally uses multiscreen filtration plates (Blevitt JM et al., *J Biomol Screen* 1999, 4:87-91; Cihlar T and Ho ES, *Anal Biochem* 2000, 283:49-55).

Ion channels mediate essential physiological functions, including fluid secretion, electrolyte balance, bioenergetics, and membrane excitability. Assays for channel activity can incorporate ion-sensitive dyes or proteins or voltage-sensitive dyes or proteins, as reviewed in Gonzalez JE *et al.* (*Drug Discovery Today* (1999) 4:431-439). Alternative methods measure the displacement of known ligands, which may be radio-labeled or fluorescently labeled (*e.g.*, SchMid EL *et al.*, *Anal Chem* (1998) 70:1331-1338).

Transcription factors control gene transcription. Electrophoretic mobility shift assay (EMSA) or gel shift assay is one of the most powerful methods for studying protein-DNA interactions. High throughput gel shift assays for transcription factors may involve fluorescence (Cyano dye Cy5) labeled oligodeoxynucleotide duplexes as specific probes and an automatic DNA sequencer for analysis (Ruscher K, et al., (2000) *J Biotechnol* 78:163-70). Alternatively high throughput methods involve colorimetric assays (Renard P, et al. (2001) *Nucleic Acids Res* 29(4):E21), or homogeneous fluorescence assays for the detection and quantification of sequence-specific DNA-binding proteins (Heyduk T, and Heyduk E (2001) *Nat Biotechnol* 20:171-6.)

High throughput assays based on photometric analysis of the activity of decarboxylase enzymes have been described (Breuer M et al (2002) *Anal Bioanal Chem* 374:1069-73).

High-throughput photometric assays for peroxidases have been described (Smith AD et al (2001) *Int J Vitam Nutr Res* 71:87-92; Smith AD and Levander OA (2002) *Methods Enzymol* 347:113-21).

High throughput adenylyl cyclase assays are known in the art, and are also commercially available (NEN[®] Adenylyl Cyclase Activation FlashPlate[®] Assay, available from Perkin Elmer Life Sciences, Boston, MA). Examples include homogeneous cellular assays that allow direct measurement of receptor-mediated adenylyl cyclase activation/inhibition.

Apoptosis assays. Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, *Nature* 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, *J. Exp. Med.* 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, *Blood* 15:4730-41). Other cell-based apoptosis assays include the caspase-3/7 assay and the cell death nucleosome ELISA assay. The caspase 3/7 assay is based on the activation of the caspase cleavage activity as part of a cascade of events that occur during programmed cell death in many apoptotic pathways. In the caspase 3/7 assay (commercially available Apo-ONE[™] Homogeneous Caspase-3/7 assay from Promega, cat# 67790), lysis buffer and caspase substrate are mixed and added to cells. The caspase substrate becomes fluorescent when cleaved by active caspase 3/7. The nucleosome ELISA assay is a general cell death assay known to those skilled in the art, and available commercially (Roche, Cat# 1774425). This assay is a quantitative sandwich-enzyme-immunoassay which uses monoclonal antibodies directed against DNA and histones respectively, thus specifically determining amount of mono- and oligonucleosomes in the cytoplasmic fraction of cell lysates. Mono and oligonucleosomes are enriched in the cytoplasm during apoptosis due to the fact that DNA fragmentation occurs several hours before the plasma membrane breaks down, allowing for accumulation in the cytoplasm. Nucleosomes are not present in the cytoplasmic fraction of cells that are not undergoing apoptosis. An apoptosis assay system may comprise a cell that expresses an MBCAT,

and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether MBCAT function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the MBCAT plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, Int. J. Cancer 38, 369; Campana *et al.*, 1988, J. Immunol. Meth. 107, 79), or by other means.

Cell proliferation is also assayed via phospho-histone H3 staining, which identifies a cell population undergoing mitosis by phosphorylation of histone H3. Phosphorylation of histone H3 at serine 10 is detected using an antibody specific to the phosphorylated form of the serine 10 residue of histone H3. (Chadlee, D.N. 1995, J. Biol. Chem 270:20098-105). Cell Proliferation may also be examined using [³H]-thymidine incorporation (Chen, J., 1996, Oncogene 13:1395-403; Jeoung, J., 1995, J. Biol. Chem. 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [³H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter). Another proliferation assay uses the dye Alamar Blue (available from Biosource International), which fluoresces when reduced in living cells and provides an

indirect measurement of cell number (Voytik-Harbin SL et al., 1998, *In Vitro Cell Dev Biol Anim* 34:239-46). Yet another proliferation assay, the MTS assay, is based on in vitro cytotoxicity assessment of industrial chemicals, and uses the soluble tetrazolium salt, MTS. MTS assays are commercially available, for example, the Promega CellTiter 96® AQueous Non-Radioactive Cell Proliferation Assay (Cat.# G5421).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook et al., *Molecular Cloning*, Cold Spring Harbor (1989)). For example, cells transformed with MBCAT are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

Cell proliferation may also be assayed by measuring ATP levels as indicator of metabolically active cells. Such assays are commercially available, for example Cell Titer-Glo™, which is a luminescent homogeneous assay available from Promega.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW et al. (1986) *Int J Radiat Biol Relat Stud Phys Chem Med* 49:237-55). Cells transfected with an MBCAT may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system such as a cell-free assay system. A cell proliferation assay may also be used to test whether MBCAT function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the MBCAT plays a direct role in cell proliferation or cell cycle.

Angiogenesis. Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether MBCAT function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the MBCAT plays a direct role in angiogenesis. U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434, among others, describe various angiogenesis assays.

Hypoxic induction. The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with MBCAT in hypoxic conditions (such as with 0.1% O₂, 5% CO₂, and balance N₂, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g. beta-catenin is

over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether MBCAT function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the MBCAT plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF, and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., *Bioconjug Chem.* 2001 May-Jun;12(3):346-53).

Tubulogenesis. Tubulogenesis assays monitor the ability of cultured cells, generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include MatrigelTM (Becton Dickinson), an extract of basement membrane proteins containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4°C and forms a solid gel at 37°C. Other suitable matrices comprise extracellular components such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, *Nature Medicine* 5:1418-1423). These assays have traditionally involved stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a candidate agent modulates. Moreover, we have found that different target genes respond differently to stimulation with different pro-angiogenic agents, including inflammatory angiogenic factors such as TNF- α . Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing an MBCAT's response to a variety of factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF- α , ephrin, etc.

Cell Migration. An invasion/migration assay (also called a migration assay) tests the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic

signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, J Biol Chem 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for determining cell number. In another exemplary set up, cells are fluorescently labeled and migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing an MBCAT's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

Sprouting assay. A sprouting assay is a three-dimensional *in vitro* angiogenesis assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900 μ l of methocel-collagen solution and pipetted into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents

are added after 30 min by pipetting 100 μ l of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to determine the cumulative sprout length per spheroid.

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the MBCAT protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbent assay (ELISA) is a preferred method for detecting MBCAT-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also be used to test antibody modulators.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance MBCAT gene expression, preferably mRNA expression. In general, expression analysis comprises comparing MBCAT expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express MBCAT) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that MBCAT mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either

the MBCAT protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve MBCAT mRNA expression, may also be used to test nucleic acid modulators.

Secondary Assays

Secondary assays may be used to further assess the activity of MBCAT-modulating agent identified by any of the above methods to confirm that the modulating agent affects MBCAT in a manner relevant to the beta-catenin pathway. As used herein, MBCAT-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with MBCAT.

Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express MBCAT) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate MBCAT-modulating agent results in changes in the beta-catenin pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use “sensitized genetic backgrounds”, which, as used herein, describe cells or animals engineered for altered expression of genes in the beta-catenin or interacting pathways.

Cell-based assays

Cell based assays may detect endogenous beta-catenin pathway activity or may rely on recombinant expression of beta-catenin pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

Animal Assays

A variety of non-human animal models of normal or defective beta-catenin pathway may be used to test candidate MBCAT modulators. Models for defective beta-catenin pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the beta-catenin pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, beta-catenin pathway activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal beta-catenin are used to test the candidate modulator's affect on MBCAT in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the MBCAT. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on MBCAT is assessed via tumorigenicity assays. Tumor xenograft assays are known in the art (see, *e.g.*, Ogawa K et al., 2000, *Oncogene* 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the MBCAT endogenously are injected in the flank, 1×10^5 to 1×10^7 cells per mouse in a volume of 100 μ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is

assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors may be utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

In another preferred embodiment, tumorigenicity is monitored using a hollow fiber assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate modulator. Implanted cells are generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate period of time, generally around six days, the implanted samples are harvested for evaluation of the candidate modulator. Tumorigenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in soft agar, the capacity of the cells to recover and replicate in vitro, etc.

In another preferred embodiment, a tumorigenicity assay uses a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as transgenic tumor assays. In a preferred application, tumor development in the transgenic model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell carcinomas (Hanahan D, 1985, *Nature* 315:115-122; Parangi S et al, 1996, *Proc Natl Acad Sci USA* 93: 2002-2007; Bergers G et al, 1999, *Science* 284:808-812). An "angiogenic switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered

at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of small tumors (e.g., for a model of intervention), or during the growth of large and/or invasive tumors (e.g., for a model of regression). Tumorogenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

Diagnostic and therapeutic uses

Specific MBCAT-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the beta-catenin pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the beta-catenin pathway in a cell, preferably a cell pre-determined to have defective or impaired beta-catenin function (e.g. due to overexpression, underexpression, or misexpression of beta-catenin, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates MBCAT activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the beta-catenin function is restored. The phrase “function is restored”, and equivalents, as used herein, means that the desired phenotype is achieved, or is brought closer to normal compared to untreated cells. For example, with restored beta-catenin function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal relative to untreated cells. The invention also provides methods for treating disorders or disease associated with impaired beta-catenin function by administering a therapeutically effective amount of an MBCAT -modulating agent that modulates the beta-catenin pathway. The invention further provides methods for modulating MBCAT function in a cell, preferably a cell pre-determined to have defective or impaired MBCAT function, by administering an MBCAT -modulating agent. Additionally, the invention provides a method for treating disorders or disease associated with impaired MBCAT function by administering a therapeutically effective amount of an MBCAT -modulating agent.

The discovery that MBCAT is implicated in beta-catenin pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of

diseases and disorders involving defects in the beta-catenin pathway and for the identification of subjects having a predisposition to such diseases and disorders.

Various expression analysis methods can be used to diagnose whether MBCAT expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47). Tissues having a disease or disorder implicating defective beta-catenin signaling that express an MBCAT, are identified as amenable to treatment with an MBCAT modulating agent. In a preferred application, the beta-catenin defective tissue overexpresses an MBCAT relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial MBCAT cDNA sequences as probes, can determine whether particular tumors express or overexpress MBCAT. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of MBCAT expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing reagents such as the MBCAT oligonucleotides, and antibodies directed against an MBCAT, as described above for: (1) the detection of the presence of MBCAT gene mutations, or the detection of either over- or under-expression of MBCAT mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of MBCAT gene product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by MBCAT.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in MBCAT expression, the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for MBCAT expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer. The probe may be either DNA or protein, including an antibody.

EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

I. Drosophila beta-catenin screen

Two dominant loss of function screens were carried out in *Drosophila* to identify genes that interact with the Wg cell signaling molecule, beta –catenin (Riggleman et al. (1990) Cell 63:549-560; Peifer et al. (1991) Development 111:1029-1043). Late stage activation of the pathway in the developing *Drosophila* eye leads to apoptosis (Freeman and Bienz (2001) EMBO reports 2: 157-162), whereas early stage activation leads to an overgrowth phenotype. We discovered that ectopic expression of the activated protein in the wing results in changes of cell fate into ectopic bristles and wing veins.

Each transgene was carried in a separate fly stock:

Stocks and genotypes were as follows:

eye overgrowth transgene: *isow*; P{3.5 *eyeless*-Gal4}; P{arm(S56F)-pExp-UAS}}/TM6b;

eye apoptosis transgene: *y w*; P{arm(S56F)-pExp-GMR}/CyO; and

wing transgene: P{arm(Δ N)-pExp-VgMQ}/FM7c

In the first dominant loss of function screen, females of each of these three transgenes were crossed to a collection of males containing genomic deficiencies. Resulting progeny containing the transgene and the deficiency were then scored for the effect of the deficiency on the eye apoptosis, eye overgrowth, and wing phenotypes, i.e., whether the deficiency enhanced, suppressed, or had no effect on their respective phenotypes. All data was recorded and all modifiers were retested with a repeat of the original cross. Modifying deficiencies of the phenotypes were then prioritized according to how they modified each of the three phenotypes.

Transposons contained within the prioritized deficiencies were then screened as described. Females of each of the three transgenes were crossed to a collection of 4 types of transposons (3 piggyBac-based and 1 P-element-based). The resulting progeny containing the transgene and the transposon were scored for the effect of the transposon

on their respective phenotypes. All data was recorded and all modifiers were retested with a repeat of the original cross. Modifiers of the phenotypes were identified as either members of the Wg pathway, components of apoptotic related pathways, components of cell cycle related pathways, or cell adhesion related proteins.

In the second dominant loss of function screen, females of the eye overgrowth transgene were crossed to males from a collection of 3 types of piggyBac-based transposons. The resulting progeny containing the transgene and the transposon were scored for the effect of the transposon on the eye overgrowth phenotype. All data was recorded and all modifiers were retested with a repeat of the original cross. Modifiers of the phenotypes were identified as either members of the Wg pathway, components of cell cycle related pathways, or cell adhesion related proteins.

II. Analysis of Table 1

BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of *Drosophila* modifiers. The columns “MBCAT symbol”, and “MBCAT name aliases” provide a symbol and the known name abbreviations for the Targets, where available, from Genbank. “MBCAT RefSeq_NA or GI_NA”, “MBCAT GI_AA”, “MBCAT NAME”, and “MBCAT Description” provide the reference DNA sequences for the MBCATs as available from National Center for Biology Information (NCBI), MBCAT protein Genbank identifier number (GI#), MBCAT name, and MBCAT description, all available from Genbank, respectively. The length of each amino acid is in the “MBCAT Protein Length” column.

Names and Protein sequences of *Drosophila* modifiers of beta-catenin from screen (Example I), are represented in the “Modifier Name” and “Modifier GI_AA” column by GI#, respectively.

Table 1

MBCAT symbol	MBCAT name aliases	MBCAT RefSeq_NA or GI_NA	MBCAT GI_AA	MBCAT name	MBCAT description	MBCAT protein length	Modifier name	Modifier gi_aa

ADCY5	ADCY5 adenylate cyclase 5 adenylyl cyclase type V adenylyl cyclase type 5	XM_171048	22212711	adenylate cyclase 5	response to hypoxia	894	CG32158	24665354
ADCY6	KIAA0422 adenylate cyclase 6 ADCY6 adenylyl cyclase type VI	NM_015270 NM_020983	10181096	adenylate cyclase 6	regulation of smooth muscle contraction; cAMP biosynthesis	1168	CG32158	24665354
ADCY8	ADCY8 ADCY3 HBAC1 Adenylyl cyclase-8, brain adenylate cyclase 8 (brain) adenylyl cyclase type VIII	NM_001115	4557257	adenylate cyclase 8 (brain)	learning and/or memory; response to drug; signal transduction	1251	CG32158	24665354
CENTG1	CENTG1 PIKE AGAP2 GGAP2 KIAA0167 centaurin gamma1 phosphoinositide 3-kinase enhancer GTP-binding and GTPase activating protein 2 Arf GAP with GTP-binding protein-like, ANK repeat and PH domains 2 centaurin, gamma 1	NM_014770 NM_023017	7661962	centaurin, gamma 1	na	836	cenG1A	24584217
CENTG2	CENTG2 AGAP1 GGAP1 KIAA1099 GTP-binding and GTPase-activating protein 1 Arf GAP with GTP-binding protein-like, ANK repeat and PH domains 1 centaurin, gamma 2	NM_014914	7662484	centaurin, gamma 2	vesicle-mediated transport; actin cytoskeleton organization and biogenesis	804	cenG1A	24584217
CENTG3	CENTG3 MRIP-1 MRIP-1 protein centaurin, gamma 3	NM_031946	16799069	centaurin, gamma 3	cell growth and/or maintenance; signal transduction; signal transduction	875	cenG1A	24584217

DDX17	DDX17 P72 RH70 probable RNA-dependent helicase p72 DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 17 (72kD) DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 17, 72kDa DEAD (Asp-Glu-Ala-Asp) box polypeptide 17	NM_006386 NM_030881	5453840	DEAD (Asp-Glu-Ala-Asp) box polypeptide 17	regulation of transcription from Pol II promoter; RNA processing	650	Rm62	24644481
DDX5	DDX5 P68 HLR1 G17P1 HUMP68 RNA-dependent ATPase DEAD/H box-5 (RNA helicase, 68kD) DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 5 (RNA helicase, 68kD) DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 5 (RNA helicase, 68kDa) p68 DEAD box-5 DEAD (Asp-Glu-Ala-Asp) box polypeptide 5	NM_004396	4758138	DEAD (Asp-Glu-Ala-Asp) box polypeptide 5	mRNA splicing	614	Rm62	24644481
FLJ10665	FLJ10665 hypothetical protein FLJ10665	NM_018173	8922581	hypothetical protein FLJ10665	Rho protein signal transduction; cell growth and/or maintenance	790	CG7323	7293694
KIAA0720	KIAA0720 KIAA0720 protein	XM_030970	29728516	KIAA0720 protein	Rho protein signal transduction; cell growth and/or maintenance	1083	CG7323	7293694
FLJ20530	FLJ20530 hypothetical protein FLJ20530	NM_017864	8923496	hypothetical protein FLJ20530	na	557	EG:EG0003.5	18484916

GABRA2	GABRA2 gamma-aminobutyric acid (GABA) A receptor, alpha 2	NM_000807	4557601	gamma-aminobutyric acid (GABA) A receptor, alpha 2	gamma-amino butyric acid signaling pathway; gamma-amino butyric acid signaling pathway; chloride transport; regulation of neurotransmitter levels	451	GluClalpha	24648249
GABRA5	GABRA5 gamma-aminobutyric acid (GABA) A receptor, alpha 5	NM_000810	4503861	gamma-aminobutyric acid (GABA) A receptor, alpha 5	gamma-amino butyric acid signaling pathway; chloride transport; synaptic transmission; signal transduction	462	GluClalpha	24648249
GLRA1	GLRA1 STHE glycine receptor, alpha 1 (startle disease/hyperekplexia, stiff man syndrome)	NM_000171	4504019	glycine receptor, alpha 1 (startle disease/hyperekplexia, stiff man syndrome)	cell surface receptor linked signal transduction; neurogenesis; chloride transport; chloride transport; synaptic transmission; synaptic transmission; synaptic transmission	449	GluClalpha	24648249
GYG	GYG glycogenin	NM_004130	20127457	glycogenin	glycogen biosynthesis; glycogen biosynthesis; glycogen biosynthesis	350	Glycogenin	24656813
GYG2	GYG2 GN2 GN-2 glycogenin 2	NM_003918	5453674	glycogenin 2	glycogen biosynthesis; glycogen biosynthesis	501	Glycogenin	24656813
LOC350705	LOC350705 similar to Glycogenin-1	XM_301113	30147855	similar to Glycogenin-1	na	233	Glycogenin	24656813

MLLT10	MLLT10 AF10 ALL1 fused gene from chromosome 10 myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog); translocated to, 10 myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 10	NM_004641	4757726	myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 10	na	1027	Alhambra	24644741
MLLT6	MLLT6 AF17 Myeloid/lymphoid or mixed-lineage leukemia, translocated to, 6 myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog); translocated to, 6 myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 6	NM_005937	5174577	myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 6	cell growth and/or maintenance; regulation of transcription, DNA-dependent	1093	Alhambra	24644741
POU2F1	POU2F1 OCT1 OTF1 Octamer-binding transcription factor-1 POU domain, class 2, transcription factor 1	NM_002697	4505957	POU domain, class 2, transcription factor 1	transcription from Pol II promoter; regulation of transcription from Pol II promoter; negative regulation of transcription; regulation of transcription from Pol III promoter	743	pdm2	24583942
POU2F2	POU2F2 OCT2 OTF2 POU domain, class 2, transcription factor 2	NM_002698	4505959	POU domain, class 2, transcription factor 2	transcription from Pol II promoter; regulation of transcription, DNA-dependent; humoral immune response; membrane fusion; pathogenesis; development	463	pdm2	24583942

POU2F3	POU2F3 OCT11 PLA-1 Epoc-1 Skn-1a POU transcription factor likely ortholog of mouse POU domain, class 2, transcription factor 3 POU domain, class 2, transcription factor 3	NM_014352	7657409	POU domain, class 2, transcription factor 3	keratinocyte differentiation; epidermal differentiation; regulation of transcription from Pol II promoter	436	pdm2	24583942
PRKCI	PKCI DXS1179E protein kinase C, iota PRKCI PKCi	NM_002740	4506071	protein kinase C, iota	anti-apoptosis; signal transduction; signal transduction; protein amino acid phosphorylation	587	aPKC	24653760

PRKCZ	protein kinase C, zeta PRKCZ PKC2 PKCz zeta protein kinase C protein kinase C zeta PKC zeta	NM_0027 44	108646 50	protein kinase C, zeta	insulin receptor signaling pathway; cytokinesis; glucose transport; antibacterial humoral response (sensu Vertebrata); activation of MAP/ERK kinase kinase; positive regulation of transcription from Pol II promoter; actin cytoskeleton organization and biogenesis; T-cell activation; protein kinase cascade; anti-apoptosis; chemotaxis; positive regulation of cell proliferation; monocyte activation; cytokine biosynthesis; cytokine and chemokine mediated signaling pathway; cell cycle arrest; signal transduction; signal transduction; signal transduction; inflammatory response; protein amino acid phosphorylation; protein amino acid phosphorylation; protein amino acid phosphorylation; protein amino acid phosphorylation	592	aPKC	24653760
-------	--	---------------	--------------	------------------------------	--	-----	------	----------

SLC1A2	SLC1A2 EAAT2 GLT-1 glutamate/aspartate transporter II excitatory amino acid transporter 2 H.sapiens mRNA for glutamate transporter glial high affinity glutamate transporter solute carrier family 1 (glial high affinity glutamate transporter), member 2	NM_004171	4759124	solute carrier family 1 (glial high affinity glutamate transporter), member 2	neurotransmitter uptake; synaptic transmission; glutamate transport; glutamate transport	574	Eaat2	17137666
SLC1A3	SLC1A3 EAAT1 GLAST GLAST1 solute carrier family 1 (glial high affinity glutamate transporter), member 3	NM_004172	4759126	solute carrier family 1 (glial high affinity glutamate transporter), member 3	neurotransmitter uptake; neurotransmitter uptake; neurotransmitter uptake; synaptic transmission; glutamate transport; glutamate transport	542	Eaat2	17137666
SLC1A7	SLC1A7 EAAT5 excitatory amino acid transporter 5 (retinal glutamate transporter) solute carrier family 1 (glutamate transporter), member 7	NM_006671	20070239	solute carrier family 1 (glutamate transporter), member 7	vision; glutamate transport	560	Eaat2	17137666
SLC1A4	SLC1A4 SATT ASCT1 glutamate/neutral amino acid transporter alanine/serine/cysteine/threonine transporter solute carrier family 1 (glutamate/neutral amino acid transporter), member 4	NM_003038	21314632	solute carrier family 1 (glutamate/neutral amino acid transporter), member 4	neutral amino acid transport; neutral amino acid transport	532	Eaat2	17137666

SLC1A6	SLC1A6 EAAT4 excitatory amino acid transporter 4 solute carrier family 1 (high affinity aspartate/glutamate transporter), member 6	NM_005071	4827012	solute carrier family 1 (high affinity aspartate/glutamate transporter), member 6	aspartate transport; chloride transport; synaptic transmission; glutamate transport; glutamate transport; glutamate transport	564	Eaat2	17137666
SLC1A1	SLC1A1 EAAC1 EAAT3 excitatory amino acid carrier 1 excitatory amino acid transporter-3 solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1	NM_004170	18252049	solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1	synaptic transmission; glutamate transport	524	Eaat2	17137666
SLC1A5	SLC1A5 R16 AAAT ATBO M7V1 RDRC ASCT2 M7VS1 RD114 virus receptor baboon M7 virus receptor neutral amino acid transporter B solute carrier family 1 (neutral amino acid transporter), member 5	NM_005628	5032093	solute carrier family 1 (neutral amino acid transporter), member 5	neutral amino acid transport; neutral amino acid transport; EGF receptor signaling pathway	541	Eaat2	17137666
SLC7A1	SLC7A1 ERR ATRC1 CAT-1 HCAT1 REC1L ecotropic retroviral receptor amino acid transporter, cationic 1 solute carrier family 7 (cationic amino acid transporter, y+ system), member 1	NM_003045	4507047	solute carrier family 7 (cationic amino acid transporter, y+ system), member 1	basic amino acid transport; amino acid transport; arginine transport; amino acid metabolism	629	CG11128	7296600

SLC7A2	SLC7A2 ATRC2 CAT-2 HCAT2 amino acid transporter, cationic 2 solute carrier family 7 (cationic amino acid transporter, y+ system), member 2	NM_003046	4507049	solute carrier family 7 (cationic amino acid transporter, y+ system), member 2	basic amino acid transport; amino acid metabolism	657	CG11128	7296600
SLC7A3	SLC7A3 ATRC3 CAT-3 FLJ14541 solute carrier family 7 (cationic amino acid transporter, y+ system), member 3	NM_032803	17939406	solute carrier family 7 (cationic amino acid transporter, y+ system), member 3	amino acid transport; amino acid transport; amino acid metabolism; L-amino acid transport; L-amino acid transport	619	CG11128	7296600
SULF2	SULF2 SULF-2 HSULF-2 KIAA1247 sulfatase 2 similar to glucosamine-6-sulfatases	NM_018837 XM_030036	29789100	similar to glucosamine-6-sulfatases	metabolism	870	Sulf1	18487383
SULF1	SULF1 SULF-1 HSULF-1 KIAA1077 sulfatase FP sulfatase 1	NM_015170	29789064	sulfatase 1	metabolism	871	Sulf1	18487383
LOC151531	LOC151531 liver-specific uridine phosphorylase	NM_173355	27597096	liver-specific uridine phosphorylase	na	317	CG6330	23172399
UP	UP uridine phosphorylase	NM_003364 NM_181597	4507839	uridine phosphorylase	nucleobase, nucleoside, nucleotide and nucleic acid metabolism; response to drug; response to drug; response to drug; response to drug	310	CG6330	23172399

RAB32	RAB32 RAB32, member RAS oncogene family	NM_006834	5803133	RAB32, member RAS oncogene family	small GTPase mediated signal transduction; organelle organization and biogenesis; protein-vacuolar targeting; exocytosis; endocytosis; mitochondrial membrane organization and biogenesis; membrane fusion	225	Rab-RP1	6561893
RAB38	RAB38 NY-MEL-1 Rab-related GTP-binding protein RAB38, member RAS oncogene family	NM_022337	11641237	RAB38, member RAS oncogene family	small GTPase mediated signal transduction; vesicle-mediated transport	211	Rab-RP1	6561893
RAB7L1	RAB7L1 RAB7, member RAS oncogene family-like 1	NM_003929	4506375	RAB7, member RAS oncogene family-like 1	na	203	Rab-RP1	6561893
C6orf37	C6orf37 FLJ20037 retinal expressed gene C6orf37 chromosome 6 open reading frame 37	NM_017633	8923042	chromosome 6 open reading frame 37	na	447	CG30497	24586361
FLJ20202	FLJ20202 hypothetical protein FLJ20202	NM_017709	8923192	hypothetical protein FLJ20202	na	391	CG30497	24586361
MGC16491	MGC16491 MGC20845 hypothetical protein MGC16491	NM_052943	16418427	hypothetical protein MGC16491	na	424	CG30497	24586361
MGC26999	MGC26999 hypothetical protein MGC26999	NM_152630	22749287	hypothetical protein MGC26999	na	389	CG30497	24586361

III. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled MBCAT peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of MBCAT activity.

IV. High-Throughput In Vitro Binding Assay.

³³P-labeled MBCAT peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate beta-catenin modulating agents.

V. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells containing the MBCAT proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at $15,000 \times g$ for 15 min. The cell lysate is incubated with 25 μ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

VI. Kinase assay

A purified or partially purified MBCAT is diluted in a suitable reaction buffer, e.g., 50 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10 $\mu\text{g/ml}$). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100 μl . The reaction is initiated by the addition of ^{33}P -gamma-ATP (0.5 $\mu\text{Ci/ml}$) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation (Mg^{2+} or Mn^{2+}) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

VII. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues are obtained from Impath, UC Davis, Clontech, Stratagene, Ardaïs, Genome Collaborative, and Ambion.

TaqMan analysis is used to assess expression levels of the disclosed genes in various samples.

RNA is extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/μl. Single stranded cDNA is then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan assay (Applied Biosystems, Foster City, CA) are prepared according to the TaqMan protocols, and the following criteria: a) primer pairs are designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis is performed using a 7900HT instrument.

Taqman reactions are carried out following manufacturer's protocols, in 25 μl total volume for 96-well plates and 10 μl total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis is prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data are normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples are compared with matched normal tissues from the same patient. A gene is considered overexpressed in a tumor when the level of expression of the gene is 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue is not available, a universal pool of cDNA samples is used instead. In these cases, a gene is considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type is greater than 2 times the standard deviation of all normal samples (i.e., $\text{Tumor} - \text{average}(\text{all normal samples}) > 2 \times \text{STDEV}(\text{all normal samples})$).

A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating

a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

Nucleic Acid and Polypeptide sequences

>gi|27480832|ref|XM_171048.2| Homo sapiens adenylate cyclase 5 (ADCY5), mRNA

ACATCGTGGGTGTCTGCACCCACTATCCGGCTGAGGTCTCCAGAGACAGGCTTTCCAGGAGACCCGAGA
GTGCATCCAGGCGGGCTCCACTCGCAGCGGAGAACCAGCAGCAGGAACGGCTCCTGTGTCTGTCTT
CCCCGTCATGTTGCCATGGAGATGAAAGCAGACATCAACGCCAAGCAGGAGGATATGATGTTCCATAAGA
TTTACATCCAGAAACATGACAACGTGAGCATCCTGTTTGCTGACATCAGGGCTGAGGGGCTCCGACTC
AGCTGGGGGTAGACGGGCTCGAATGTGGCCTGGGAGAGCCTAGGGGGCCCCAGGGGTCTGCTTTTCTATG
TGAGCCTTTAAACTTCAGACAGGCCACCACCCTGCACCTGCAGGGGCTTTGGCACAGGAGTGCTGGCTTT
GGAGGGACTGTGGCCTTCATCGTGGTCTCTGCCCACACCTCCACGCACACAGACAGTGCCCTAGGAGGG
AAACAGAACTAATTACGAGGGGGAGGCAAGAGGACGCCAAGCAAGGAGTGGTGATTCTGAGAAAAATATT
TATTAAATAAAACAAAACAAGTTCTCCGTGCCCTTCTTTAGACTATGCTAGTTGTATGCGTGTAAGAGAC
ACACAAGCAAACGAGGACGCCACTCGGGGGGAGGGCGGGGATCCCCACTTGTCTTTTTTGTATTTTTTAT
TTTGTATTATTGAAAGCCTTGGAGATCTCACAGATAGATATGCCAAATTCTATATTTTGTAAATTCTCTA
TATTAGAAAACAGCTGTGTCACAGCAGGGCGGGTGTGCTCATTTGTAAGTGTGTATGTCGGTGTATGTAC
TGGTGTATATGTGTGTGTGTTTCATGCTGTGGAAGTGGTCTCACACAGGATGTGTTCCTTCATTTTCAGAT
TTGGCAGTTTGGGTTTTTCCAAGGTACCACCAGAGCAGTGGGTGTGTGCTTTTGGGGTACCATATGCTCA
GATTAAGTAGGAGGATGCATGGACACACTGCCCATCTTTTCTGACACACGCACACGTATGTACACACAT
GCACACACCTCCTTCCCCTAAGCAAAACGCAGATGGAATAAGAAAACAAAAGCTGCTTTCCCATCCCCA
GGCCGAGCTGGAACCAAGGGAAGCAATCTCATCTGCGACAGGCAGTGTGGTGCCTTCCACACCTTGAGA
TTTCAGACGTTTGGCGCTTACAGAGGCAGCGCCACAGATTCAGAGTGCTTACAGAAGGCCAGGTGCTT
TGAGGCTGGGACGAGGAAGCCAAGCCTCCCTGGCCTACTCAGTTGGCCAAGGTGCAGGTGGCTCTTCTCT
GGAGATGTTTCACTCAGACTGGGGGATGCAATGTGCAGCCTTCAGGTTTGCAGAAAGGGAGTGGCCTTGAC
CTCCACCGGCAAAACAGGCAGAGGAATGGGTAGAGCCAGCTTTAGAGTCCACAGGGAAAGCTAGCAGGA
ATTTTGTTTTGTGGAGGGGGCAGTTAAACATACCAAGAAAAAATACTATTTTATAACCTATGAGGA
AGACATTTGAAAATGATACTCTAGCACAGAATTCAGTGAATCCTTAGGGCCCATGCCCAATCTTTCC
ATTGCTTCTCAGGTTAGAATGATCTTCACCTCCAACATGAGCTTGGAGGTGATGAGGCAGTGGCTCTGTG
CCAGCTGCCACAATGTGACTTTGATGTCCACCTGTACCACCTCTCACTGGGCTCTAGCACACCTCCCC
TCCCCGCACACCAACTGAACACAGCTCTGAGAAGCAAAGTGTGTGGACCCAAACTGCCAAGCCTGAGTC
TGTCCCGTGCTTCTGCTGCTCCATCCTTTGAGTTCTGCATTGCCATCCTGACGTCGGCCACAGGAGGCCT
GCTTTCTTCCAGCTGTTGTTCTCAAGTTCCCTGCCCCCTCCACATCGCCCCCAGTGGTGTGGGTTTTCG
TTCTGCTTCCAACCTGAGTAAAGTGTGTGTGCTGAGTTTCATCCCATGTTCTCCCATGGTCATGGCTTCCC
GGCCCCATGGGGACCCCTCTCCCATCCCAGCAGTGACTGGTGACAGTGTGCAGGTGCAGTGCTAGCTCTT
CGTTCCCTCTAAAGGGTGTGCACTCTTTTTATTCTACTCTTGCAAAAACAGATACGATTATGATTTCCC
ATGGAAATTGAAAAGTCTATTTAAATAATTTAACTATTAAACACTTTCACTGGT

>gi|10947059|ref|NM_015270.2| Homo sapiens adenylate cyclase 6 (ADCY6), transcript variant 1, mRNA

CCGGTCTATAGGTGCATACCACACAGGCTAATTTTTTTTTTTTGGAGACGGAGTCTCACTCTGTACAC
CCAGGCTGGAGTGCAATGGCGCGATCTTTGCTCACTGCAACCTCTGCCACCCGGGTTCAAGCAATTCTCC
TGCTCAGCTCCTGAGTAGCTGGGATTACAGGCGCCCGCCACCAACGCCCGGCTAATTTTTGTATTTTATAG
TAGAGACGAGGTTTACCATCTTGGCCAGGCTGGTCTTGAACCTCTGACCTTGTGATCTACCTGCCAGGG
CCTTCCAAAGTGCTGGGATTACGGACGTGAGCCACCACGCTGGCCAGGCTAATTTTTTGAATTTTAA
TAGAGGCAGGGTCTTGTGTATTGCCCAGGCTGCTCTCACATTCCTAGCCTTAAGCCATCCTCCTATCTC
GGCCTCACAAAGTGCTGGGATTACAGGTGTAAGCCACTGTGCTGGTTTTGGGCAAGTTTCTTGTCTGTA
CTGTGTCTTGGGTTCTCACCTGTTGATTGGGGATACCCTTGTTGCCTACCTCCAAGTAATTTTATGAGGA
TGAAACTCATAGCTTGTGAGTGGGAAGTGCTTAGGTAATGAGCTACTAGTGGTGGCAGTGGCTGTGATGG
TGGTATTAATACCACTCCTACTGGATGGTCTCTAATACTATCTCCTTCCTTCCCTACCAGCAACATGTCA
TGGTTTGTGGCTCCTGGTCCCTAAAGTGGATGAACGGAACAGCCTGGGGCGAACGCAATGGGCAGA
AGCGTTTCGCGGCGCCGTGGCACTCGGGCAGGTGGCTTCTGCACGCCCCGCTATATGAGCTGCCTCCGGGA
TGCAGAGCCACCCAGCCCCACCCCTGCCGGCCCCCTCGGTGCCCCCTGGCAGGATGACGCCTTCATCCGG
AGGGGCGGGCCAGGCAAGGGCAAGGAGCTGGGGCTGCGGGCAGTGGCCCTGGGCTTCGAGGATACCGAGG
TGACAACGACAGCGGGCGGGACGGCTGAGGTGGCGCCCGACGCGGTGCCAGGAGTGGCGGATCTGCTGTG
GCGCGTCTGGTGAGGTGTTCCAGTCAAGCAGTTCCGTTTCGGCCAAGCTGGAGCGCTGTACACGCGG
TACTTCTTCAGATGAACAGAGCAGCTGACGCTGCTGATGGCGGTGCTGGTGTGCTCACAGCGGTGC
TGCTGGCTTTCCACGCGCACCCGCCCCCCTCAGCCTGCCTATGTGGCACTGTTGGCCTGTGCCCGCGC

CCTGTTTCGTGGGGCTCATGGTGGTGTGTAACCGGCATAGCTTCCGCCAGGACTCCATGTGGGTGGTGAGC
 TACGTGGTGTCTGGGCATCCTGGCGGCAGTGCAGGTTCGGGGGCGCTCTCGCAGCAGACCCGCGCAGCCCCCT
 CTGCGGGCCTCTGGTGCCCTGTGTTCTTTGTCTACATCGCCTACACGCTCCTCCCCATCCGCATGCGGGC
 TGCCGTCCTCAGCGGCCTGGGGCTCTCCACCTTGCAATTTGATCTTGGCCTGGCAACTTAACCGTGGTGAT
 GCCTTCCTCTGGAAGCAGCTCGGTGCCAATGTGCTGCTGTTCCCTCTGCACCAACGTCATTGGCATCTGCA
 CACACTATCCAGCAGAGGTGTCTCAGCGCCAGGCCTTTTTCAGGAGACCCGCGGTACATCCAGGCCCCGGCT
 CCACCTGCAGCATGAGAATCGGCAGCAGGAGCGGCTGCTGCTGTCGGTATTGCCCCAGCACGTTGCCATG
 GAGATGAAAGAAGACATCAACACAAAAAAGAAGACATGATGTTCCACAAGATCTACATACAGAAGCATG
 ACAATGTCAGCATCCTGTTTGCAGACATTGAGGGCTTCACCAGCCTGGCATCCAGTGCACTGCGCAGGA
 GCTGGTCTAGACCCTGAATTGAGCTCTTTGCCCCGTTTGACAAGCTGGCTGCGGAGAATCACTGCCTGAGG
 ATCAAGATCTTGGGGGACTGTTACTACTGTGTGTGTCAGGGCTGCCGGAGGCCCGGGCCGACCATGCCCACT
 GCTGTGTGGAGATGGGGGTAGACATGATTGAGGCCATCTCGCTGGTACGTGAGGTGACAGGTGTGAATGT
 GAACATGCGCGTGGGCATCCACAGCGGGCGCGTGCCTGCGGCGTCTTGGCTTGCGGAAATGGCAGTTC
 GATGTGTGGTCCAATGATGTGACCCTGGCCAACCACATGGAGGCAGGAGGCCGGGCTGGCCGCATCCACA
 TCACTCGGGCAACACTGCAGTACCTGAACGGGGACTACGAGGTGGAGCCAGGCCGTGGTGGCGAGCGCAA
 CGCGTACCTCAAGGAGCAGCACATTGAGACTTTCTCATCTCGGGCGCCAGCCAGAAACGGAAAGAGGAG
 AAGGCCATGCTGGCCAAGCTGCAGCGGACTCGGGCCAACCTCCATGGAAGGGCTGATGCCGCGCTGGGTTC
 CTGATCGTGCCTTCTCCCGGACCAAGGACTCCAAGGCCTTCCGCCAGATGGGCATTGATGATTCCAGCAA
 AGACAACCGGGGCACCCAAGATGCCCTGAACCTTGAGGATGAGGTGGATGAGTTCCCTGAGCCGTGCCATC
 GATGCCCCGAGCATTGATCAGCTGCGGAAGGACCATGTGCGCCGGTTTCTGCTCACCTTCCAGAGAGAGG
 ATCTTGAGAAGAAGTACTCCCGGAAGGTGGATCCCGCTTCGGAGCCTACGTTGCCTGTGCCCTGTTGGT
 CTTCTGCTTCATCTGCTTCATCCAGCTTCTCATCTTCCCACACTCCACCCTGATGCTTGGGATCTATGCC
 AGCATCTTCTGCTGCTGCTAATCACCGTGTGATCTGTGCTGTGTACTCCTGTGGTTCTCTGTTCCCTA
 AGGCCCTGCAACGTCTGTCCCGCAGCATTGTCCGCTCACGGGCACATAGCACCGCAGTTGGCATCTTTTC
 CGTCTGCTTGTGTTTACTTCTGCCATTGCCAACATGTTACCTGTAACCACACCCCATACGGAGCTGT
 GCAGCCCGGATGCTGAATTTAACACCTGTGCATCATCTGCCCTGCCACCTGCAGCAGCTCAATTAATCTCTC
 TGGGCCTGGATGCTCCCCTGTGTGAGGGCACCATGCCACCTGCAGCTTTCCTGAGTACTTCATCGGGAA
 CATGCTGCTGAGTCTCTTGGCCAGCTCTGTCTTCTGCACATCAGCAGCATCGGGAAGTTGGCCATGATC
 TTTGTCTTGGGGCTCATCTATTTGGTGTGCTTCTGCTGGGTCCCCAGCCACCATCTTTGACAACTATG
 ACCTACTGCTTGGCGTCCATGGCTTGGCTTCTTCCAATGAGACCTTTGATGGGCTGGACTGTCCAGCTGC
 AGGGAGGGTGGCCCTCAAATATATGACCCCTGTGATTCTGTGCTGTTTGGCTGGCGCTGTATCTGCAT
 GCTCAGCAGGTGGAGTCGACTGCCCGCCTAGACTTCTCTGGAACTACAGGCAACAGGGGAGAAGGAGG
 AGATGGAGGAGCTACAGGCATACAACCGGAGGCTGCTGCATAACATTCTGCCCAAGGACGTGGCGGCCCA
 CTTCTGCGCCGGGAGCGCCGCAATGATGAACTCTACTATCAGTCGTGTGAGTGTGTGGCTGTTATGTTT
 GCCTCCATTGCCAACTTCTCTGAGTTCTATGTGGAGCTGGAGGCAACAATGAGGGTGTGAGTGCCTGC
 GGCTGCTCAACGAGATCATCGCTGACTTTGATGAGATTATCAGCGAGGAGCGGTTCGGGCAGCTGGAAAA
 GATCAAGACGATTGGTAGCACCTACATGGCTGCCTCAGGGCTGAACGCCAGCACCTACGATCAGGTGGGC
 CGCTCCCACATCACTGCCCTGGCTGACTACGCCATGCGGCTCATGGAGCAGATGAAGCACATCAATGAGC
 ACTCCTTCAACAATTTCCAGATGAAGATTGGGCTGAACATGGGCCCAGTCGTGGCAGGTGTCTCGGGGC
 TCGGAAGCCACAGTATGACATCTGGGGGAACACAGTGAATGTCTCTAGTCGTATGGAAGCAGCAGGGGGTC
 CCCGACCGAATCCAGGTGACCACGGACCTGTACAGTTCTAGCTGCCAAGGGCTACCAGCTGGAGTGTG
 GAGGGGTGCTCAAGGTGAAGGGCAAGGGGAGATGACCACCTACTTCTCAATGGGGGCCCCAGCAGTTA
 ACAGGGCCCAAGCCACAAATTCAGCTGAAGGGACCAAGGTGGGCATTGAGTGGACTCTGTGCTCACTGGGT
 GGAGCTGTGGCAGGGGGCACTGAGCCTCCAGACCCTGCTAACCAACAAAGGGAACATCCAGCAGGCTGT
 GCTTGGATCATGCTCGTCTGCCCTCAAGCTGGAAAACAAGGGGCTACCTACCGAGAGGATTATGCAAGTG
 ACTTTCTTTCTTACTTGGGGTAGGGCTGTTCCCTCTCCAATCTTCCAGCCTTTGGGAGCAGGGGAGGGGT
 CAGTAGCAGAAGCAGAGGGAGGCCTCTTGCTGAGGGATTAAATGGCAGCTTGCCATGCCTACCCTTCC
 CTGTCTGTCTGGGCAGCAGGTTTACGGGCTGAGCCCTTCTTTTCCCTCTTTTTTCTGGGAATATTTTGTA
 CAATATTTTGTACAAAGACAGGCATGAGGAGTGCCTATTCCATGCTTGCCTTTGCAATACCTGCATCCCC
 AGCACTGGTCTTGGGCATTTCCCCACCCAGCCAGGTGTCCCTCCTATGCACAGAGCAGAGGAGGGAGAA
 GCTCTGGGGAGCCAGCTTTGGCCATATTTTCAGGAGAATGTTTCCATGTGCCAAATCTTAGTCCCATGATC
 TGTCCCCAAAGGGGAACAAAGGGACCTCTGACAGCTTAGATTTAGCCCCAGTTCTTGACGCTCCAGGGA
 ACGGGGTGTCTGGCCTCACTGGTACTGTGAAAAATGCTCAGAGAGCAAGCCTGTGTGTGGGGATGTGAGG
 TCAGGAGCTGGAAGTTCACCTGCAGGTGCCAAAGAGCAGGCCCGGCCAGGGCTGGGGCAGTGCCAGACTCT
 GATCTGAGGACCCCGTCGGGGTCCAGATCAGGTCACTCTGCCCCAGTGTCTCTTGTGTCTGTGCTGACAA
 GGGGGCATGGAGCATCTCTTCTCTTCTGTTGCCAAATAGAAAAGGTTCAGGGCATGGAGAAAGGTGACC
 CTGATCCCAAACCTGCCCTCCCAAGTCTCTGGTGTGGGGAGGGCCCGTGTGTTTGTGTAACGTGTGTGTG

CATGTTGGTCTTTGTGTGCATATCTGTTTTCCAGGTCTATGTGAGTCCTTGTGCTCCTGCTCCTCAGCTC
TCCACCCAGGTTGCCTCTCTCCTGTGGGCCTCTGTCTTCGGGAATAAAGCAGGGTTTCTACTTCAGG
GGATGTAGAGAGATGCCCAGGTTGCACAGGAGTGGGATGGGGTGTGGTAGCAAAAGGAGGGAGAGGAGTC
CTTTTTGTGCCAAATCCCTAAGTGCCGTTCCGGGGGCCATGTGTGCAGCATGACTCTCCCTGTCTGTGGCA
GGGACCCAAGCGCTTGCTTAAGCCCCAGTGCTCCATGCCAGCACTTGAAGTGTCTGGGGTTTGATGGACA
GAGGCTGAGGAATTTCTGGCTTCCCCAGATAGTGTCTGGGACATGGTATGCTTTGGGGCTGGGGTAGC
ATGGAATCCCTCTGAGGACCTGGATACTGGTAC TACGGGGTGGGAAGAGGAACCTTAAACTTGGCTTTC
CCCAGCCTTCAGCCTGAGTCTAGCATGTTTCTAGCTCCCCAGTCCCTTGTGAAGCCTTGAGGGCTGGCAG
AAGGGTTAGGAGTTGAACCTAGATTCCCTTCTGTCTTTGCCTTCCTTTTACCCTTTTCCCTGCAACC
TCCTTGACTCTGGCCTGAATTTGTTGGTGCCTCAGTTTCTCTGTCTGTACCTATTTAAGCCAAAGGCACT
AGCCTGAATTTTGCTTGAAGATCACTTTGTCTTGGAAATGACTAGAGAGGCAGAGGAGAAGGGTTTCCAG
AGTTGCTAGGTTTGGGAGTGAAGGGGCAGGCAGTGCCTTGCCCCCTCCTCATGCCCTTCTGACACCAG
CTCCCTGTGGAGGCCTGGTTTCTGGGTAATGCCTCCCTTGGGCATCTTCATGCATCAACCAATGGGCCA
TCAGGTACTTCATTAGTCATGGCAGAAGGAGGGGAAAAGACTTGTTTTCCAGACAGAAAATCTACTCCCC
TGTCCCCAGCCATATCCCTGGATAGGAAGGGATAGGAAGAGACTACTTGGTGCCATGGGGTAGGGGTGAG
GGTATAAGTAGATCAGAGTGGGAAGACCTCAGCCTTGGGTGGCTTGTCTCTGCTTCTTGCCAGGTGGGAG
GGCCTGTCCACACCCTGGATCCCCGTAC CACAGTGCCAGCCATGCCCTTCCCCTGGGCTACCATTGTCCC
TTTCCTCACCCAGTTGGTAGAGGAGTCAGGAGGTGGGAGGCCGTGGGGCTTTGGTTTTATAATGTAACCAC
TGTGGGGGTGGGGGAGGATGGTGAACCATGTATTTCAGTGAAATATTTAATATATTTAAATATCAATAAA
ATCAAACCTCTTTGT

>gi|4557256|ref|NM_001115.1| Homo sapiens adenylate cyclase 8 (brain)
(ADCY8), mRNA

GACTGGCTGCAGCCGAGTCTTGGTGGAGGAGGTGGTGACCACCACCGCTCCTCCACCTGCATCCGGCTG
CGGGACTGCGCGCGCGTCTTGGCTTGCAGACCCTGCACCCCGGGACGCGGCTCATCTGTATTAGCAC
CGGCACTAACCTCCACCGCTCAGCGACTTGGTCCGCCGCAAGCTCCGCCGAGGCTTTGTCCGCTAGCG
CTCGGCTGAGTCTGGGCGGGCGGGAACCTGGGCTAGGGCGAGGCGGGGCCCCCTGGACATGCCCTTCTCC
ACGTCCGCTCCTCGACCCTATTGTAAGCGGAGAACTTAGTGTGCGAGGCAGGCAGGGGACGTCCCCAT
TTTGATGTCCGTAACCTCCACCCCTTCCGATCGAGGTAGTAAAGAGGCTCCTGTAGGAACTGACTGC
CTCTATGATTGCGGCTCTTGGGGGATTTTGCCTTTAGCCCGAAAGTTGGCTTTGCCAAAAGACGCACGG
GTAGGAAGGGCGAAAAGGAAACCTGTATTCCGTCGCGCTGGGCTCTCCGAGTCCGTGCGCAAAGCGGCC
TACGAGTCTTGGCTCCGCACCTGCAGAGGACAAGAGCCAATGCC TAAAAAAGAACAGCGGAGGAACCGGC
TGGCGCGGCCAGCTGGAACGCTGGATCGCAGTGCGCCAGGGAAGGCCGGGGCGCCCGCCGGCCCTAGC
CCTCAGTGGTCTCTCCACGCCGGCCCGCGCTGCCTCTGCC TACAAGACCTGGGGCGTCTTGCCAGA
TCTGGATGGCAGGTCCCTCCGCCACCCCGGCCCGGTTCCGGGGGCGTGGCTTGGCGCGGGGGCGGGTTT
AAGTCACCGCGGGTGTCTGACCACTCTGACAGGTCTCCAAATTTCTCCAGTCCGCTGGCGCCCGGGTG
CGTTTTAGAGCTCCAGGGTGGCACGCGGCGGCGCTTCCCTAGATCCAGAGGCGTCTCTGTGCACTTCCAC
GCGGCTCGGGCTCCCTTCTCCTCCAAACCTTCGCCTCATCCGCCAAGCTTCGGTTCTCAGCCTCAGAT
ATCCGCACCGGCGGCTTCTCTTCTGTTCTCGGAGCTCTTGGCTTTGGAGCCCTCACCCTTTTCTTCCC
CCGCTCTCCTGATCTCTTAGCTCCGAGCCAAATGGACTCCAAAGAACGAAATAAAGGGGATGAGAACT
GTGTGCTGCGACCTTTCGAAAGCACAGCTGAAAGCGTTGACCTCGTCTTATAGATCAGGCTGGGACCTG
GGGCGAGAGTCCCACACCCCTCCGGGAGGGATGCTTCTGGCCAGAGCCAGCGCTGCGCTGTCACTCCT
TGCTCCCGAATAGGAAAGAGCCTAGGAGGGAGCCTCAGCATACCCCTTCTTCCAAATTAATATTTGGT
GAATTGTTAGCGCCGAGGCTACCACCTCTCCAACCTGTGCGGGGCGCGCCGCCACCTCACCGTGACCC
CCTCCTCCTCTTCTTTGCCACCGCCCCAGCTCCGCCCTGCTCCCCATCCCGGCGCAATGGAGTTCTC
CGAAGGGCGATGATTCCAGCCACATCTGCTAACTTCGCACCCATCGCTGCCGCGGTCACCGCCGGCCAG
GCCCCCTGCAGCCGCGGAGCAGTGGGCGTCCAAAGCCAGTGCAGCAGCCAGGACCCGCCCCGACGCGCAG
CAGAAGCACGGCGCCAGGCGCTTAGGCGTCTCTTGGAGAGCAAAGGCTGCGCCAAAACGCTGAGCCTAG
AATCAACCAAGGAGCCTGAGCCAGGAAGGGGCTGCGTGGCTCACAGCGCTGCGGCTCCTGAGGACAAAT
AGCCACTGCCGCTGCGTACCCAAGCTGCGCCGGCTGGCGGGAGAGCAGCACGCAAGGACGCCGAGGTCCG
CCGCGATCTTCCAGGTGCCCTTTGCCCTTGGGCACAGTATGACCCGACCTACAGGGAGCCCTAGCGCAGG
GCTCCTGCAACGGGTACGCTAGGATAAAAAGGATCCTTGCCAAGCTCCTACCAGGCCGCTTTGAGTCT
TTAGGAACCCCTCCTCCGGCTGCCTCCCCAAGGTTCTGGGCTCCTTCCCTGCGGGCCAGAGCCATGGAG
CTCTCCGATGTGCGCTGCCTTACAGGCAGCGAGGAACCTTACACCATCCACCCGACGCCCCCGGCCGGCG
ACGGCAGGAGCGCCTCCCGGCCGACGGGCTGCTGTGGCAGACGGCGGTGCGACACATCACGGAGCAGCG
CTTCATTACGGGCACCGGGGAGGACGCGGACGCGGAGTGGAGGCTCGGGCAAAGCCTCGGACCTGCG
GGCGGCGGCCCAACCAACACGCGCCGAGCTGTCAGGCGACTC GCGCTGCCCTCTACTCGCTGGGCC

CGGGAGAGCGAGCGCACAGCACCTGCGGCACCAAAGTCTTCCCGGAACGCAGCGGGAGCGGCAGTGCCAG
 CGGCAGCGGAGGCGGGGGCGACCTGGGCTTCCTGCACCTTGACTGTGCCCTTAGCAACTCGGATTTCTTT
 CTTAATGGGGGCTATAGCTACCGAGGGGTCAATTTCCCAACCTGCGCAACTCCTTCAAATCTCGGGATT
 TGGAACGCCTCTACCAGCGCTATTTCTTGGGCCAAAGGCGCAAATCGGAAGTGGTGATGAACGTGCTGGA
 CGTGCTGACCAAACCTCACTCTCTTGGTCCTACACTTGAGCCTGGCCTCGGCCCCATGGACCCGCTCAAG
 GGCATCCTGCTGGGCTTCTTACCGGCATTGAGGTAGTGATCTGCGCCCTGGTGGTGGTCAGGAAGGACA
 CCACCTCCCACACGTACCTGCAGTACAGCGGCGTGGTCACCTGGGTGGCCATGACCACCCAGATCCTGGC
 AGCAGGCTCTGGCTACGGGCTCCTGGGCGACGGCATAGGCTACGTGCTCTTCACGCTCTTCGCCACCTAC
 AGTAGCTGCGGCTACCGCTCACCTGGGCCATCCTGGCCGGCTGGGCACCTCGCTGCTGCAGGTCAATCC
 TCCAAGTGGTCATACCCCGGCTGGCGGTCAATTTCCATCAACCAGGTTGTGGCCAGGCAGTGCTATTTCAT
 GTGTATGAACACAGCTGGAATCTTCATCAGTTACCTGTGACACGGGCCAGCGCCAAGCTTTCTGGAG
 ACTCGGAGGTGTGTGGAGGCCAGGCTGCGCCTGGAGACAGAGAACCAAAGACAGGAGCGGCTCGTGCTTT
 CTGTGCTCCCCCGGTTTGTGTCTGGAATGATCAACGACATGACCAATGTGGAAGATGAGCACCTGCA
 GCACAGTTCCATCGGATCTACATCCATCGCTATGAGAACGTCAGTATTCTTTTTGCAGATGTTAAAGGA
 TTTACCAACCTCTCCACGACCTTGTCTGCTCAGGAGCTGGTCAGGATGCTCAACGAGCTCTTTGCCAGAT
 TTGATCGACTGGCCCATGAGCATCACTGCCTTCGTATTAAATCCTGGGGGACTGCTACTACTGCGTGTC
 TGGACTTCCTGAGCCCCGCCAGGACCATGCCACTGCTGTGTTGAAATGGGTCTCAGCATGATCAAAACC
 ATCAGGTATGTGCGGTCAAGGACAAAACACGATGTTGACATGAGGATTGGAATCCACTCCGGCTCGGTGC
 TGTGCGGTGTTTTGGGACTACGGAAGTGGCAGTTTGATGTCTGGTCTTGGGATGTGGATATTGCAAAACA
 ACTCGAATCTGGAGGAATCCCCGGGAGGATTCACATTTCCAAAGCCACGCTGGACTGTCTCAACGGTGAC
 TATAACGTGGAAGAGGGCCATGGTAAAGAGAGGAATGAATTCCTGAGGAAGCATAATATCGAACTTACT
 TAATTAAGCAGCCTGAGGACAGTCTGCTGTCTTGCCTGAAGATATCGTCAAGGAGTCAGTGAGCTCCTC
 AGACCGGAGAAACAGTGGGGCCACATTCAGTGAAGGATCCTGGAGCCCTGAAGTGGCTTTGATAATATC
 GTGGGGAAACAGAATACCTGCGCTGCCCCAACAAGAAATTCATAAATCTGCTTCCAAACCATCTTGAC
 AAGCTTTGATGTCCAGTCTGGGCTGAGGAAATTAACAAGAGAATAGAACATACCATCGACTTGCGGAG
 TGGCGATAAATTGAGAAGAGAGCATATCAAGCCATTCTCACTGATGTTTAAAGACTCCAGCCTGGAGCAC
 AAGTATTCTCAAATGAGGGATGAAGTGTTCAGTCAAACCTGGTCTGTGCATTTATCGTTCTTCTATTTA
 TCACGGCAATACAAAGTTTTGCTTCTTCTTCAAGAGTGATGCCAATGACCATCCAGTTCTCCATTCTGAT
 TATGCTGCACTCGGCTCTGGTCTCATCACACAGCAGAGGATTATAAATGTTTGCCCTCATCTCCGG
 AAAACTTGCTGTTGGATTAATGAGACCTATTTGGCCCGGAACGTCATCATCTTTGCATCCATTTTGATTA
 ATTTCTTGGGTGCCATCTTAAATATCCTGTGGTGTGATTTTGACAAGTCGATACCCTTGAAGAACCTGAC
 TTTCAATTCTCAGCTGTGTTTACAGATATCTGCTCTACCCAGAGTACTTTGCTTTCACGGGGGTGTTG
 GCCATGGTGACCTGTGCAGTTTTCTTCCGGCTGAAGTCCGTCTGAAGCTGGCAGTGCTGCTGATCATGA
 TTGCCATCTATGCCCTGCTCACTGAGACCGTCTACGCAGGCCTCTTTCTGCGTTATGACAACCTCAACCA
 CAGTGGAGAAGATTTCTTGGGGACCAAGGAGGTATCACTGCTACTGATGGCCATGTTCTCTCTGGCTGTG
 TTCTACCATGGACAGCAGCTGGAGTACACAGCCCGCTGGACTTCTTTGGCGAGTACAGGCCAAAGAGG
 AGATCAATGAGATGAAGGAGCTGAGGGAACACAATGAGAATGCTCCGGAATATCTTACCCAGCCATGT
 GGCCCGCCATTTCTTAGAGAAGGACCGAGACAATGAGGAGCTGTATTCTCAATCCTATGATGCTGTGGG
 GTGATGTTTGCCTCCATCCAGGATTTGCGGACTTTTACTCTCAGACTGAAATGAATAACAGGGAGTG
 AATGCCTGCGCTTGCTCAATGAGATCATTTGCTGACTTCGATGAGTTGCTTGGTGAAGACCGATTTCAGA
 CATTTGAAAAGATTAAAGACATTGGCAGCACCTACATGGCCGTGTGTCAGGCCTGTACCTGAAAAACAGCAA
 TGTGAAGACAAGTGGGGACATTTGTGTGCTCTGGCTGACTTCTCACTCGCCCTGACAGAAAGCATAACAGG
 AGATCAACAAGCATTTCATTCAACAATTTTGAAGTCCGGATTGGCATCAGCCACGGCTCAGTGGTAGCTGG
 CGTTATCGGCGCTAAGAAACCACAGTATGACATTTGGGGCAAACTGTGAACCTGGCAAGCCGAATGGAC
 AGCACGGGGGTAGTGGCCGGATCCAAGTCCCAGAGGAGACCTATCTCATCTGAAAGGACAGGGCTTTG
 CCTTTGATTACCGAGGGGAGATCTATGTGAAGGGTATCAGTGAACAGGAAGGAAAAATCAAAACGTAATT
 TCTTCTGGGAAGAGTCCAACCCAACCCATTTCATCTTGCCCCAAGAAGACTGCCTGGGCAGTACTCCCTG
 GCCGCGGTTGTCTGGGACTTGTCCAGTCCCTCAATAGGCAAAGGCAGAAGCAGCTACTCAATGAGAACA
 ACAACACAGGAATCATCAAGGGTCATTACAACCGGCGGACTTTGTTGTACCCAGCGGCACAGAGCCTGG
 AGCCCAGGCTGAAGGCACCGACAAATCTGATTTGCCATAAAAGCATTTTCTTTCTGTTTTTTTTTTTTTT
 TGTATTTCTTTTATATATAAAATAAATACTAATAAAAAGGTTTAATTTTTTTTTTAGAACAAAAA
 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAACCCCAAAAAAAAAAAAAA

>gi|7661961|ref|NM_014770.1| Homo sapiens centaurin, gamma 1 (CENTG1), mRNA

AAGGGGCTTCTGAGGTTTGGGGGCTGTAGGGCCATGGGCTCAGGGCCAGAGGTGGTGTAGCCTGGC
 AAGACAGGTCTGGGCAACATGCATGCCCAGAGGCAGTTCTGTTGTAGCTGCAGTGAGAGCAGAAGTCAGAC

GACATGAGGTGGCCAAGCAGGCTCTAAACCGCCTCAGGAAGCTGGCAGAGAGGGTGGACGACCCCGAACT
 CCAGGACAGCATCCAGGCCTCATTGGACAGCATTTCGAGAGGCTGTGATCAATAGCCAGGAATGGACTTTG
 AGCCGCTCCATTCTGAACTGCGCCTGGGTGTGCTGGGCGATGCCAGGAGTGGGAAGTCATCGCTCATCC
 ACCGATTCTGACTGGCTCATACAGGTGCTGGAGAAGACAGAGAGTGAGCAGTACAAGAAAGAAATGTT
 GGTGGATGGACAGACACATCTGGTGCTAATCCGAGAGGAAGCTGGGGCACCTGATGCCAAGTTCTCAGGC
 TGGGCAGATGCTGTGATCTTCGTCTTCAGCCTGGAGGATGAGAACAGTTTCCAGGCTGTGAGCCGTCTCC
 ATGGGCAGCTGAGTTCCCTTCGCGGGGAGGGACGAGGAGGCCTGGCCTTGGCACTGGTGGGGACACAAGA
 CAGGATCAGTGCTTCCCTCCCTCGGGTGGTGGGAGATGCTCGTGCCAGAGCTCTGTGCGCGGACATGAAA
 CGCTGCAGCTACTATGAGACTTGTGCAACCTATGGGCTCAATGTGGATCGGGTCTTCCAGGAGGTGGCCC
 AGAAGGTGGTGACCTTTCGCAAGCAGCAACAGCTTCTGGCTGCCTGCAAGTCCCTGCCCAGCTCCCCAAG
 CCACTCAGCTGCATCCACTCCGGTAGCTGGCCAGGCTAGTAACGGGGGCCACACTAGCGACTACTCTTCT
 TCCCTCCCGTCTCACCGAATGTTGGTCAACGGGAGCTCCGAGCCGAGGCAGCTGCAGTGGCTGGATTGA
 GCACCCAGGGTCCCTGCACCGGGCAGCCAAGCGCAGGACCAGCCTTTTTCGAATCGTCGGGGTAGTGA
 CTCCGAGAAACGAAGCTTGGATAGTCGGGGAGAGACAACAGGGAGTGGGCGAGCCATCCCCATCAACAG
 AGCTTCTACTAAACGAAGTGGCAATTCTTGAACAAAGAATGGAAGAAGAAATATGTAACCTGTCCA
 GTAATGGCTTTCTACTCTACCAACCCAGTATTAACGATTACATCCACAGTACCCACGGCAAGGAGATGGA
 CTTGCTGCGAACAACAGTCAAAGTCCCGGGCAAGCGGGCCCCGAGGGCCATCTCTGCCTTTGGCCCCCTCA
 GCCAGCATTAACGGGCTCGTCAAGGACATGAGCACTGTCCAGATGGGTGAAGGCCTGGAAGCCACTACTC
 CCATGCCAAGCCCCAGCCCCAGCTTCCCTGCAGCCACCACCAGATCAGACATCCAAACACCTGTCT
 GAAGCCAGACCCGAATTTGGCCCCGAGCCCTCAGCACGGACTGTACCCCATCTGGAGACCTGAGCCCCCTG
 AGTCGGGAACCCCCCTCTTCTCCCATGGTGAAGAAGCAGAGGAGGAAAAATGACAACACCATCCAAGA
 CTGAAGGCTCGGCTGGGCAGGCTGAAGAGGAAAACTTTGAGTTCTTGATCGTGTCCAGCACGGGTCTAGAC
 GTGGCACTTTGAGGCAGCCAGTTTGGAGAGCGGGATGCCTGGGTCCAGGCCATCGAGAGTCAGATCCTA
 GCCAGTCTGCAATGCTGTGAGAGCAGCAAGGTCAAGCTGCGCACAGACAGCCAAAGCGAGGCCGTGGCCA
 TCCAGGCGATCCGGAACGCCAAGGGGAATTCAATCTGCGTGGACTGCGGGGGCCCCAACCCACGTGGGC
 CAGCTTGAACCTGGGCGCCCTCATCTGCATCGAGTGTCTTGGCATCCACCGCAACCTGGGCACACACCTG
 TCCCGCGTCTCGCTCGCTGGACTTGGACGACTGGCCACGGGAGCTGACCTGGTGTGACGGCTATTGGCA
 ACGACACGGCCCAACCGCTGTGGGAAAGCGACACGCGAGGCCGTGCCAAGCCCTCGCGGGACTCTTCGCG
 GGAGGAGCGCGAGTCTGTGGATTCTCGCCCAAGTACGAGCAGCTACTGTCTCTGGCGCCGCTGAGCACCTCG
 GAGGAGCCGCTGGGCCGCCAGCTGTGGGCCGCCGTGCAGGCCAGGACGTGGCTACCGTTCTCCTGTCTTT
 TGGCCCATGCGCGACACGGGCCGCTCGACACCAGCGTAGAGGACCCACAGCTGCGCTCCCCACTCCACCT
 GGCGGCCGAGCTCGCCACGTCGTCATCACGCAACTGCTGTGTGGTACGGCGCGGACGTGGCGGCCCGT
 GACGCCCAGGGCCGACGGCGCTGTTCTACGCCCCGCCAGGCTGGAAGCCAGCTGTGCGCCGACATCCTTC
 TCCAGCACGGCTGCCCCGGGTGAGGGCGGCAGCGCGGCCACCACGCCAGCGCGGCCACCACGCCCAGCAT
 CACCGCCACGCCAGCCCCCGCCGCCGGAGCAGCGCCGCTAGCGTGGGCCGCGCCGACGCCCCGGTTGCG
 CTGGTATAGTTGCCCAGCGGGAGAGACACCCCCATCCCCACGCGGGCCGGGCACGACCACACCGGGCGGA
 CCGCTGGACAGACGCACCCACTCACCTCTCCGATCCGACCCCGCCCCACGGGAGCACTTCTTACCCCCA
 CGAGGGCACAGCCCCACGCCCTTCCAGAAGACACAACTCACCTCCCTCTCCCCAACGAGGCATGGAGAC
 CCCAGCTCAACACGCCCCCTCTCCATTGTTTCCACACTCGGATTGCTCTGGGTCTACCCGCTCGGGGTTC
 GCGGAAGGGGAGGTCCCCAGCGGTACGGGCTCCTAGGCGCTTGGACATGCCCGCTTGTCGCCCCCTAGGGG
 CGGAAAGAAGACCCGGTCTTCCCCGAGCTCAACAATCCCAGGGCGGGAGCCCATACCCAGGCTGGCTCCT
 GGGGCGACGCCACGCCAGAGGGAGGGTTTAGTACATGGGAGGGCTAGCCCCGGGACTTGGGGCCAATACG
 GAAACCCTACCCCTTCCATGCTCATCTCACTGCCCAGTGAAGGGGGCAAGGGCAGCGAGGGGCAGGACCC
 CTGCTGCTCATGGGGGTGGGGGTTCCTCAACCCTTTCCGTGAAAGGGACCGTGAGTAGTGAAACCAGGAC
 GTCCCCACACCTACCCATGGGTGGAAGCTAAAGAGTCCGGGGGAACCAAGAGCCTGGGTCCCTCCAC
 TACTATCTTGGGAGGGAGAAGGGATGGAGCAGGAGCGGGCTAAGAGGGCTGCGGGGGCTTGTAATTTAT
 TGCTTTGTTCCTCAACATGGGGATGGGGGCGGGCAGGGGCGTGGGGAGGGCGTTTTGGGGTAGGGGAAGC
 CCAGGGTGGGCAGGGGGTGGGTGGGGTGGGGTGGCTCTCGGGTTGTGTCTGTGACCGTGACTGCGTACC
 TGTGACTGTGCAGCGCCGTGGTGATCTGGGGTAGGGGGCACCCCTACAGTGGGACCCCTCCCCATTAT
 TCTTTCTGTCCAGCCCCCTCCCTCCCACTGGAGCAGCTCCAGAGCCATTCTTCAACCCCGTGACCTCTCCC
 AGCCAGGGCTGAGAGGATTCGAGGTGGCTGGGAGGGTTTCCAGGCCCCCTGCCCCCTTCCCCACAACTGG
 GTGATAGGGCGGACTTACCTGGCCCAGCCCTGCCTCCCTCAGCCAACCCAGCCCTTGGGCTGTCTGTGTC
 AGTGGTTTCCGGTCTTTTCTTTTCTTTTCTTTTCTTTTCTTTTCTTTTCTTTTCTTTTCTTTTCTTTTCT
 GAGGGGAGAGAGGTGGGGGCAAGGGGGAATGCCCCCATCTCTTGGAGGCAGTGGTGTGAATCTTCT
 TCAACAGCAATTAAAGAGGAAGTGATTTTG

>gi|7662483|ref|NM_014914.1| Homo sapiens centaurin, gamma 2 (CENTG2), mRNA

CAAGGCGCCTGCGACTCGGTCCCAGGTTCGGCGGGCGGGCGCGCGGGCGGGCTCGCGCGGGGGCCCCGGCGCG
 CCGGGCGGGCAGTACGCAGCGCGCGGACCCACGCCACGGCCAGGAGCCAGAGCAGCGCGGCCACACTG
 CCCAGGGGTTCGGCCTTCGGCCCCGGCGCTCGGAGCGCGGGCGGTGCCTGGGCTTTAATGGCTGCTCCGCG
 GAGCAGCGCCTAGGGCTGGAAGGCGGCTGCGGCTCAGGAAGTCACCCGAGCAAGCCTCCTTCGGGGCCGG
 CCGCACCCGCCCGCGCGCGCTCCATGGGGGCGCGCTCCCCCGGGCGGCCGCTGACCCGGGACGCCGGG
 GCCCGCTCGCTCGCCGGCCGCGCGCTCCCGGCCATGAAGTTCAGCCCGCGGGCCAGCCCCGCCCTGCTCCG
 CCCGCGCCTTTCTTCTCGCGCCTCCTCCGCCCGCCGCGGGCCCGGCTCCCCGGGGGCTCGCGCGCC
 CCGGGCTCGGGCGGGCCCCGGGGCGCGGGGCGGGCGGGCGGGGGGCGCGCGGCTCCGGGCGCG
 GCGCCTGCACCATGAACTACCAGCAGCAGCTGGCCAAGTTCGGCTGCCATCCGGGCCGAGATCCAGCGCTT
 CGAGTCGGTCCACCCCAACATCTACTCCATCTACGAGCTGCTGGAGCGCGTGGAGGAGCCGGTGTGCAG
 AACCAGATCCGGGAGCACGTTCATCGCCATCGAAGATGCCTTCGTGAACAGCCAGGAATGGACGCTGAGTC
 GATCTGTCCCGGAGCTCAAAGTGGGAATTGTGGGTAAGTTCGGCCAGCGGCAAGTCTGCCCTGGTGCACCG
 GTACCTGACGGGCACATATGTCCAGGAGGAGTCTCCGGAAGGTGGCAGGTTCAAGAAAGAGATTGTCTGT
 GATGGACAGAGCTATCTGTCTGCTGATCAGAGATGAAGGGGGCCCCCGGAGGCGCAGTTTGCCATGTGGG
 TGGACGCTGTTATATTTGTCTTCAGCTTGGAGGATGAAATAAGTTTCAGACCGTTTACCACTACTACAG
 TCGAATGGCCAAGTATCGGAACACGAGCGAGATTCTCTGTTTCTGGTGGGAACCCAGGATGCCATAAGT
 TCTGCTAACCCGAGGGTCATCGATGACGCCAGGGCGAGGAAGCTCTCCAACGACCTGAAACGGTGCACGT
 ACTACGAGACGTGTGCTACATACGGGCTGAATGTGGAGAGGGTCTTCCAGGACGTTGCCCAGAAGATTGT
 TGCCACAAGGAAGAAGCAGCAGCTGTCCATAGGACCCTGCAAGTCGCTACCTAATTCTCCAGCCATTCC
 TCCGTCTGTTCCGCGCAGGTGTCTGCCGTGCACATCAGCCAGACAAGTAATGGAGGTGGGAGTTTAAGCG
 ACTATTCCTCCTCCGTTCCATCGACTCCCAGCATCAGCCAGAAGGAACCTTCGGATCGATGTTCTCTCCAC
 TGCCAACACGCCCACGCCCGTTTCGCAAGCAGTCTAAGCGCCGGTCCAACCTGTTACCTCTCGGAAAGGG
 AGCGACCCAGACAAAGAGAAGAAAGGCCTGGAGAGTCTGTCGGACAGCATTGGGAGCGCCGAGCCATCC
 CAATTAACAGGCAATGCTGTTGAAGCGAAGTGGCAATCGTTGAATAAAGAGTGGAAAAAGAAATATGT
 CACCTGTGTGACCAATGGCGTGTGACCTATCATCCCAGTTTACATGATTACATGCAGAATGTTTCATGGT
 AAGGAGATTGACCTTCTGAGAACCCTGTGAAAGTCCCAGGGAAGAGGCCACCCCGAGCCACGTACGCT
 GCGCACCCATCTCCAGCCCTAAAACCAATGGCCTATCCAAGGACATGAGCAGTTTACACATCTCACCCAA
 TTCAGACACAGGGCTGGGTGACTCCGTATGCTCCAGCCCCAGTATCTCCAGCACACCAGCCCCAAGCTC
 GACCCGCCCCCTCCCTCAGCCCAACAGAAAGAAGCACCAGGAAGAAAAGCACTAGCAACTTCAAAG
 CCGACGGCCTGTCCGGCACTGCTGAAGAACAAGAAGAAAATTTTGAAGTTTATCATTGTGTCCCTCACTGG
 CCAAACATGGCACTTTGAAGCCACGACGTATGAGGAGCGGGACGCCTGGGTCCAAGCCATCGAGAGCCAG
 ATCCTGGCCAGCCTGCAGTCGTGCGAGAGCAGCAAGAACAAGTCCCGGCTGACGAGCCAGAGCGAGGCCA
 TGGCCCTGCAGTCGATCCGGAACATGCGCGGGAACCTCCACTGTGTGGACTGCGAGACCCAGAATCCCAA
 CTGGGCCAGTTTGAACCTTGGGAGCCCTCATGTGCATCGAATGCTCAGGGATCCACCGGAATCTTGGCACC
 CACCTTTCCCGAGTCCGATCTCTGGACCTGGATGACTGGCCAATCGAGCTCATCAAGGTGATGTATCCA
 TCGGGAACGAGCTAGCCAACAGCGTCTGGGAAGAGAGCAGCCAGGGGCGGACGAAACCATCGGTAGACTC
 CACAAGGGAAGAGAAGGAACGGTGGATCCGTGCCAAGTACGAGCAGAAGCTCTTCTGGCCCCGCTGCCC
 TGCACGGAGCTGTCCCTGGGCCAGCACCTGCTGCGGGCCACCGCCGACGAGGACCTGCGGACGGCCATCC
 TGCTGCTGGCACACGGCTCCCGGGACGAGGTGAACGAGACCTGCGGGGAGGGAGACGGCCGACGGCGCT
 GCATCTGGCCTGCCGAAGGGGAATGTGGTCTCTGGCGCAGCTCCTGATCTGGTACGGAGTGGACGTCACG
 GCCCCGAGATGCCCCACGGGAACACAGCTCTGGCCTACGCCGCGCAGGCCTCCAGCCAGGAGTGCATCGACG
 TGCTGCTGCAGTACGGCTGCCCCGACGAGCGCTTCGTGCTCATGGCCACCCCTAACCTGTCCAGGAGAAA
 CAATAACCGGAACAACAGCAGTGGGAGGGTGCCACCATCATCTGAGGAACAGCCGTGCCCGCCTGCTCG
 CCGCACCTGGGACGCGGCAGCCTCGCCGATTTCTCGCTCAGAAGTCGCAGCACGTGAGTCCCGTCGCATC
 CCTCCCTCTTCTGGTGGCCACCTCCCTCCCGCCACCCACTCTCACCCAAAACAAATCACAAAACCT
 GGACATCCCTCAAGGGGCGAAGAGGCGGCCGGGAGACTGCAGAAGTGGCTCCTTTTCATAAACTCCCCTA
 AACCACACACAGGAGAGAGCGACGGCCCTCGGCCCTTTGATGATAGCACATGGCGCAGGACCCTTGTCTT
 GGTGGCACAAGGGATGGGGACGCGAGGGGGAGGGGAGGCGAGGAACAAGGAGAAGGGGCAACTTTCTCTTA
 ACTGGCAGTTGAGCACATAGTACATTTCCCTCTACCAAACGGAACACTTGGATTCCATCTCTTCTCTGA
 GGAGCTCGACGGCATAAATCAGAAGCAAGCACAGAGTTTGTGAGGTTTGAAGCCCCCTATGATGGTGTGTG
 TCAAATCAGTTGTAGCTAATCTGTCCAGGGAGAATACTGGCTTCATTACACTTGTACAGCCGAGTTCTTC
 CCGCATTAAGTGTGTTAATAGAAGCTGATTAGTCATCGCCGAGAAGAAAGCATATTAGCCGAGGAGGTA
 GTCACGCGGCACGCGCCGGTGATTGCCACGATGTGATTGCAATACTCTTAGAAGCACCATATTATCCAG
 ACATGTTCTTTCAAGCCCTTGGAGCCCTCTCTAAATTCATGTATCATTTAGTATCTGTTTAAATTTTC
 AGTCCAAAGAGAGGAAATCAGTCGCTGAGTATTATTGACTCCGGTCTCCTTGGTGCAAAAACAAAATGG

GAAAAATAAATAAGAATAACTCAGAACTCAAAAGGAAACCACAAATTCAGCTAATAATAGCATTTTCGAG
TATATTTTCGTAAACTAAGGAAATACACAAAAGGCTGTTTTTTTCCGACTGTAAGAGATATTTGATGTCCT
TTTGCCGAGGTGGATGTGTTAGTCTCAGGCCCTCCTGGACCACGTTGCCCAAGTCACACAGGCTTCTGTG
TTATGTATTTAGATAAGATGTGTGAAAAATATATTTGAATAAAAGAAGTTCAT

>gi|32307157|ref|NM_031946.3| Homo sapiens centaurin, gamma 3 (CENTG3), mRNA

GCCATGAAC TTCCAGGCGGGCGGGGGG CAGAGCCCGCAGCAGCAGCAGCTGGCCGGCGGGCCCCC
AGCAGTTCGCGCTCTCCAAC TCCGCGGCCATCCGGGCGGAGATCCAGCGCTTCGAGTCCGTGCATCCCAA
TATCTACGCCATCTACGACCTGATCGAGCGCATCGAGGATTTGGCGCTGCAGAACCAGATCCGGGAGCAC
GTCACTCCATCGAGGACTCGTTTGTGAACAGCCAGGAGTGGACGCTGAGCCGCTCCGTACCGGAGCTTA
AAGTGGGCATAGTGGGGAACCTGTCTAGCGGGAAGTCAGCCCTGGTGCACCGCTATCTGACGGGGACCTA
TGTCCAGGAGGAGTCCCCTGAAGGGGGCGGTTTAAAGAAGGAGATTGTGGTGGATGGCCAGAGTTACCTG
CTGCTGATCCGAGATGAAGGAGGCCCCCTGAGCTCCAGTTTGTGCTGGGTGGATGCAGTGGTGTGTTG
TGTTTACGCTGGAGGATGAAATCAGTTTCCAGACGGTGTACAAC TACTTCTGCGTCTCTGCAGCTTCCG
CAACGCCAGCGAGGTGCCCATGGTGCTTGTGGGCACGCAGGATGCCATCAGCGCTGCGAATCCCCGGGTT
ATCGACGACAGCAGAGCCCCGAAGCTCTCCACAGATCTGAAGCGGTGCACCTACTATGAGACGTGCGCGA
CCTACGGGCTCAATGTGGAGCGTGCTTCCAGGACGTGGCCAGAGGTAGTGGCCTTGCGAAAGAAGCA
GCAACTGGCCATCGGGCCCTGCAAGTCACTGCCCCAAGTCGCCCAGCCACTCGGCCGTGTCCGCCGCTCC
ATCCCCGGCCGTGCACATCAACCAGGCCACGAATGGCGGGCGGAGCGCCTTCAGCGACTACTCGTCTTCAG
TCCCCCTCCACCCCCAGCATCAGCCAGCGGGAGCTGCGCATCGAGACCATCGCTGCCTCCTCCACCCCCAC
ACCCATCCGAAAGCAGTCCAAGCGGGCGCTCCAACATCTTCACGTCTCGGAAGGGTGTGACCTGGACCGG
GAGAAGAAGGCTGCCGAGTGCAAGGTGGACAGCATCGGGAGCGGCCGCGCATCCCCATCAAGCAGGGGA
TCCTGCTAAAGCGGAGCGGCAAGTCCCTGAACAAGGAGTGAAGAAGAAGTATGTGACGCTCTGTGACAA
CGGGCTGCTCACCTATCACCCAGCCTGCATGATTACATGCAGAACATCCACGGCAAGGAGATTGACCTG
CTGCGGACAACGGTGAAGTGCCAGGGAAGCGCTGCCCCGAGCCACACCTGCCACAGCCCCGGGCACCA
GCCCCCGTGCCAACGGGCTGTCCGTGGAGCGGAGTAACACACAGCTGGGTGGGGGCACAGGTGCCCCCA
CTCGGCCAGCAGCGCATCCCTGCACTCTGAGCGCCCCCTCAGCAGCTCGGCCTGGGCTGGCCCCGCGCCT
GAGGGGCTGCACACAGCGCTCCTGCTCCGTTTCCAGCGCCGACCAGTGGAGTGGAGGCCACCACTTCCCTGC
CCCCAGGCATGCAGCACCTTGCAGTGGCCAGCTGAGGTACTCAGTTCCAGCCCCAAGCTGGATCCTCC
CCCATCTCCCCACTCCAACCGGAAGAAGCACCGGAGGAAAAAGAGCACCGGGACCCCCGACCAGACGGC
CCCAGCAGTGCTACTGAAGAGGCAGAGGAGTCGTTTGAATTTGTGGTGGTGTCCCTCACTGGGCAGACGT
GGCACTTCGAGGCTTCAACGGCGGAGGAGCGGGAGCTGTGGGTTTCAAGTGTGCAGGCCAGATCCTTGC
CAGCCTGCAAGGCTGCCGAGTGCCAAGGACAAGACTCGACTGGGGAACGAGAACGCAGCTCTGGCTGTG
CAGGCCGTCCGCACCGTCCGCGGCAACAGCTTTTGTATCGACTGCGATGCACCCAATCCAGACTGGGCCA
GCCTGAACCTGGGTGCCCTGATGTGCATTGAGTGCTCAGGCATCCACCGACACCTGGGGGCTCACCTGTC
CCGGGTGCGCTCCCTTGACCTCGATGACTGGCCGCTGAGCTGCTGGCTGTGATGACTGCCATGGGCAAT
GCCCTCGCCAACAGCGTCTGGGAGGGGGCCTTGGGTGGCTACTCCAAGCCAGGGCCTGATGCCTGCAGAG
AGGAGAAGGAACGCTGGATACGGGCCAAGTATGAACAGAAGCTCTTCTGGCCCCACTGCCAAGCTCAGA
TGTGCCACTGGGGCAGCAGCTGCTCCGGGCCGTGGTGAAGATGACCTGCGGCTGTTGGTGATGCTCCTG
GCACATGGCTCCAAAGAGGAGGTGAATGAGACCTATGGGACGGGGACGGGCGGACGGCTCTACATCTCT
CCAGTGCCATGGCCAACGTTGTCTTCACGCAGCTGCTCATCTGGTACGGGGTGGACGTGAGGAGCCGGGA
CGCCCCGGGCGCTGACTCCACTGGCATATGCTCGCCGGGCGGCGAGCCAGGAGTGTGCAGACATCTTGATC
CAGCATGGCTGCCCTGGGGAGGGCTGTGGCTTAGCGCCTACCCCCAACAGAGAGCCTGCCAATGGCACCA
ACCCCTCTGCTGAGCTGCACCGTAGTCTTAGCCTCCTATAAGGCCCAGGAAGAGGGCAGAGGGGCCAGAA
GGACTCCATGGCCCAAAGACCTCCTCCCTGCAGGCACTGTGGGAACAGACACAGAGATGGAGAAGCAGG
GACATGCTGAGAGGACGAAGCCAAGGAAATTAGGGAGGAGAGTCAAAGGGATCAAGGAGAGTTGGGGATT
TGAGCTGCAGCAGAGAGGGATGAGGGATTTAGCCCTCTGCCCTAAGGTGCCATTGAAAAGGGACAGGACC
CTTCGAGAGTGCCTGTGAGGAGAGGGGAGCAGGACCTCTCCCTCCTCCAGATCCCTGCCTCCTAGTGCCA
GCCCCTCACACGCCTTCATCTGAAACAGGAAGAGGACGGCACCAAGTTGGGGGTGCTGGATGAAAGAGA
CGAGGGGTGATCTGTGAGTCCCATGTAACTTTGTACATTGGAATATTTATGTTTGTGTACATATTTGAT
GTGTGTGTGTATGATGAGCCAATAAACCAGACTGTGTGCGTGGCAAAAAAAAAAAAAAAAAAAAAAAAAA
AA

>gi|13787202|ref|NM_006386.2| Homo sapiens DEAD (Asp-Glu-Ala-Asp) box
polypeptide 17 (DDX17), transcript variant 1, mRNA
ATTTTGTGCAGTCGCTGGGAAGGAAGGAGACGCCTAAACCGCGGCACTGCCCGGTTTGAGCGTAGCCAAA

CCTGCCACCGGCTTTGTAGCCCCGATTCTCTGTGTTTGTCTCCGCTCTCCGACGAGAGAGGCGGGCAGC
 GTGGCGTCTGCGACGGGAGACAGCGCTCGGAGCGAGAGCGCTGCGCCTGCCGCCGCCAACAGCGG
 AGGCGCCGCCGCCATCGGTCGTACACAGACCGGAGCCGAGGCCCTCCCGAGCCCGGCCATCCGTGCCCC
 GCTCCAGATCTCTATCCTTTTGGGACCATGCGCGGAGGAGGCTTTGGGGACCAGGACCGGGATCGTGAC
 CGTGGAGGATTTGGAGCAAGAGGTGGTGGTGGCTTCCCCGAAGAAATTTGGTAATCCTGGGGAGCGTT
 TGCCTAAAAAAGTGGGATTTGAGTGAGCTCCCCAAGTTTGAGAAAAATTTTATGTGGAACATCCGGA
 AGTAGCAAGGCTGACACCATATGAGGTTGATGAGCTACGCCGAAAGAAGGAGATTACAGTGAGGGGGGA
 GATGTTTGTCTAAACCCGTGTTTGCCTTCCATCATGCTAACTTCCACAATATGTAATGGATGTGTTGA
 TGGATCAGCACTTTACAGAACCAACTCCAATTGAGTCCAGGATTTCCGTGGCTCTTAGTGGCCGGGA
 TATGGTGGGCATTGCTCAGACTGGCTCTGGGAAGACGTTGGCGTATCTCCTGCCTGCAATTGTTTCATATT
 AACCACCACTACTTTGGAAAGGGGAGATGGCCCAATCTGTCTAGTTCTGGCTCCTACCAGAGAGCTTG
 CCCGCAAGTGAAGGTGGCCGATGACTATGGCAAATGTTCTAGATTGAAGAGTACTTGTATTTATGG
 AGGTGCTCCTAAAGGTCCCCAGATTGAGACTTGGAAAGAGGTGTTGAGATCTGCATAGCCACTCCTGGA
 CGTCTGATAGATTTTCTGGAGTCAGGAAAGACAAATCTTCGCCGATGTACTTACCTTGTATTGGACGAAG
 CTGACAGAATGCTTGATATGGGGTTTGAACCCAGATCCGTAATTTGTTGACCAATCAGGCCTGATAG
 GCAGACACTGATGTGGAGTGCAACCTGGCCAAAAGAAGTAAGACAGCTTGCAGAGGATTTCTTTCGTGAT
 TACACCCAGATCAACGTAGGCAATCTGGAGTTGAGTGCCAACCAACATCCTCCAGATAGTGGATGTCT
 GCATGGAAAGTGAAAAAGACCACAAGTTGATCCAACATAAGGAAGAAATAATGGCTGAAAAGGAAACAA
 AACAATAATATTTGTGGAGACAAAGAGACGCTGTGATGATCTGACTCGAAGGATGCGCAGAGATGGTTGG
 CCAGCTATGTGTATCCATGGAGACAAGAGTCAACAGAAAGAGATTGGGTACTTAATGAGTTCCGTTCTG
 GAAAGGCACCCATCCTTATTGCTACAGATGTAGCCTCCCGTGGGCTAGATGTGGAAGATGTCAAGTTTGT
 GATCAACTATGACTATCCAAACAGCTCAGAGGATTATGTGCACCGTATTGGCCGAACAGCCCGTAGCACC
 AACAAGGGTACCGCTATACCTTCTTCACCCAGGGAACCTAAAACAGGCCAGAGAGCTTATCAAAGTGC
 TGGAAGAGGCCAATCAGGCTATCAATCCAAACTGATGCAGCTTGTGGACCACAGAGGAGCGGGGAGG
 CGGGGGTGGTCTTCTCGTTACCGGACCACTTCTTCAGCCAACAATCCAATCTGATGTATCAGGATGAG
 TGTGACCGAAGGCTTCGAGGAGTCAAGGATGGTGGCCGGAGAGACTCTGCAAGCTATCGGGATCGTAGTG
 AAACCGATAGAGCTGGTTATGCTAATGGCAGTGGCTATGGAAGTCCAATTTCTGCCTTTGGAGCAAAAGC
 AGGCCAATACCTATGGTCAAGGCACCTATGGGGCAGCTGCTTATGGCACCAGTAGCTATACAGCTCAA
 GAATATGGTGCTGGCACTTATGGAGCTAGTAGCACCACCTCAACTGGGAGAAGTTCACAGAGCTCTAGCC
 AGCAGTTTGTGGGATAGGCCGGTCTGGGCAGCAGCCACAGCCACTGATGTCAACAGTTTGCACAGCC
 TCCGGGAGCTACCAATATGATAGGTTACATGGGGCAGACTGCCTACCAATACCTCCTCCTCCTCCCCCT
 CCTCCTCCTTACGTAATGAAACCACTCAAGTGGTAGTGACTCCAGCAGACTTAATTACATTTTAAAGGA
 AACTGTCTTTCTTTTTTTTCTCCTCTTCGCCTTTTCTTTTTTTTTTCTTTTTTTTTTTTTTTTTTT
 AATTTTTTCCCCCAACCATCGTGATTTGTCTTTTCATGCAGATTAGTTAGAATTCAGTGCCAGGTTTCTT
 CTGCCCCACCAAAATGATCCAGTCTGGAATAACATTTTGTAAAAAATAAAAAAAAAA

>gi|13514826|ref|NM_004396.2| Homo sapiens DEAD (Asp-Glu-Ala-Asp) box
 polypeptide 5 (DDX5), mRNA

ACCTCATTCAATTTCTACCGGTCTCTAGTAGTGCAGCTTCGGCTGGTGTATCGGTGTCTTCTCCGCTG
 CCGCCCCGCAAGGCTTCGCCGTATCGAGGCCATTTCCAGCGACTTGTGCGACGCTTTTCTATATACTT
 CGTTCCCCGCCAACCGCAACCATTTGACGCCATGTGGGTTATTGAGTGACCGAGACCGCGGGGACC
 GAGGGTTTGGTGCACCTCGATTTGGAGGAAGTAGGGCAGGCGCTTATCTGGAAAGAAGTTTGGAAACCC
 TGGGGAGAAATTAGTTAAAAAGAAGTGGAATCTTGATGAGCTGCCATAATTTGAGAAGAATTTTATCAA
 GAGCACCTGATTTGGCTAGGCGCACAGCACAGAGGTGGAACATACAGAAGAAGCAAGGAAATTACAG
 TTAGAGGTACAACTGCCCGAAGCCAGTTCTAAATTTTATGAAGCCAATTTCCCTGCAATGTATGGA
 TGTTATTGCAAGACAGAATTTCACTGAACCCACTGCTATTCAAGCTCAGGGATGGCCAGTTGCTCTAAGT
 GGATTGGATATGGTTGGAGTGGCACAGACTGGATCTGGGAAAACATTGTCTTATTGCTTCTGCCATTG
 TCCACATCAATCATCAGCCATTCTAGAGAGAGGCGATGGGCCTATTTGTTTGGTGTGGCACCAACTCG
 GGAAGTGGCCCAACAGGTGCAGCAAGTAGCTGTGAATATTGTAGAGCATGTGCTTGAAGTCTACTTGT
 ATCTACGGTGGTGTCTCTAAGGGACCACAAATACGTGATTTGGAGAGAGGTGTGGAAATCTGTATTGCAA
 CACCTGGAAGACTGATTGACTTTTATAGAGTGTGGAATAACCAATCTGAGAAGAACAACCTACCTTGTCTT
 TGATGAAGCAGATAGAATGCTTGATATGGGCTTTGAACCCCAATAAGGAAGATTGTGGATCAAATAAGA
 CCTGATAGGCCAACTCTAATGTGGAGTGCAGCTTGGCCAAAAGAAGTAAGACAGCTTGTGAGATTTC
 TGAAAGACTATATTATATAAACATTGGTGCAGTTGAAGTGAAGTGAAGACCAACATCTTCAGATTGT
 GGATGTGTGTCATGACGTAGAAAAGGATGAAAACCTATTTCGTCTAATGGAAGAGATCATGAGTGAGAAG
 GAGAATAAAACCATTTGTTTTGTGGAAACCAAGAAGATGTGATGAGCTTACCAGAAAAATGAGGAGAG
 ATGGGTGGCCTGCCATGGGTATCCATGGTGACAAGAGTCAACAAGAGCGTACTGGGTTCTAAATGAATT

CAAACATGGAAAAGCTCCTATTCTGATTGCTACAGATGTGGCCTCCAGAGGGCTAGATGTGGAAGATGTG
 AAATTTGTCAATATGACTACCCTAACTCCTCAGAGGATTATATTCATCGAATTGGAAGAACTGCTC
 GCAGTACCAAAACAGGCACAGCATACACTTTCTTTACACCTAATAACATAAAGCAAGTGAGCGACCTTAT
 CTCTGTGCTTCGTGAAGCTAATCAAGCAATTAATCCCAAGTTGCTTCAGTTGGTTCGAAGACAGAGGTTCA
 GGTCGTTCCAGGGGTAGAGGAGGCATGAAGGATGACCGTCGGGACAGATACTCTGCGGGCAAAGGGGTG
 GATTTAATACCTTTAGAGACAGGGAAAATTATGACAGAGGTTACTCTAGCCTGCTTAAAAGAGATTTTGG
 GGCAAAACTCAGAATGGTGTTTACAGTGTGCAAATTACACCAATGGGAGCTTTGGAAGTAATTTTGTG
 TCTGCTGGTATACAGACCAGTTTTAGGACTGGTAATCCAACAGGGACTTACCAGAATGGTTATGATAGCA
 CTCAGCAATACGGAAGTAATGTTCCAAATATGCACAATGGGATGAACCAACAGGCATATGCATATCCTGC
 TACTGCAGCTGCACCTATGATTGGTTATCCAATGCCAACAGGATATTCCCAATAAGACTTTAGAAGTATA
 TGTAAATGTCTGTTTTTTCATAATTGCTCTTTATATTGTGTGTTATCTGACAAGATAGTTATTTAAGAAAC
 ATGGGAATGTCAGAAATGACTGCAGTGCAGCAGTAATTATGGTGCACTTTTTCGCTATTTAAGTTGGATA
 TTTCTCTACATTCTGAAACAATTTTTAGGTTTTTTTTGTACTAGAAAATGCAGGCAGTGTTCACAAA
 AGTAAATGTACAGTGATTGAAATACAATAATGAAGGCAATGCATGGCCTTCCAATAAAAAATATTTGAA
 GACTGAAAAAAAAA

>gi|8922580|ref|NM_018173.1| Homo sapiens hypothetical protein FLJ10665
 (FLJ10665), mRNA

AAGCGCTCGCGGGGGTCTGGCCCGGAGTGGAGGGCGCGGGTCCCAGCCCTCCCGCTCGGCGGGCGGGT
 GTCGAGTTCAGCCCTAGGGGACCTCTTTCTCCTGGACATTGAAGATATGGCCCTTTGGAGGTGACCCAGG
 AGAGAAGGGATGAAGGCCTTTGGTCTCCACATGAGGGCCCCCTCCAAGGACTCGTGGCCTCCCGCATTG
 AGACTTATGGGGGCCGGCATCGAGCCTCTGCTCAGAGCACTACTGGCAGACTCTATCCCCGAGGATACCC
 TGTGCTGGATCCCAGTCGCCGACGCCTCCAGCAGTATGTCCCTTTGCCAGGGGTTCTGGCCAGGCCGGA
 GGCTGTCAACCATGAGACTGCGAGATCCAGAGCCCGAGAAGAGGCACGGGGGCCATGTGGGGGCTGGCC
 TGCTTCACTCCCCAACTCAAGGAATCAACAAAGGCCATGAGCTGGAGGTGAGGCTGCACACTTTTCAG
 CATGTTTGGGATGCCCCGGCTGCCCCCTGAGGACCGCGGCACTGGGAGATAGGAGAGGGTGGCGACAGT
 GGCTGAGCCATGAGGAGTCTGGAGGGAGCTGGTGGCTGGGCACAAGGAGATGAGCCAGGAGCTCTGCC
 ACCAACAGGATGCCCTGTGGGAGCTCCTGACCACCGAGCTGATCTACGTGAGAAAGCTCAAGATCATGAC
 TGATCTGCTAGCCGCCGGCCTGCTGAACCTGCAGCGAGTGGGACTGCTGATGGAAGTGTGAGCTGAGACC
 CTGTTTGGAAATGTCCCCAGCCTGATTGCAACCCACCGGAGCTTTTGGGATGAGGTGCTGGGGCCACCC
 TGGAGGAGACTCGGGCCTCGGGCCAGCCTCTGGACCCATTGGTCTGCAAAGTGGCTTCCTGACGTTTGG
 CCAGCGGTTCCACCCCTATGTCCAGTACTGCCTCCGAGTGAAGCAGACCATGGCTTACGCCCCGAGAACAG
 CAAGAACTAACCTCTCTTCCATGCCTTCGTGCAGTGGTGTGAGAAGCACAAGCGCTCTGGGAGGCAGA
 TGCTCTGTGACTTGCTTATCAAGCCCCACAGCGCATACCAAGTACCCACTGCTGCTCCATGCTGTGCT
 CAAGAGGAGCCCCGAGGCACGAGCCCAAGAGGCCCTGAATGCCATGATTGAAGCCGTGGAGTCATTCTTG
 CGACATATCAATGGGCAGGTCCGCCAGGGCGAAGAGCAAGAGAGCTTGGCGGCTGCAGCACAAACGCATCG
 GGCCCTACGAGGTGCTGGAGCCACCCAGTGATGAGGTGGAGAAGAACCCTGCGCCCATTTCTCCACCTTGA
 CCTGACGTCCCCCATGCTGGGGGTTGCATCTGAGCACACCAGACAGCTGCTGCTGGAGGGGCTGTGCGA
 GTGAAGGAGGGACGAGAAGGGAAGCTGGACGTGTACCTGTTCTCTTCTCTGATGTGCTCCTTGTGACCA
 AGCCCCAGCGCAAGGCGGACAAAGCCAAGGTCATCCGACCCCTCTCATGCTGGATAAGCTCTGTGTGCCA
 ACCCTGCGAGACCCCAACAGCTTCTCTGCTGATCCACCTCACTGAATTCCAGTGTGTCTCCAGCGCCCTC
 CTTGTGCACTGTCCAGTCTTACAGACCGTGCCCAAGTGGCTGGAGAAGACCCAGCAGGCCAGGCCGCC
 TACAGAAGCTGAAGGCAGAGGAGTATGTTCAACAGAAGAGGAGCTCCTGACCCTCTATCGGGACCAGGA
 CAGGGAGTCCCCCAGCACCAGGCCCTCCACGCCTTCCCTGGAGGGCTCTCAGAGCAGCGCAGAGGGGAGG
 ACTCCTGAGTTCTCGACCATATATCCCCACCTGGTGGTGACAGAAGACACAGATGAAGATGCTCCCTTG
 TGCCAGATGATACCTCAGACTCTGGCTACGGCACTTTGATCCAGGCACCCCAAGGGGTCCCGCTCCCC
 ACTGAGCCGTCTACGCCAAAGAGCCCTTCGGCGGGACCCTCGCCTCACCTTCTCCACCTGGAACCTCCGG
 GACATCCCTCTGCGTCCCCACCTCCCCGACCCCAAGCTCCCCAACGCCGAAGCGCCCCGAAGTCCCGG
 AAGGAATCTAAAAGGAGGCAGTCTTCCCAGGAAGACCCACCAACCTGGTCTGAGGAAGAAGATGGGGC
 CTCCGAGCGAGGGAATGTGGTGGTGGAAACACTCCACAGGGCCCGGCTTCGGGGCCAGCTTCCCTCTCC
 CCAACCCATGCTGACTCTGCGGGGAAAGCCCTGGGAGTCTCAGGGGAGGAGGAAGAAGAGGGGCCTC
 TGTTCCTGAAAGCTGGCCACACATCCCTGCGCCCAATGCGGGCTGAGGACATGCTCAGAGAGATCCGGGA
 GGAGCTGGCCAGCCAAAGGATTGAGGGGGCCGAGGAGCCCCGGGACAGCAGGCCACGGAAGCTGACTCGG
 GCCAGCTGCAGAGGATGCGGGGGCCCCACATCATTAGCTGGACACCCCTCTGTCCGCATCAGAGGTAT
 GAGGAATGCAGAGGACCTTTGGCATGCATCTCTCCAGAGGAGATCTCTCCCAGTAGTGTGTTCAACCC
 TCCGGCATCTGTGACTCTACCTCAAGGACCACATTTCCCAAAGGAAGCTGGCCCAAGGACCCCTGCCTCC
 TGCTCTGTTTGGGGATCAAGAACTGTAAATTTATGTATCATAGGTGCACCTGAGCCCCACAGAAAGTTGT

GCATAAAATGACTGCCCTGGCTGGGCATGGCTGCCGTGTAATCCCAGCACTTTGGGAGGCTGAGGTGGGA
GGATCCCTTGAGCCCAGGAGTTCCAGACCAGCCTGGGCAATATAGGGAACCCCTGTCTTTACAAAAAA
ATTTTAAAAATT

>gi|29728515|ref|XM_030970.9| Homo sapiens KIAA0720 protein (KIAA0720),
mRNA

AAACTAAGGCCTTCCTGGTAGCCAAGGAGCCAGCAGAGGGGCTGCTGCTGGCTGCTGGGATTGCAGAGG
CCGGGGTCTCTGTGGTGTTCCTTTCCTGGAAAGGAGGACGGCACCCTCCAGGGCAGCCAGCAGTGTGGGA
GCCCATTCCAGGGCGCAAATGGGCACTCAGGAATGGAAAATCCAAGAAGGAACCAACCACCAACAAGGAA
ATGCAGTGTGAGCGGCGGCTGCGAAGTCTGTGCCCTTCAAATATACAGCGCCAGGGGACCTGTGTTGGAC
GGGAGCAAAATTGGCAGGGCTGAGAGCGCCTTGGTTTTCTGTGCAGACAGGTCCAGCTGCAGAAATTAAG
ACTGGGCTGAGTCGGGGAGAAGCCTTCCATGTTGAGATGGCAACTGGGCAAAGCAGGACGGAGACCTGT
GTGTCTGATTAGAGAGGCCTGTGGCATTGGCATGTGGCTGGAGGTCCAGGCAGGGTGCCACCACCGCAGA
GGTCAAGCATGAATTCAGTCTTGACAAAACATGGGTCTCCACCTCGAAGCTGGCTCAGCCTTTGTTTCAGG
AACCGACGACCAGAGCCCCGTGAAAAGAAGGGACTGCGCTGTCAGAACCCCGCTGCATGGACAAGGGG
CGGGCGGCCAAGGTATGTCACCACGCCGACTGCCAGCAGCTGCACCGCCGGGGGCCCTCAACCTCTGCG
AGGCCTGTGACAGCAAGTTCCACAGCACCATGCATTATGATGGGCATGTCCGCTTCGACCTTCCCCACA
AGGCTCTGTGCTGGCCCCGAACGTGTCCACCCGGTCATGCCCGCCGCGCACCAGCCCCGAGTGGACTTG
GAGGAGGAGGAGGAGGAGAGCTCTGTGGATGGCAAAGGGGACCGGAAGAGCACAGGCCTGAAACTCTCCA
AGAAGAAAGCAAGGAGGAGACACACGGATGACCCAAGCAAGGAATGCTTCACCTTGAAATTTGACCTGAA
TGTGGACATTGAGACAGAGATCGTCCCAGCCATGAAGAAGAAGTCACTGGGGGAGGTGCTGCTGCCCTGTA
TTTGAAAGGAAGGGCATTGCGCTGGGCAAAGTGGACATCTACCTGGACCAGTCCAACACACCCCTGTCCC
TCACCTTCGAGGCCTACAGGTTTCGGGGACACTACCTTCGTGTCAAAGCCCCAGCCAAGCCTGGAGATGA
GGGCAAGGTGGAGCAGGGCATGAAGGACTCCAAGTCCCTGAGTTTGCCGATTCTGCGGCCAGCTGGGACC
GGGCCCCCGCCCTGGAGCGTGTGGACGCCCAGAGCCGCCGGGAGAGCCTGGACATCTTGGCCCCCTGGCC
GCCGCCGCAAGAACATGTGCGAGTTCTTGGGGAGGCGAGCATCCCCGGGCAGGAGCCCCCACGCCCTC
CAGCTGCTCTCTGCCCAGCGGCAGTGGCAGCAGCAACACTGGCGACAGCTGGAAGAACCGGGCGGCC
AGTCGCTTCAGCGGCTTTTTAGCTCCGGCCCCAGCACCAGCGCCTTTGGCCGGGAGGTAGACAAGATGG
AGCAGCTGGAGGGCAAGCTGCACACCTACAGCCTCTTCGGGCTGCCCAGGCTGCCCGGGGGCTGCGCTT
CGACCATGACTCCTGGGAGGAGGAGTACGATGAAGACGAGGATGAGGACAATGCCCTGCCTGAGGCTGGAG
GACAGCTGGCGGGAGCTCATTGATGGGCATGAGAAGCTGACCCGGCGGCAGTGCCACCAGCAGGAGGCGG
TGTGGGAGCTGCTGCACACGGAGGCCTCTACATCAGGAACTGCGGGTGATCATCAACCTGTTCTCTGTG
CTGCCCTCTGAACCTGCAAGAGTCAGGGCTGCTGTGTGAGGTGGAGGCGGAGCGCCTGTTTCAGCAACATC
CCGAGATCGCGCAGCTGCACCGCAGGCTGTGGGCTAGCGTGATGGCGCCGGTGCTGGAGAAGGCGCGGC
GCACGCGAGCGCTGCTACAGCCCCGGGACTTCTCAAAGGCTTCAAGATGTTTCGGCTCGCTCTTCAAGCC
CTACATCCGCTACTGCATGGAGGAGGAGGGCTGCATGGAGTACATGCGCGGCCTGCTGCGCGACAACGAC
CTCTTCCGGGCCTACATCACGTGGGCGGAGAAGCACCCACAGTGCCAGAGGCTGAAGCTGAGCGACATGC
TGGCCAAACCCACCAGCGGCTCACCAAGTACCCGCTGCTGCTCAAGTCGGTGCTGAGGAAGACCAGGA
GCCGCGCGCCAAGGAGGCCGTGCTCGCCATGATCGGCTCCGTGGAGCGCTTCATCCACCACGTGAACGCG
TGCATGCGGCAGCGGCAGGAGCGGCAGCGGCTGGCGGCCGTGGTGAGCCGCATCGACGCCTACGAGGTGG
TGGAAAGCAGCAGCGACGAAGTGGACAAGCTCCTGAAGGAATTTCTGCACCTGGACTTGACAGCGCCCAT
CCCTGGCGCCTCCCCGAGGAGACGCGGCAGCTGCTGCTGGAGGGGAGCCTGAGGATGAAGGAGGGGAAG
GACAGCAAGATGGATGTGTACTGCTTCTCTTTCAGCTGCTGTTGTTGGTGACCAAGCAGTGAAGAAGG
CAGAGAGGACCAGGGTCATCAGGCCACCCCTGCTCGTGGAACAAGATTGTGTGCCGGGAGCTACGGGACCC
TGGGTCTCTTCTTATCTACCTGAATGAGTTTACAGTGCTGTAGGGGCCTACACGTTCCAGGCCAGT
GGCCAGGCCTTGTGCCGTGGCTGGGTGGACACCATTTACAATGCCCAGAACCAGCTGCAACAGCTGCGTG
CACAGGAGCCCCCAGGCAGTCAGCAGCCCCCTGCAGAGCCTGGAAGAGGAGGAGGATGAGCAGGAGGAGGA
AGAGGAGGAGGAGGAGGAGGAGGAGGAAGGCAGGACAGTGGCACTTCAGCTGCCAGCTCCCCCTACCATC
ATGCGGAAAAGCAGCGGCAGCCCCGACTCTCAGCACTGTGCCTCAGATGGCTCCACGGAGACCCCTGGCCA
TGGTTGTGGTAGACCTGGGGACACGCTGTCTCCCCGAGTTTCGACAGCGGTCTTTTCAGCTCCCAAGTC
TGATGAGACCTCTCTCAGCACCCTGCTCATCTGCCACGCCCCACAGTGAGCTGCTGCCCCCTGGGTCCG
GTGGACGGCCGCTCTGCTCCATGGACTCTGCCTACGGCACCCCTCTCCCCAACCTCTTACAAGACTTTG
TGGCCCCAGGCCCAATGGCAGAGCTAGTGCCCTCGGGCCCCAGAGTCCCCACGAGTTCTTCCCCCTCCACC
CTCGCCCCGTCTCCGCCGCCGACCCCTGTCCAGCTGTTGAGCTGCCCCCCCCACCTGCTCAAGTCTAAG
TCCGAGGCCAGCCTCTCCAGCTGCTGGCAGGGGCTGGCACCCATGGGACACCCCTCTGCCCCCAGCCGCA
GCCTGTGAGAGCTCTGCTTGGCTGTTCCAGCCCCAGGTATTAGGACTCAGGGCTCCCCCTCAGGAAGCTGG
GCCAGCTGGGATTGCCGAGGGGCCCTAGCCCTGGCAGCGGTCTGGGCTAGTCGGCTGCCTGGCCGGG

GAAGAGAAACATGGCTGCTTTTTCTAAAGAATGTGTGTCTGGGGTTGGAAGATCTGCAGTATGTTTTTCATG
ATTTCTTTCACATGAGCTTTTTATTACATTGTTGAAAGATGAAGAACGAAAGCTACTTGTGTGATCAGATGA
GGAAGAGATCCCCTAGAGTAAATCTGTGCATTAAACCTGTAACCTTCATTTTATGATATCCAGCTTCAGC
AAGTGTCAACATTGGTCAGTTAGAGCATCAACTTATATTGTCACTGGATCCTTGGAGGATTAGACAAATT
TTAATTGAATTACATGGTATGACTTCAGAGCGCCAGTTCTGGACAGTGTCTAATAAGTGGGAAGTACCTT
CTGTCTATAGTGGTGTATCCTGGGAATTAAGACAATTTAACAAGAGATTGGTTTATATTCCTTATGGC
CAAAGGTTTGCAGTCTGCTGTTAAGGACTTTTCCCATGCTAAACAGCTCTTTGCTGCTTGTGGAG
TTGGTAACAGAGTTCTCACCGAAGCTTCGTGAGGTCATGCTGAATGAGATGTTGCTTTTGGATATTCTATA
CACACGAAGCTGGGACAGGGCAGGCAGGAGAGAGACCGCCATCCGACCTTATAAGTAGAGTACGAGGCTA
TCTGGAAATGAGGCTTCCTGATATTCTCTTCGTCAAGTTTATAGCTGAGGAATGTGTTGCCCTTATGTTA
AACTGGAGAGAAAATGAATACCTTACACTCCAAGTCTCTGCATTTTTGCTTCAGAGTAATCCATATGTAA
AGCTTTGGACAGCTTTTACGACTACATGCAAAAGAACTCCAGGCCCTAAAGAAAAGTAGACGGACTGCCAA
AGACCTTTTGCGAAGTTTGTGTTCAAACTCTGTAGTGTGTCCAGTCAGCACAAACGAGGAAATGATGGCAGA
GTTAGTTTAAATAAAACAGAGGGAATCTACGTTAGGTATCATGTATCGGAGTGAAC TGCTTTC TTTTATCA
AAAAATTACGAGAAACCACTCGTTTTGACTATTATTTTATCACTCTTTGCGAAACTTCACAATGTTCCGGGA
GGACATTGTGAATGATATTACAGCTGAACACATTTCTATTTGGCCATCTTCCATTCCCAACCTCCAGTCT
GTGGACTTTGAAGCTGTGGCAATCACAGTGAAAGAGCTAGTTTCGATATACACTCAGTATAAATCCAAATA
ACCATTCTTGTTAATTATCCAGGCAGATATTTACTTTGCAACGAATCAGTATTCAGCAGCTCTTCATTA
TTACCTCCAGGCAGGAGCTGTGTGTTCTGACTTCTTTAACAAGGCTGTGCCCTTGATGTTTATACAGAC
CAGGTAATAAAACGAATGATAAAATGTTGTTCTTTGCTGAATTGCCACACACAGGTGGCCATTTTATGTC
AGTTCCTCAGAGAAATTGACTACAAAAACAGCGTTTAAATCTCTGCAAGAACAAAACAGTCATGATGCTAT
GGACTCCTACTACGACTACATATGGGATGTTACCATTTTGGAACTACTGATTTATCTTCATATAAAAGA
GGAGAAACAGATAAAAGACAAATTGCAATCAAAGCCATCGGCAGACAGTGTGAATGCAAGCAATCCAG
AAGAAGTGTTACAGCTGGCAGCGCAGGAGAAGGAAAAAAGTTTCTCC AAGCAATGGCAAAAC TTTACTT
TTAAGCAGTTAAATTTTTTAACTGTTATTTTAAACAATGGGCTAAAAATAAACAGTATTTAAAGGTT
AAGTTTATATAAAAAAAAAAAAAA

70

CTCTCCCCAAAGTGGCTTATGCAACTGCCATGGACTGGTTTATTGCTGTTTGTATGCATTTGTGTTCTC
 TGCCCTAATTGAATTTGCAACTGTTAATTACTTCACCAAAAGAGGATGGACTTGGGATGGGAAGAGTGTA
 GTAAATGACAAGAAAAAGAAAGGCTTCCGTTATGATACAGAACAACGCTTATGCAGTGGCTGTTGCCA
 ATTATGCCCCGAATCTTTCAAAGATCCAGTTCTCTCCACCATCTCCAAGAGTGCAACCACGCCAGAACC
 CAACAAGAAGCCAGAAAAACAGCCAGCTGAAGCAAAGAAAACTTTCAACAGTGTAGCAAAATTGACAGA
 ATGTCCAGAATAGTTTTTCCAGTTTTGTTTGGTACCTTTAATTTAGTTTACTGGGCTACATATTTAAACA
 GAGAACCTGTATTAGGGGTCAGTCCTTGAATTGAGACCCATGTTATCTTTGGGATGTATAGCAACATTAA
 ATTTGGTTTTGTTTTGCTATGTACAGTCTGACTAATAACTGCTAATTTGTGATCCAACATGTACAGTATGT
 ATATAGTGACATAGCTTACCAGTAGACCTTTAATGAGAGACATGCATTTGCTAACTCATGGAACGTCAGAC
 AGAAAGCACTCCATGCGAAAAACGCCATTGCCTTTTTTAAAGATTTACCCTAGGACCTGATTTAAAGTGA
 ATTTCAAGTACCTTGATTAATTTCTATTCTTCCAAATGAGATGAAAATGGGGATCCTGTACAACCCCTTT
 GTGGACCCCTTTGGTTTTAGCTCTTAAGTAGGGGTATTTTCTACTGTTGCTTAATTATGATGGAAGATAAC
 ATTTGTCATTCTAGATGAATCCTTTGAAGTAACAAACATTGTATCTGACATCAGCTCTGTTTCATGAGTGC
 TCAGAGTCCCTGCTAATGTAATTGGAAGCTTGGTACACATAAGAAAACTAGAGATTTGAAATCTAGCTA
 TGAATTACTCTATATAGTATCTATAGCCATGTACATATTACAGCATGACAAGCTCGAAATAATTATGAGT
 CAGCCCGAAAGATGTTAAT

>gi|6031207|ref|NM_000810.2| Homo sapiens gamma-aminobutyric acid
 (GABA) A receptor, alpha 5 (GABRA5), mRNA

GAAGATGCTGTTGAGGGCCCTGGAGAACTTCAGCAGAACAGGGCCTCTCCCTTGCAGGCCGAGCCGGG
 GCCCTGCGCCCTCCCCCTCCGCCAGCTCGGCCAAGGGCGCATTGTGCTGAGCGTCTGGCGGCCCTCTACCG
 GAGCACCTCTGCAGAGGGCCGATCCTCCAGCCCAGAGACGACATGTGGCGCTCGGGCGAGTGCCCTGCAG
 AGAGAGGAGTAGCTTGCTGGCTTTGAACGCGTGGCGTGGCAGATATTTCAAGAAAGCTTCAAGAACAAAGCT
 GGAGAAGGGAAGAGTTATTCTCCATATTCACCTGCTTCAACTACTATTCTTATTGGGAATGGCAATGG
 AATGTTCTCTGGTTTTATCATGATCAAAAACCTCCTTCTCTTTTGTATTTCCATGAACCTTACCAGTCAC
 TTTGGCTTTTCACAGATGCCAACCAGTTTCAAGTGAAGAGATGAGACCAATGACAACATCAGATATTTACCA
 GGATCTTTGGATGGGCTCTTGGATGGCTACGACAAGACTTTCGGCCCGGGCTGGGAGAGCGCATCACTCA
 GGTGAGGACCGGACCTTACGTCAACAGCTTTCGGCCCGGTGTCCGACACGGAATGGAGTACACCATAGAC
 GTGTTTTTCCGACAAAGCTGGAAAGATGAAAGGCTTTCGGTTTAAAGGGGCCCATGCAGCGCTCCCTCTCA
 ACAACCTCCTTGCCAGCAAGATCTGGACCCCAGACACGTTCTTCCACAAACGGGAAGAAGTCCATCGCTCA
 CAACATGACCACGCCCAACAAGCTGCTGCGGCTGGAGGACGACGGCACCTGCTCTACACCATGCGCTTG
 ACCATCTCTGCAGAGTGCCCCATGCAGCTTGAGGACTTCCCATGGATGCGCACGCTTGCCCTCTGAAAT
 TTGGCAGCTATGCGTACCCTAATTCTGAAGTCGTTTACGTCTGGACCAACGGCTCCACCAAGTCGGTGGT
 GGTGGCGGAAGATGGCTCCAGACTGAACAGTACCACCTGATGGGGCAGACGGTGGGCACCTGAGAATC
 AGCACCAGCACAGGCCAATACACAATCATGACAGCTCACTTCCACCTGAAAAGGAAGATTGGCTACTTTG
 TCATCCAGACCTACCTTCCCTGCATAATGACCGTGATCTTATCACAGGTGTCCTTTTGGCTGAACCGGGA
 ATCAGTCCCAGCCAGGACAGTTTTTGGGGTCAACACGGTGCTGACCATGACGACCTCAGCATCAGCGCC
 AGGAACTCTCTGCCCCAAAGTGGCCTACGCCACCGCCATGGACTGGTTTCATAGCTGTGTGCTATGCCTTCG
 TCTTCTCGGCGCTGATAGAGTTTGCCACGGTCAATTACTTTACCAAGAGAGGCTGGGCCTGGGATGGCAA
 AAAAGCCTTGGAAGCAGCCAAGATCAAGAAAAAGCGTGAAGTCATACTAAATAAGTCAACAAACGCTTTT
 ACAACTGGGAAGATGTCTCACCCCCAAACATTCCGAAGGAACAGACCCAGCAGGGACGTCGAATACAA
 CCTCAGTCTCAGTAAACCCCTCTGAAGAGAAGACTTCTGAAAGCAAAAAGACTTACAACAGTATCAGCAA
 AATTGACAAAATGTCCCGAATCGTATTCCAGCTCTTGTTCGGCACTTTCAACTTAGTTTACTGGGCAACG
 TATTTGAATAGGGAGCCGGTGATAAAAAGGAGCCGCTCTCCAAAATAACCGGCCACACTCCCAAACTCCA
 AGACAGCCATACTTCCAGCGAAATGGTACCAAGGAGAGGTTTTGCTCACAGGGACTCTCCATATGTGAGC
 ACTATCTTTTCAGGAAATTTTGCATGTTTAATAATATGTACAAATAATATTGCCTTGATGTTTCTATATG
 TAACTTCAGATGTTTCCAAGATGTCCCATTGATAATTCGAGCAAACAACCTTCTGGAAAAACAGGATACG
 ATGACTGACACTCAGATGCCAGTATCATACGTTGATAGTTTACAAACAAGATACGTATATTTTAACTG
 CTTCAGTGTTACCTAACAATGTTTTTTATACTTCAAATGTCAATTCATACAAATTTTCCAGTGAATAA
 ATATTTTAGGAACTCTCCATGATTATTAGAAGACCAACTATATTGCGAGAAACAGAGATCATAAAGAGC
 ACGTTTTCCATTATGAGGAACTTGGACATTTATGTACAAAATGAATTGCCTTTGATAATTCTTACTGTT
 CTGAAATTAGGAAAGTACTTGCATGATCTTACACGAAGAAATAGAATAGGCAAACCTTTTATGTAGGCAGA
 TTAATAACAGAAATACATCATATGTTAGATACACAAAATATT

>gi|4504018|ref|NM_000171.1| Homo sapiens glycine receptor, alpha 1
 (startle disease/hyperekplexia, stiff man syndrome) (GLRA1), mRNA
 CGGGAGGCAACAGACACGCTGGAGTTTAACAAACAGCAATACTCTTCGCGCTCCTGAAAAGCAGGTCTGG

ACGCTCTCCGTGGTGTCTGAAACGCCCTCGCAGCCGCCGCTGTCCGTGGTATCTACGACCCCCCTCGCTCCAA
 TTTCCCTGGGGCTCTCCCTCCGCGCCCCCTGTTCCCGCCCTCCCTTTTAACATCTGGATTATTTTTTGCAA
 TAGCGCTTTCTGGTTTTGTAAAGTGCCAATTTGAAACATTTTTTGCCCCCATAACTCGTGGACTACAAAGC
 ACAAAGGACCTGAAAAATGTACAGCTTCAATACTCTTCGACTCTACCTTTTCGGGAGCCATTGTATTCTTC
 AGCCTTGCTGCTTCTAAGGAGGCTGAAGCTGCTCGCTCCGCAACCAAGCCTATGTCACCCCTCGGATTTCC
 TGGATAAGCTAATGGGGAGAACCTCCGGATATGATGCCAGGATCAGGCCCAATTTTAAAGGTCCCCCAGT
 GAACGTGAGCTGCAACATTTTCATCAACAGCTTTGGTTCCATTGCTGAGACAACCATGGACTATAGGGTC
 AACATCTTCTGCGGCAGCAATGGAACGACCCCCGCCCTGGCCCTATAATGAATACCCTGACGACTCTCTGG
 ACCTGGACCCATCCATGCTGGACTCCATCTGGAAACCTGACCTGTTCTTTGCCAACGAGAAGGGGGCCCA
 CTTCCATGAGATACCAACAGACAACAAATTGCTAAGGATCTCCCGGAATGGGAATGTCTCTACAGCATC
 AGAATCACCCTGACACTGGCCCTGCCCATGGACTTGAAGAATTTCCCATGGATGTCCAGACATGTATCA
 TGCAACTGGAAAGCTTTGGATATACGATGAATGACCTCATCTTTGAGTGGCAGGAACAGGGAGCCGTGCA
 GGTAGCAGATGGACTAACTCTGCCCCAGTTTATCTTGAAGGAAGAGAAGGACTTGAGATACTGCACCAAG
 CACTACAACACAGGTAAATTCACCTGCATTGAGGCCGGTTCCACCTGGAGCGGCAGATGGGTTACTACC
 TGATTGAGATGTATATTCCAGCCTGCTCATTGTCTCCTCTCATGGATCTCCTTCTGGATCAACATGGA
 TGCTGCACCTGCTCGTGTGGGCCTAGGCATCACCCTGTGCTCACCATGACCACCCAGAGCTCCGGCTCT
 CGAGCATCTCTGCCCAAGGTGTCTTATGTGAAAGCCATTGACATTTGGATGGCAGTTTGCCTGCTCTTTG
 TGTTCTCAGCCCTATTAGAATATGCTGCCGTTAACTTTGTGTCTCGGCAACATAAGGAGCTGCTCCGATT
 CAGGAGGAAGCGGAGACATCACAAGGAGGATGAAGCTGGAGAAGGCCGCTTTAACTTCTCTGCCTATGGG
 ATGGGGCCAGCCTGTCTACAGGCCAAGGATGGCATCTCAGTCAAGGGCGCCAAACAAGTAACACCACCA
 ACCCCCCCTCCTGCACCATCTAAGTCCCCAGAGGAGATGCGAAAACCTTTCATCCAGAGGGCCAGAAGAT
 CGACAAAATATCCCGCATTGGCTTCCCCATGGCCTTCCTCATTTTCAACATGTTCTACTGGATCATCTAC
 AAGATTGTCCGTAGAGAGGACGTCCACAACAGTGAAGGGTCTGAAAGGTTGGGGGAGGCTGGGAGAGGG
 GAACGTGGGAATAGCACAGGAATCTGAGAGACGGT

>gi|20127456|ref|NM_004130.2| Homo sapiens glycogenin (GYG), mRNA
 CTCTGAGTCACCAACCTGAGGCTGCCCCGGCCGCTGCGCACCCGGCAGCACCATGACAGATCAGGCCTT
 TGTGACACTAACCACAAACGATGCCTACGCCAAAGGTGCCCTGGTCTGGGATCATCTCTGAAACAGCAC
 AGGACCACCAGGAGGCTGGTCTGTGCTCGCCACCCCTCAGGTCTCAGACTCCATGAGAAAAGTTTTAGAGA
 CAGTCTTTGATGAAGTCATCATGGTAGATGTCTTGGACAGTGGCGATTCTGCTCATCTAACCTTAATGAA
 GAGGCCAGAGTTGGGTGTACGCTGACAAAGCTCCACTGCTGGTCTGCTTACACAGTATTCAAAATGTGTA
 TTCATGGATGCAGATACTCTGGTCTTAGCAAATATTGATGATCTTTTTGACAGAGAAGAATTGTGTCAGCAG
 CACCAGACCCAGGGTGGCCTGACTGCTTCAATTCCGGAGTCTTCGTTTATCAGCCTTCAGTTGAAACATA
 CAATCAGCTGTTGCATCTTGCTTCTGAGCAAGGTAGTTTTGATGGTGGGGACCAAGGCATACTGAACACA
 TTTTTTAGCAGCTGGGCAACAACAGATATCAGAAAACACCTGCCGTTTATTTATAACCTAAGCAGCATCT
 CTATATACTCCTACCTCCCGGCATTTAAAGTGTTTTGGTGCAAGTGCCAAAGTTGTGCATTTCTGGGACG
 AGTCAAACCATGGAATTATACTTATGATCCCAAAACAAAAAGTGTCAAAAGTGAGGCCCATGATCCCAAC
 ATGACTCATCCAGAGTTTCTCATCTGTGGTGGAAACATCTTTACCACCAACGTTTACCTCTGCTTCAAC
 AATTTGGCCTTGTCAAAGACACCTGCTCATATGTAAATGTGCTTTCAGACTTGGTCTATACACTGGCTTT
 CTCTTGTGGCTTCTGTAGAAAGGAAGATGTCTCAGGAGCCATATCACATCTGTCCCTTGGGGAGATCCCA
 GCTATGGCACAGCCGTTTGTATCCTCGGAAGAACGGAAGGAACGATGGGAACAGGGCCAGGCTGATTATA
 TGGGAGCAGATTCTTTTGACAACATCAAGAGGAAACTTGACACTTACCTCCAGTAGAAACACTGCATTTT
 TCTGTGAACACATCCACTTCAACAGCCTTGTTTCTGATACTTAGTATCTAGAGCTGGGTTGAGAAAAGTC
 TGTTACAGTTGCTAGAGGTTTTTCATTAAAACTTATCAGATGAGAGGCTTTTTTAGGATAAGAGGTGAGAA
 CTGGGCAAAAGTTGTGAAGCAGCAATTCTGTTATATGGACAGTGTCTGCTTTTTAATCCTATTTAGCTT
 GTTTCAGAAATTCTCACTTTTGTGACTGCCAACATACAAAGTAAGGGAAACTCAAGATATTAAGATGGC
 TGTATCAGTTCTTAAATCTGCAGAGCCTGGTTCAAAATCAGTCACTCCCTTCAGAAGCAGACATGGCAT
 CTGTTCTTGTCTGCTTGTGTTGGTTGTGTACCTTTCACGAGACCTGAATTTTAGAATTGCCAGTGCTGCC
 AGAGTGAGTGAGTGTAATTCTCCTTTCAGGTAAAGATAGGCTATCTCAACACTGCTGAGTGATTTCATAAA
 CATATCAACCAATAGCATTAAACCATTTTATTTCTGTCTTGTAGTGTCTGAAGATGCTCACCAGTTTTCT
 GTGTACAGTAAGGCAGCATGCTAAAATGCTTTTGTTCAGTTCTGGATATTTGAAAATAGCAGTGTGTTCT
 CTGATGGTTACCTGCAGTGGCACCCCTGTACAAAAAATAAAAGACTTATTGGTGTA

>gi|5453673|ref|NM_003918.1| Homo sapiens glycogenin 2 (GYG2), mRNA
 CGCGGGCTTGCGGGCAGGGGCTGCAGGGAGGAGGAGGACCCGCGCCCGGGGGCTGGGCGGAGGCGG
 GGCCGGGCTTCCGGACAGAGGCCAATCGCTGCCCTCGGGGCTCCAGCGCCGGCTCTGGGCGGAGGCAGC
 CAGAGCGCGGAAGAGGCCTGGAAATCCACGCGGATTCCCGGAGACGGCGCTCTGCTCTGCGGGTTCTGTG

GCGAGGAAGTCCACCCACTGCTCCCGGGCGCAGGTCTGCAGGTCCGCGCCCACTGCCCCGCGGCGCCACTG
 ACCATGTGCGGAGACAGAGTTTACCATGGTGCCAGGCTGGTCTCGAAGTCTCTGAGGTCAAGCAATTCAC
 CCACCTCAGCCTCCCAAAGTGTGGAATGACAGTGAAGTATCAGGCTTTTGTACACTAGCCACCAATGA
 CATCTACTGCCAGGGCGCCCTGGTCTTGGGGCAGTCACTGAGGAGACACAGGCTGACGAGGAAGCTGGTG
 GTGTTGATCACTCCTCAGGTGTCCAGCCTGCTCAGGGTCACTCTCGAAGGTGTTTCGATGAAGTCATTG
 AAGTGAATCTAATCGATAGTGGCGACTACATCCACCTGGCCTTTCTGAAGAGACCTGAGCTCGGGCTCAC
 CCTCACCAAGCTTCACTGTTGGACTCTCACTCACTACAGCAAGTGTGTCTTCTGGATGCAGACACTGTG
 GTGCTGTCCAATGTGATGAGCTGTTTGACAGGGGAGAGTTTCTGCGGCCCGGACCCCGGATGGCCGG
 ATTGCTTCAATAGCGGGGTGTTTGTCTTCCAGCCTTCTCTCCACACGCATAAACTCCTGCTACAGCACGC
 CATGGAACACGGCAGCTTTGACGGGGCAGACCAAGGCTTACTGAATAGTTTCTTCAGGAACCTGGTCGACC
 ACAGACATCCACAAGCACCTGCCGTTTATCTATAACTTGAGTAGTAACACGATGTACACTTACAGCCCTG
 CCTTCAAGCAATTCGGTTCCAGTGCAAAGGTCGTCCACTTTTGGGGTCCATGAAACCTTGAACCTACAA
 GTACAATCCACAGAGTGGCTCGGTGTTGGAGCAAGGCTCAGTGTCCAGCAGCCAGCACCAGGCGGCATT
 CTTTATCTCTGGTGGACGGTCTACCAGAACACGTGCTGCCCCCTTATAAAAGCGTCCAAGCGGGGAAG
 CACGCGCGTCTCCTGGTCAACACTTTGCCACAGTGTGTGGGGGGGCGGTGTGCGGATTACAGCCTCTGG
 TGTTGGAGAGCCGTGTGAAAATTCAACACCCAGTGCAGGGCGTCCGTGTGCAAATTCACCACTGGGTCT
 AACCAGCCTGCTCAGGGCCTTCCGGAGCCGACCCAGATAGTGGATGAGACCCTGTCCCTACCTGAAGGAC
 GCCGTTTCAAGATATGATAGCTTGTCTGAAACTGAGACTCCTGCCGTGATAACGTGTGACCCACTGTC
 CCAGCCTTCCCCTCAGCCTGCAGACTTACAGAGACTGAAACCATCTTGCAGCCAGCAAATAAAGTCGAA
 AGTGTCTCATCCGAGGAAACCTTTCGAACCAAGCCAGGAACCTCCCTGCTGAGGCTCTCAGGGACCCAGTC
 TGCAGGATGCACTGGAGGTGACCTGGCCGTCTCTGTTTCCAGATCTCCATCGAAGAGAAGGTGAAGGA
 ATTGAGCCCCGAGGAAGAGAGGAGGAAGTGGGAGGAAGGCCGTATCGACTACATGGGGAAGGACGCGTTT
 GCTCGCATCCAGGAGAAGCTGGACCGGTCTCTGCAGTAATCCGGCAGCTGGTGGGCGTTGTGTGTAGTTA
 GACAATGTCTGTTGGGTGGTCTCTGTTGCGTGGAGATCTCTCTGGTCTTTCAAAGGGAACAGCTGTTG
 AACCTTGTGCTCTATTTATGCTTAATCCATTTGAGTGCCTCACAAAAAACGTAGAGTATAGAAATCC
 ACCTTAAAGCCCCCTCGCCCCAACCTCTCACCAACGCCTTCTGGGCTTTCTTTCAGAGGTCACTTCTACCC
 TTGAAGTGTCTGGCAAAAGCGAGCAGTAATAACATTCTAGTAGACTCTCGATGGTGGTCTCCGCTCTTGC
 CCGAAGGACCTCTGAAGTACGCTGGATCTGTGTTGTACAGGTGCTGTGAGACCTACCCTATTGAGAAATTA
 AACCTCACTGCAAATTTCTCCCATCACGAAGCTAACAACACTAATATACGTATTTAGCACCTCTGAGGC
 TTTGCCATGGAGACCATTCTGTAGGGCTAAGGAAACATTTAGACGTGGTGAAGTACTTTTCAATTTGGACT
 TGGCGAAGTGTATCTGAGAAACACCTCGGCTGTGGTCTCTCTGCTTTAAATCCTAACAGGACTTCTTAGA
 GCGTTGACAGAAATTTCTACTCGTGGACGTTGGGAAGAAAGATTGTAGGTGGCTTGGGGAATGTGGGTGGC
 TTAGAGGATCTAAACCGATTCACTTCTGGTTGAGAAGCAACGAGGGCTTGCTCTAAATCTGTTTAGAGGA
 TAACAGGATCTAGAGATGCTCTCTGCTTGACAACAAAAGTCAGGGTGCAGTCGGTCCACCCTTGACTGCT
 CTTGGCTTGGTCTCTACCTCACTACCTCAGTTCTCAATAACTTAGTGAATCACTGCCCTCCTCAAAGCC
 ATTTCCACTCAGCTCTTTCCAGAGAATTCTCAGTTTTATGAGACGGGAACTTTATTTACAGAGAAAGCC
 TCATTGTGAGAAGTATCTTCAATTCAATGGGCACAATATGCTGTGTATCTCACCAGGTAGCTGTGAGGGC
 CACCGAGAGTGTGCTTAAAAATGGGCATCGTTGTAATAAAGGAGGAAAGTGCAGCTTTTGAAATGTTTGG
 AAGGTTTATTTCTCATGCACATTCCAGGGAAGAGCAGAGAGTAAATTAGAGACGGGATAGGAAGGCCGTG
 GGAGAACTCGATCCTAGCCTGTGTGACGCTGGATGTGTTTACGTGGAGAGGCGTGCCACTTTTATGTTCA
 CCTGAAGCAGTTTAGCCTTTGGATAGAGGAACCTGCCTGAATTTATGGCATTAGTGGTGGCATTGTTTTTG
 TGTACAAGATGTGGGTGATGGAGGGGCTGTTCTTTTTCCGTGTGGGTGGTTAATAATCGTCAAGTCTCGG
 AGGGCGATGCTCGTAGGATATTTCAAGGTGAGTCAGGGTTGGATGGTCATCGGCTTTTCAGAGGGAGACCAC
 GGAATGTTTCAGGGAAACAATGTCTAGCTTCTCTGAGGACCAGAATTC

>gi|30147854|ref|XM_301113.1| Homo sapiens similar to Glycogenin-1
 (LOC350705), mRNA

ATGACAGATCAGGTCTTTGTGACACTGACCACAAATGATGCCTACACCAAAGGTGCCCTGGTCTCTGGGCT
 CATCTCTGAAACAGCACAGGACCACCAGAAGACTAGTCATGCTCGCCACCCCGAGGTCTCAGACTCCAT
 GAGAAAAGTTTTAGAGACAGTCTTTGATGAAATCATCGTGGTAGATGTCTTGGACAGTGGTGATTCTGCT
 CATCTAACCTTAATGAAGAGGCCAGAGTTGGCATTAAAGTGTGGTGGCAAGTGCCAAAGTTGTGCATT
 TCCTGGGACAAGTCAAACCATGGAATTATACTTATGATCCCAAAACAAAAGTGTCAAATGAGTCCCA
 CCATCCCAACGTGACTCATCCAGAGTTTCTCATCTATGGTGGAACTCTTTACCACCACCGTTTTTACCT
 CTGCTTCAACAATTTGGCCTTGTCAAAGACACCTTCTCATACGTAAATGTGGAAAATGTCTCAGGAGCCA
 TATCACATCTGTCCCTTGGGGGGATCCAGCTATGGCACAGTCTGTGTATCTCAGAAGAATGGAAGGAG
 CGGTGGGAACAGGGCTAGACCGATTATATGGGAGCAGATTCTTTGACAACATCAAGAGGAAGCTTGACA
 CTTACCTCCAGTAGAAACACTGCATTTTCTGTGGACACATCCACTTCAAGCCTTGTTTCTGATACTT

AG

>gi|4757725|ref|NM_004641.1| Homo sapiens myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 10 (MLLT10), mRNA

GCCCTCTTGATTATGTGTGCCCTCTCCGGGCGCCCGCGTTAGCGGCCGGGTGGAGGTGGGGAGGGAAGAC
GCTGAGGAGGAGGAGGAGGCGGAGGAGCGGTGGAGGGGAGGTGGGGGGAATCAGCAAGGACATGGCTCC
TGACTCCTGTGCGGAACGTGAGTGACTGAGCGGCAAGCCCCGAATGGTCTCTAGCGACCGGCCCGTGTCA
CTGGAGGACGAGGTCTCCATAGTATGAAGGAGATGATTGGAGGCTGTTGCGTTTGCTCAGACGAGAGAG
GCTGGGCGGAGAACCCGCTGGTTTATTGCGACGGGCACGGCTGCAGCGTCGCGGTGCATCAAGCTTGCTA
TGGCATTGTTCAAGTACCCACTGGACCGTGGTTTTGTCAGGAAATGTGAATCTCAGGAGAGAGCAGCCAGA
GTGAGATGTGAACTTTGTCCCCATAAGGATGGAGCTTTAAAAAGAACAGATAATGGGGGTGGGCCCATG
TGGTTTGTGCCCTGTATATTCCAGAGGTACAATTTGCCAATGTTTCCACAATGGAACCAATTGTTTTACA
GTCTGTTCCGCATGATCGTTATAATAAGACTTGCTACATTTGTGATGAACAAGGAAGAGAAAGCAAAGCA
GCCACTGGTGCTTGCATGACATGTAATAAACATGGATGTCGACAGGCTTTTCATGTAACATGCGCTCAGT
TTGCCGGACTGCTTTGTGAAGAAGAAGGTAATGGTGCCGATAATGTCCAATACTGTGGCTACTGTAAATA
CCATTTTAGTAAGCTGAAAAAGAGCAAACGGGGATCTAATAGGTCATATGATCAAAGTTTAAGTGATTCT
TCCTCTCACTCTCAGGATAAACATCATGAGAAAGAGAAAAAAATATAAAGAGAAGGACAAACACAAAC
AGAAACACAAGAAGCAGCCAGAACCATCACCTGCATTGGTTCCATCCTTGACTGTTACTACAGAAAAAC
TTATACAAGCACTAGCAACAACCTCTATATCTGGATCATTGAAGCGCTTGAAGATACTACTGCACGATTT
ACAAATGCAAAATTTCCAGGAAGTCTCTGCACACACCTCTAGTGGAAGATGTTTCAGAGACTAGAGGGT
CAGAGGGCAAAGGGAAGAAATCTTCAGCTCACAGCTCAGGTCAAAGGGGAAGAAAGCCTGGTGGTGGAAG
AAATCCAGGAACAACCTGTGTGTCAGCAGCTAGCCCTTTTCTCAAGGCAGTTTTTCAGGAACCTCAGGCAGT
GTAAAGTCATCTTCTGGAAGTTCACTGCCAGCTCAGTCAAGGATTTCTGAGCTTTACAGACTCAGATCTGC
GTAATGACAGTTACTCTCACTCCCAACAGCTCATGAGAACCAAGATGTACATAAAGGAGAGCTGGAAG
CCAGGAAGGGGGGTAAATAGTTTTAGTACCTTAATTGGCCTCCCTTCAACCTCAGCTGTTACTTTCACAG
CCTAAAGCTTTGAAAATTCACCTGGAGATTTGGGTAATTCCAGCCTTCTTACAGCAGGATATAAGCGGG
CTCAAACTTCTGGCATAGAAGAAGAACTGTAAAGGAAAAGAAAAGGAAAGGAAATAAACAAAGTAAGCA
TGGGCTTGGCAGACCCAAAGGAAACAAAATCAAGAGAATGTTTCTCATCTCTCAGTTTTCTTGCTTCA
CCAACATCATCTGTAGCATCAGCTGCAGGAAGCATAACAAGCTCTAGTCTGCAGAAATCTCCTACATTGC
TCAGGAATGGAAGTTTACAGAGCCTCAGTGTGGCTCATCTCCAGTTGGTTGAGAAATTTCCATGCAGTA
TCGGCATGATGGAGCTTGGCCAACTACGTTCTCAGAGTTGCTGAATGCAATACACAACGACAGAGGT
GACAGTTCTACACTAACAAAGCAAGAACCTAAATTCATAGGTATTTATAACAGCAATGATGTAGCAGTAT
CGTTTTCCAAATGTAGTATCTGGCTCGGGATCTAGTACTCCTGTCTCCAGCTCTCACTTACCTCAGCAGTC
TTCGGGCATTTGCAACAAGTAGGAGCGCTCTCTCCCTCAGCTGTGTCTATCTGCAGCCCCCTGCTGTTGCT
ACAACTCAGGCAAAATACTCTATCTGGATCTTCTCTCAGTCAGGCACCATCTCATATGTATGGCAATAGAT
CAAATTCATCAATGGCAGCTCTTATAGCTCAGTCTGAAAACAATCAAACAGATCAAGATCTTGGAGACAA
TAGCCGCAACCTAGTTGGCAGAGGAAGCTCACCCCGAGGAAGTCTCTGCCACGATCCCTGTAAAGCAGC
TTACAGATTCGCTATGATCAACCAGGCAACAGCAGTTTGGAAAATCTGCCCTCAGTAGCAGCCAGCATAG
AACAGCTTTTGGAGAGGCAGTGGAGTGAAGGACAGCAATTTTAC TAGAACAGGGTACTCCTAGTGACAT
TTTAGGAATGCTGAAGTCATTACACCAACTTCAAGTTGAAAACCGAAGATTAGAGGAACAATTAATAAAC
TTGACTGCCAAAAGGAACGGCTTCAGTTATTGAATGCACAGCTTTTCAAGTGCCTTTTTCCAACAATAACAG
CAAATCCTAGTCCGCTCTCATCAAATACACACATTTTTCAGCACAGACTGCTCCTACTACTGATTCTTTGAA
CAGCAGTAAGAGCCCTCATATAGGAAACAGCTTTTTTACCTGATAATTTCTCTTCTGTATTAAATCAGGAC
TTAACCTCCAGTGAGCAAGTACCAGCAGCTCATCAGCTCTTTCTACCCACCTCCTGCTGGGCAGAGTC
CGGCTCAACAAGGCTCAGGAGTGAGTGGAGTTTACAGAGGTCAATGGCGTGACAGTGGGGGCACTAGCTAG
TGGAATGCAGCCTGTAACCTCCACCTTCTGCGGTGCTGTCAGTGGGTGGAATAATTGGAGCTTTGCCA
GGTAACCAACTGGCAATTAATGGCATTGTAGGAGCTTTAAATGGGGTTATGCAGACTCCTGTCACAATGT
CCCAGAACCCTACCCCTCTCACCCACACAACCGTACCACCTAATGCAACACATCCAATGCCAGCTACACT
GACTAACAGTGCTCAGGACTAGGATTACTTTCTGACCAGCAACGACAAATACTTATTCATCAACAGCAG
TTTCAGCAGTTGTTAAATTTCTCAACAGCTCACACCAGTACACAGGCACCCCCACTTCACACAGCTACCAC
CAACCCATTTCTCACCATCCATGGAGATAATGCAAGTCAGAAAGTAGCAAGACTTAGTGATAAACTGGG
CCTGTAGCTCAAGAGAAAAGTTGACACCTGAGAAACATCTAGAAATTGCCTATCCTGCTGTTCTAGCACT
TCATCTGGCTGCCTTTGTCAGTCTTTTACTACAGCTATGAAGAAACGCAACAAGAACTCAATGCACAAC
AAAGGATTAATTGCTGCAAGGACATTCTTGTAAGGCTTTGATTAGTTTCTTGTGCTTTGTTGCACTGA
AATGGAATTCCTATGCCCTTACCCCTTACCCAGTTTTTTGAACATGGAAGAAAATTTAATAACTTTTT
AAAGTGACATAATTTACATGCAATATGTTTATCAACTCAAGAATTTAATATAGTTGGTACACAAC TAGTT

TTGTTTATAAATTGGAGATGCAAAATAGCAAACTAAATACTTGCTCCATTTACAACTACTTGATTTTAT
TGTACAAGTTGAAATATGCTCTTTTGTGGGTTACAGTATGCTTGCTCTAAGTCAAATTCGAAGAACT
AATTTCTTCTCCTGGAGTTGCATTGATTAGTATTACAAATATATAGCACATCACCTGGGAC

>gi|5174576|ref|NM_005937.1| Homo sapiens myeloid/lymphoid or mixed-
lineage leukemia (trithorax homolog, Drosophila); translocated to, 6
(MLLT6), mRNA

ATGAAGGAGATGGTAGGAGGCTGCTGCGTATGTTTCGACGAGAGGGGCTGGGCCGAGAACCCGCTGGTCT
ACTGCGATGGGCACGCGTGCAGCGTGGCCGTCCACCAAGCTTGCTATGGCATCGTTTCAGGTGCCAACGGG
ACCCTGGTTCTGCGGAAATGTGAATCTCAGGAGCGAGCAGCCAGGGTGAGGTGTGAGCTGTGCCACAC
AAAGACGGGGCATTGAAGAGGACTGATAATGGAGGCTGGGCACACGTGGTGTGTGCCCTCTACATCCCCG
AGGTGCAATTTGCCAACGTGCTCACCATGGAGCCCATCGTGCTGCAGTACGTGCCTCATGATCGCTTCAA
CAAGACCTGTTACATCTGCGAGGAGACGGGCCGGGAGAGCAAGGCGGCCTCGGGAGCCTGCATGACCTGT
AACCGCCATGGATGTGACAAGCTTTCCACGTACCTGTGCCCAAATGGCAGGCTTGCTGTGTGAGGAAG
AAGTGCTGGAGGTGGACAACGTCAAGTACTGCGGCTACTGCAAATACCACTTCAGCAAGATGAAGACATC
CCGGCACAGCAGCGGGGGAGGCGGAGGAGGCGCTGGAGGAGGAGGTGGCAGCATGGGGGGAGGTGGCAGT
GGTTTCATCTCTGGGAGGAGAAGCCGGTCAGCCTCACCATCCACGCAGCAGGAGAAGCACCCACCCACC
ACGAGAGGGGCCAGAAGAAGAGTCAAGAGACAAAGAACGCCTTAAGCAGAAGCACAAGAAGCGGCCTGA
GTCGCCCCCAGCATCCTCACCCCGCCCGTGGTCCCCACTGCTGACAAGGTCTCCTCCTCGGCTTCTCT
TCTTCCCACACGAGGCCAGCACGCAGGAGACCTCTGAGAGCAGCAGGGAGTCAAAGGGGAAAAAGTCTT
CCAGCCATAGCCTGAGTCATAAAGGGAAGAACTGAGCAGTGGGAAAGGTGTGAGCAGTTTTACCTCCGC
CTCCTCTTCTTCTCCTCCTCTTCTCCTCCTCTGCGGGGGCCCTTCCAGCCTGCAGTCTCGTCCCTGCAG
AGCTCCCCTGACTTCTCTGCATTCCCCAAGCTGGAGCAGCCAGAGGAGGACAAGTACTCCAAGCCACAG
CCCCCGCCCCCTCAGCCCCCTCCTTCTCCTCAGTCCCCAGCCCCCAAGGCTGACCTTTTTTGAGCAGAA
GGTGGTCTTCTCTGGCTTTGGGCCCATCATGCGCTTCTCCACCACCTCCAGCTCAGGCCGGGCCCGG
GCGCCTCCCCTGCGGACTATAAGTCTCCCCACGTACGGGGTCTGGGGCCTCGGCAGGCACCCACAAAC
GGATGCCCCGCACTGAGTGCCACCCCTGTGCCTGCTGATGAGACCCCTGAGACAGGCCTGAAGGAGAAGAA
GCACAAAGCCAGCAAGAGGAGCCGCCATGGGCCAGGCCGTCCCAAGGGCAGCCGGAACAAGGAGGGCACT
GGGGGCCAGCTGCCCCATCCTTGCCAGTGCCAGCTGGCTGGCTTTACCGCCACTGCTGCCTCACCT
TCTCTGGAGGTTCCCTGGTCAGCTCCGGCCTGGGAGGTCTGTCTCCCGAACCTTTGGGCCTTCTGGGAG
CTTGCCAGCTTGAGCCTGGAGTCCCCCTTACTAGGGGCAGGCATCTACACCAGTAATAAGGACCCCATC
TCCACAGTGGCGGGATGCTGCGGGCTGTCTGCAGCACCCCTCTCTCCTCCAGCCTCCTGGGGCCCCCAG
GGACCTCGGCCCTGCCCCGCTCAGCCGCTCCCCGTTACCAGCACCTCCCTCCTCTTCTGCTTCTAT
CTCCACCCTCAGGTGTTTTCTCTGGCTGGCTCTACCTTTAGCCTCCCTTCTACCCACATCTTTGGAACC
CCCATGGGTGCCGTTAATCCCCCTCCTCTCCCAAGCTGAGAGCAGCCACACAGAGCCAGACCTGGAGGACT
GCAGCTTCCGGTGTCGGGGGACCTCCCCCTCAGGAGAGTCTGTCTTCCATGTCCCCCATCAGCAGCCTCCC
CGCACTCTTCGACCAGACAGCCTCTGCACCCTGTGGGGGCGGCCAGTTAGACCCGGCGGCCCCAGGGACG
ACTAACATGGAGCAGCTTCTGGAGAAGCAGGGCGACGGGGAGGCCGGCGTCAACATCGTGGAGATGCTGA
AGGCGCTGCACGCGCTGCAGAAGGAGAACCAGCGGCTGCAAGAGCAGATCCTGAGCCTGACGGCCAAAAA
GGAGCGGCTGCAGATTCTCAACGTGCAGCTCTCTGTGCCCTTCCCTGCCCTGCCTGCTGCCCTGCCTGCC
GCCAACGGCCCTGTCCCTGGGCCCTATGGCCTGCCTCCCCAAGCCGGGAGCAGCGACTCCTTGAGCACA
GCAAGAGCCCTCCGGGAAAGAGCAGCCTCGGCCTGGACAACCTCGCTGTCCACTTCTTCTGAGGAGCCACA
CTCAGGCTGCCCGAGCCGAGCAGCTCGTCTGTCTTCCACAGCAGCCCCCACCCTGCCCCCTCCTC
CAGCAGAGCCCTGCCACTCTGCCCCCTGGCCCTGCCTGGGGCCCCCTGCCCCACTCCCGCCCCAGCCGAGA
ACGGGTGGGGCCGGGCACCCGGGGCAGCGGGGCTGGGGGCCATGCCATGGCTGAGGGGCTGTTGGGGGG
GCTGGCAGGCAGTGGGGGCTGCCCCCAATGGGCTCCTTGGGGGGTTGAATGGGGCCGCTGCCCCAAC
CCCGCAAGCTTGAGCCAGGCTGGCGGGGCCCCACGCTGCAGCTGCCAGGCTGTCTCAACAGCCTTACAG
AGCAGCAGAGACATCTCCTTACAGCAGCAAGAGCAGCAGCTCCAGCAACTCCAGCAGCTCCTGGCCTCCCC
GCAGCTGACCCCGGAACACCAGACTGTTGTCTACCAGATGATCCAGCAGATCCAGCAGAAACGGGAGCTG
CAGCGTCTGCAGATGGCTGGGGGCTCCAGCTGCCCATGGCCAGCCTGCTGGCAGGAAGCTCCACCCCGC
TGCTGTCTGCGGGTACCCCTGGCCTGCTGCCACAGCGTCTGCTCCACCCCTGCTGCCCGCTGGAGCCCT
AGTGGCTCCCTCGCTTGGCAACAACACAAGTCTCATGGCCGAGCAGCTGCAGCTGCAGCAGTAGCAGCA
GCAGGCGGACCTCCAGTCTTCACTGCCCAGACCAACCCCTTCTCAGCCTGTGGGAGCAGAGGGCAGTG
GCGGTGGCCCCAAAGGAGGGACCGCTGACAAAGGAGCCTCAGCCAACCAGGAAAAAGGCTAA

>gi|4505956|ref|NM_002697.1| Homo sapiens POU domain, class 2,
transcription factor 1 (POU2F1), mRNA

CGTCAACATGAGTTGGGCCTGGGGCAGATGAGGCTGGCTGGCGGGGCGGGCAGCATGGTTCACTCCAGCA
TGGGGGCTCCAGAAATAAGAATGTCTAAGCCCCTGGAGGCCGAGAAGCAAGGTCTGGACTCCCCATCAGA
GCACACAGACACCGAAAGAAATGGACCAGACACTAATCATCAGAACCCCCAAAATAAGACCTCCCCATTCTCCGTGTCCCAACTGGCCCCAGTACAAAGATCAAGGCTGAAGACCCCAGTGGCGATTTCAGCCCCAGCAG
CACCCCTGCCCCCTCAGCCGGCCCAGCCTCATCTGCCCCAGGCCCAACTCATGTTGACGGGCAGCCAGCT
AGCTGGGGACATACAGCAGCTCCTCCAGCTCCAGCAGCTGGTGCTTGTGCCAGGCCACCACCTCCAGCCA
CCTGCTCAGTTCTCTGCTACCGCAGGCCCAGCAGAGCCAGCCAGGCCTGCTACCGACACCAAATCTATTCC
AGCTACCTCAGCAAACCCAGGGAGCTCTTCTGACCTCCCAGCCCCGGGGCCGGGCTTCCACACAGCCCCC
CAAATGCTTTGGAGCCACCATCCCACCCCCGAGGAGCCAGTGATCTGGAGGAGCTGGAGCAATTGCCCCGC
ACCTTCAAGCAACGCCGCATCAAGCTGGGCTTCACGCAGGGTGATGTGGGCCTGGCCATGGGCAAGCTCT
ACGGCAACGACTTCAGCCAGACGACCAATTTCCCGCTTCGAGGCCCTCAACCTGAGCTTCAAGAACATGTG
CAAACCTCAAGCCCCCTCTTGAGAGAGTGGCTCAACGATGCAGAGACTATGTCTGTGGACTCAAGCCTGCC
AGCCCCAACCAGCTGAGCAGCCCCAGCCTGGGTTTCGACGGCCTGCCGGCCGGAGACGCAAGAAGAGGA
CCAGCATCGAGACAAACGTCCGCTTCGCCTTAGAGAAAGATTTTCTAGCGAACAGAAGCCTACCTCAGA
GGAGATCCTGTGATCGCCGACAGCTGCATCGAGAGGAAAGTAGTACCGCGTCTGGTTCTGCAACCCG
CGCCAGAAGGAGAAACGCATGACACCCCTGCAGTGGCGGCCCATGTGCCCCAGCCAGCCGAAGCCGCGCA
GCTACAGCCCCCATATGGTACACCCCCAAGGGGGCGCGGGGACCTTACCGTTGTCCCAAGCTTCCAGCAG

TCTGAGCACAAACAGTTACTACCTTATCCTCAGCTGTGGGGACGCTCCACCCCAGCCGGACAGCTGGAGGG
 GGTGGGGGGCGGGGGCGGGGCTGCGCCCCCCTCAATTCCATCCCCCTCTGTCACTCCCCCACCCTCGGCCA
 CCACCAACAGCACAAACCCAGCCCTCAAGGCAGCCACTCGGCTATCGGCTTGTCAAGGCTGAACCCAG
 CACGGGGCCCTGGCCTCTGGTGGAACCTGCCCCCTTACCAGCCTTGATGGCAGCGGGAATCTGGTGCTGGG
 GGCAGCCGGTGCAGCCCCGGGGAGCCCTGGCCTGGTGACCTCGCCGCTCTTCTTGAATCATGCTGGGCTG
 CCCCTGCTCAGCACCCCGCCTGGTGCTGGGCTGGTCTCAGCAGCGGCTGCGGCTGTGGCAGCCTCCATCT
 CCAGCAAGTCTCCTGGCCTCTCCTCCTCATCCTCTTCATCCTCATCCTCCTCCTCCTCCACTTGCAGCGA
 GACGGCAGCACAGACCCCTGGAGGTCCAGGGGGGCCCCGAGGCAGGGTCCAAACCTGAGTGAGGGCCAGCC
 ATGCCCTCCCCCTCCCATTCTCTGGTCCCTGCCTTGGTCCCTTGCCTGGGAAGAGGGGCAGGAGGCCAGTG
 GTGGGGAGCGCAGAGGGTCTCAGAGCAGGAGTGACAAGGGAGGAAAGACCAAAAAACAACCAACCAAAA
 AAAAAAAAAAAAAAGGAAAGAACTAACCAACAAAAAGAGAAACCAAAAAATAATCACAAACAGAAACCAGC
 TGCCCCAAAGGAACAGAGGTGAAAAACAAACAAAAAAACCAAAAACAAACCAAAAAAAAAAAAAAAC
 TCTACCCCTCTAGAGCC

>gi|7657408|ref|NM_014352.1| Homo sapiens POU domain, class 2,
 transcription factor 3 (POU2F3), mRNA

GGGGAGGATGGTGAATCTGGAGTCCATGCACACAGATATCAAGATGAGTGGGGATGTAGCCGATTCCACG
 GATGCTCGCAGCACTCTCAGCCAGGTGGAGCCAGGAAATGATCGAAAAGGCCTAGATTTCAACAGGCAGA
 TTAAAACCGAAGATCTCAGTGACTCCCTGCAGCAGACCCCTCTCCCATCGGCCATGCCACCTGAGTCAAGG
 ACCTGCCATGATGTCCGGAACCAAAATGTCTGGGCTAAATGCCAGCCCATGTCAAGGACATGGCTTCCCTC
 CATCCGCTCCAGCAGCTTGTGCTGGTTCCTCGGCCACTTACAGTCTGTATCCAGTTCTGCTATCTCAGA
 CCCAGCCTGGGCAGCAAGGTCTGCAGCCAAATCTCCTCCCTTTCCACAGCAACAAAGCGGTCTCCTCCT
 CCCACAGACTGGGCCGGGACTGGCATCCCAGGCATTTGGGCACCCTGGGCTGCCAGGATCCTCTTTAGAA
 CCCACCTGGAAGCATCCAGCATCTCCCAGTGCCCAAGCATCTACCAGCTCTGGAGGGCCGATGGAGC
 CCAGTGACCTCGAGGAGCTGGAGAAGTTTGCCAAGACCTTCAAGCAGAGGCGCATTAAGCTGGGCTTCAC
 ACAGGGAGATGTGGGCTGGCGATGGGAAAGCTTATGGCAACGACTTCAGCCAGACCACCATCTCACGA
 TTTGAGGCCCTCAACCTGAGCTTCAAGAACATGTGCAAGCTCAAGCCCCCTGCTGGAGAAGTGGCTGAATG
 ATGCAGAGTCTCTCCGTACAGCCCTCAGTGAGCACGCCAGCTCCTACCCAGCCTCAGTGAAGTATT
 TGGTAGGAAGAGAAAGAAACGGACCAGCATCGAGACCAACATCCGCTGACTCTGGAGAAGAGGTTTCAA
 GATAACCCAAAACCCAGCTCGGAGGAGATCTCCATGATTGCAGAGCAGTTGTCCATGGAGAAGGAGGTGG
 TGAGGGTCTGGTTCTGCAACCGACGCCAAAAGGAGAAGCGAATCAACTGCCCTGTGGCCACACCCATCAA
 ACCACCTGTCTACAACCTCCGGCTGGTATCTCCCTCAGGGTCTCTGGGCCCCCTCTCTGTCCCTCCTGTC
 CACAGTACCATGCCTGGAACAGTAACGTCATCTGTTCCCTGGGAACAACAGCAGGCCTTCATCTCCTG
 GCTCAGGACTCCACGCCAGCAGCCCCACTGCATCTCAAATAACTCCAAAGCAGCAGTGAACCTCCGCCTC
 CAGTTTTAACTCTTCAGGATCTTGGTACCGATGGAATCATTCCACCTACCTCCACTGAGACCAAAAAGTT
 TCTCCTACTCCAGCTGGCCCTGTATTCCCCTGGAAGGAAGGGAATCATGCCTTCTATATACAGACAGATT
 GCCTTCAGAAGAGTGGAAGAAAATCTCCACTATCAATGAACCCAGACTCTTGTCTTCTTCAAGAGCAAGG
 GCCTCCGGAGATCCAAACTGTGATTGAACCAAGTGCAGACTCCTAATGCTCTTGAAATACACAGCCCCCTC
 CTAGGAGCTTACCATTTTACCTTCCCTTGCCCTATGCCCTTGCCCTTCTAGTTCCAAATATTTTAGCCAGCT
 TCACTGTGGCAATAGTCTTTCAGAGAAAAGACTTCTTGCTGTTATTCTCCAACCTATCCGCTGGGCTTCTG
 GGGACAGCCATTTGGCTGGGGTGCCAAACACCAGAAGGGAGATAATAGTTTTGACTCTGAACCTTGCCCA
 CAACCCCTGAACCTGATCCCAAAATCTGTGAAAAGATTGAATCTGATATCTCCACCAAGCCTTGATGTT
 TTCTCTGTACAGCTAAGTTTTCTGATGGAATCTTCATCTACCCCATTTTCTTTTTTAACCTCGCCCCCTT
 TTCTACAATCAAATCCATTATTATTGCGCCCTCCAGGTCCCCCTCCTTTTTTGCAAGAGGTGAAAAGAGC
 TGCCCTTGGTGCGTCCCTTGAGCATGTTGCAACGCCTGCTGGTGCCCTTCTGTCTGTCCACGTCTGCCT
 AGGGCTCAGGAGTGCTTGTCTTTCCCTGGCGCCCTGGGTTTCTCTTTATCCTGTCCCTCTCCTTTCTC
 CTGGCAGCAGAAAAGGAAGAACAAGAGCAACACACAACAATGATGGTGGCAAAGTACCTGTATATAG
 CCCTTTCCCATTTTTGCATTATGTTCTTAACTCCTTAATGTGGAATTTTCCAGCAAAATGTTTAACTCAG
 GTGTGATTTTTGAAAATGTCCTGTGAATATTTTATCTTCTTTAAAAAAAGGGGGGAAATACCAAGTG
 TGAAATACTGTGCTACCTTCCAGAGTTCACATGGGCTGGGATTTTTTTTTTCTTTTTTTTTTTTTT
 TTTTTTTGAGACAGTCTTGCTCCAGCCAGGCTGGAGTGCAAGTGGCATGATCTTGGCTCACTGCATGCTC
 TGCCACTTGGGTTCAAGCGATTCTCCTGCCTCAGCCTCTCAAGTAGCGGATTACAGGCGCCACCACCAA
 AAATTAGCCAGCTAATTTTTGTATTTTGTAGAGATGGGGTTTACCAGTGTGGTCAAGGCTGGTCTTG
 AACTCCTGACCTCCTGATCCACCCACCTCAGCGTCCCGAAGTGCTGGGATTACAGGTGTGAGCCACCGCG
 CCTGGCCTGGGCTGGGAATCTTTGTTGACTTCTTTGGGAACATTATCTCTAGAGTCTGCTTTTTTGTGGG
 CCAGTTTGTAGAGATGTAACATGAAGCCATTCCAGAAATGGTAACCTTAAAAAAAAGACTACATGTCT
 AGGAAACTGTTGAGATTCAAAGGAATCTCACTGTTCTAAATAAATGTGTATAGTGATTATGAAAAGT

>gi|31543438|ref|NM_002740.2| Homo sapiens protein kinase C, iota (PRKCI), mRNA

AGCGGTTTTGGGCCCCGGCGGCTGTAGAGGCGGCGGCCTACGGGCAGTGGGAGGAGCCGCGCGGTTCC
GGCTGCTCCGGCGAGGCGACCTTGGGTGCGGCGCTGCGGGCGAGGTGGGCAGGTAGGTGGGCGGACGGCC
GCGGTTCTCCGGCAAGCGCAGGCGGCGGAGTCCCCACGGCGCCCGAAGCGCCCCCGCACCCCCGGCC
TCCAGCGTTGAGGCGGGGAGTGAGGAGATGCCGACCCAGAGGGACAGCAGCACCATTGTCCACACGGTC
GCAGGCGGCGGCGAGCGGGGACCATTCCCACAGGTCCGGGTGAAAGCCTACTACCGCGGGGATATCATGA
TAACACATTTTGAACCTTCCATCTCCTTTGAGGGCCTTTGCAATGAGGTTTCGAGACATGTGTTCTTTTGA
CAACGAACAGCTCTTACCATGAAATGGATAGATGAGGAAGGAGACCCGTGTACAGTATCATCTCAGTTG
GAGTTAGAAGAAGCCTTTAGACTTTATGAGCTAAACAAGGATTCTGAACTCTTGATTTCATGTGTTCCCTT
GTGTACCAGAACGTCCTGGGATGCCTTGTCAGGAGAAGATAAATCCATCTACCGTAGAGGTGCACGCCG
CTGGAGAAAGCTTTATTGTGCCAATGGCCACACTTTCCAAGCCAAGCGTTTCAACAGGCGTGCTCCTACTGT
GCCATCTGCACAGACCGAATATGGGGACTTGAGCGCCAAGGATATAAGTGCATCAACTGCAAACTCTTGG
TTCATAAGAAGTGCCATAAACTCGTCACAATTGAATGTGGGCGGCATTCTTTGCCACAGGAACCAAGTGAT
GCCCATGGATCAGTCATCCATGCATTCTGACCATGCACAGACAGTAATTCCATATAATCCTTCAAGTCAT
GAGAGTTTGGATCAAGTTGGTGAAGAAAAAGAGGCAATGAACACCAGGGAAAGTGGCAAAGCTTCATCCA
GTCTAGGTCTTCAGGATTTTGAATTTGCTCCGGGTAAATAGGAAGAGGAAGTTATGCCAAAGTACTGTTGGT
TCGATTAAAAAAAACAGATCGTATTTATGCAATGAAAGTTGTGAAAAAAGAGCTTGTTAATGATGATGAG
GATATTGATTGGGTACAGACAGAGAAGCATGTGTTTGAGCAGGCATCCAATCATCCTTTCTTGTGGGC
TGCATTCTTGCTTTTCAGACAGAAAGCAGATTGTTCTTTGTTATAGAGTATGTAAATGGAGGAGACCTAAT
GTTTCATATGCAGCGACAAAGAAAACCTCCTGAAGAACATGCCAGATTTTACTCTGCAGAAATCAGTCTA
GCATTAAATTATCTTCATGAGCGAGGGATAATTTATAGAGATTTGAACTGGACAATGTATTACTGGACT
CTGAAGGCCACATTAACTCCTGACTACGGCATGTGTAAGGAAGGATTACGGCCAGGAGATCAACACAG
CACTTTCTGTGGTACTCCTAATTACATTGCTCCTGAAATTTTAAGAGGAGAAGATTATGGTTTCAGTGT
GACTGTTGGGCTCTTGGAGTGCTCATGTTTGAGATGAGTGAGGAGGTCCTCATTGATATTGTTGGGA
GCTCCGATAACCCTGACCAGAACACAGAGGATTATCTCTTCCAAGTTATTTTGGAAAAACAAATTCGCAT
ACCACGTTCTATGTCTGTAAAAGCTGCAAGTGTTCTGAAGAGTTTTCTTAATAAGGACCTTAAGGAACGA
TTGGGTTGTCTTCTCAACAGGATTTGCTGATATTCAGGGACACCCGTTCTTCCGAAATGTTGATTGGG
ATATGATGGAGCAAAAACAGGTGGTACCTCCCTTTAAACCAATATTTCTGGGGAATTTGGTTTGGACAA
CTTTGATTCTCAGTTTACTAATGAACGTGTCCAGCTCACTCCAGATGACGATGACATTGTGAGGAAGATT
GATCAGTCTGAATTTGAAGGTTTTGAGTATATCAATCCTCTTTTGATGTCTGCAGAAGAATGTGTCTGAT
CCTCATTTTTCAACCATGTATTCTACTCATGTTGCCATTTAATGCATGGATAAACTTGCTGCAAGCCTGG
ATACAATTAACCATTTTTATATTTGCCACCTACAAAAAACACCCAATATCTTCTCTTGTAGACTATATGA
ATCAATTATTACATCTGTTTTACTATGAAAAAAAATTAATACTACTAGCTTCCAGACAATCATGTCAAA
ATTTAGTTGAACTGGTTTTTTCAGTTTTTAAAAGGCCTACAGATGAGTAATGAAGTTATCTTTTTTTGTTTA
AAAAAAAAAAAAAAAA

>gi|10864649|ref|NM_002744.2| Homo sapiens protein kinase C, zeta (PRKZ), mRNA

ATGCCCAGCAGGACCCCAAGATGGAAGGGAGCGGCGGCCGCGTCCGCCCTCAAGGCGCATTACGGGG
GGGACATCTTCATCACCAGCGTGACGCCGCCACGACCTTCGAGGAGCTCTGTGAGGAAGTGAGAGACAT
GTGTCGTCTGCACCAGCAGCACCCGCTCACCTCAAGTGGGTGGACAGCGAAGGTGACCCTTGCACGGTG
TCCTCCCAGATGGAGCTGGAAGAGGCTTTCCGCCTGGCCCGTCAGTGCAGGGATGAAGGCCATCATCTC
ATGTTTTCCCGAGCACCCCTGAGCAGCCTGGCCTGCCATGTCCGGGAGAAGACAAATCTATCTACCGCCG
GGGAGCCAGAAGATGGAGGAAGCTGTACCGTGCCAACGGCCACCTCTTCCAAGCCAAGCGCTTTAACAGG
AGAGCGTACTGCGGTCAAGTGCAGCGAGAGGATATGGGGCTCGCGAGGCAAGGCTACAGGTGCATCAACT
GCAAACTGCTGGTCCATAAGCGCTGCCACGGCCTCGTCCCGCTGACCTGCAGGAAGCATATGGATTCTGT
CATGCCTTCCCAAGAGCCTCCAGTAGACGACAAGAACGAGGACGCCGACCTTCTTCCGAGGAGACAGAT
GGAATTGCTTACATTTCTCATCCCGGAAGCATGACAGCATTAAGACGACTCGGAGGACCTTAAGCCAG
TTATCGATGGGATGGATGGAATCAAAATCTCTCAGGGGCTTGGGCTGCAGGACTTTGACCTAATCAGAGT
CATCGGGCGGGGAGCTACGCCAAGGTTCTCTGGTGCGGTTGAAGAAGAATGACCAATTTACGCCATG
AAAGTGGTGAAGAAAGAGCTGGTGCATGATGACGAGGATATTGACTGGGTACAGACAGAGAAGCAGTGT
TTGAGCAGGCATCCAGCAACCCCTTCTGGTGGGATTACACTCCTGCTTCCAGACGACAAGTCGGTTGTT
CCTGGTCATTGAGTACGTCAACGGCGGGGACCTGATGTTCCACATGCAGAGGCAGAGGAAGCTCCCTGAG
GAGCACGCCAGGTTCTACGCGGCCGAGATCTGCATCGCCCTCACTTCTTGCACGAGAGGGGGATCATCT
ACAGGGACCTGAAGCTGGACAACGTCTCTGGATGCGGACGGGCACATCAAGCTCACAGACTACGGCAT

GTGCAAGGAAGGCCTGGGCCCTGGTGACACAACGAGCACTTTCTGCGGAACCCCGAATTACATCGCCCC
 GAAATCCTGCGGGGAGAGGAGTACGGGTTCAGCGTGGACTGGTGGGCGCTGGGAGTCCTCATGTTTGAGA
 TGATGGCCGGGCGCTCCCGTTTCGACATCATCACCGACAACCCGGACATGAACACAGAGGACTACCTTTT
 CCAAGTGATCCTGGAGAAGCCCATCCGGATCCCCGGTTCTTGTCGTCGTCAAAGCCTCCCATGTTTTAAAA
 GGATTTTTTAATAAGGACCCCAAAGAGAGGCTCGGCTGCCGGCCACAGACTGGATTTTCTGACATCAAGT
 CCCACGCGTTCTTCCGCAGCATAGACTGGGACTTGCTGGAGAAGAAGCAGGCGCTCCCTCCATTCCAGCC
 ACAGATCACAGACGACTACGGTCTGGACAACCTTTGACACACAGTTTACCAGCGAGCCCGTGCAGCTGACC
 CCAGACGATGAGGATGCCATAAAGAGGATCGACCAGTCAGAGTTTGAAGGCTTTGAGTATATCAACCCAT
 TATTGCTGTCCACCGAGGAGTCGGTGTGAGGCCGCGTGCCTCTCTGTCGTGGACACGCGTGATTGACCTT
 TTAACGTATCCTTAACCACCGCATATGCATGCCAGGCTGGGCACGGCTCCGAGGGCGGCCAGGGACAGA
 CGCTTGCGCCGAGACCGAGAGGGAAGCGTCAGCGGGCGCTGCTGGGAGCAGAACAGTCCCTCACACCTG
 GCCCGGCAGGCAGCTTCGTGCTGGAGGAACCTTGCTGCTGTGCCTGCGTCGCGGCGGATCCGCGGGGACCC
 TGCCGAGGGGGCTGTATGCGGTTTCCAAGGTGCACATTTTCCACGGAAACAGAACTCGATGCACTGACC
 TGCTCCGCCAGGAAAGTGAGCGTGTAGCGTCTGAGGAATAAAATGTTCCGATGAAAAAAAAA

>gi|4759123|ref|NM_004171.1| Homo sapiens solute carrier family 1
 (glial high affinity glutamate transporter), member 2 (SLC1A2), nuclear
 gene encoding mitochondrial protein, mRNA

GATAGTGCTGAAGAGGAGGGGCGTTCCAGACCATGGCATCTACGGAAGGTGCCAACAAATATGCCCCAAGC
 AGGTGGAAGTGCGAATGCCAGACAGTCATCTTGGCTCAGAGGAACCCAAGCACCGGCACCTGGGCCTGCG
 CCTGTGTGACAAGCTGGGGAAGAATCTGCTGCTCACCTGACGGTGTGTTGGTGTATCCTGGGAGCAGTG
 TGTGGAGGGCTTCTTCGCTTGGCATCTCCCATCCACCTGATGTGGTTATGTTAATAGCCTTCCCAGGGG
 ATATACTCATGAGGATGCTAAAAATGCTCATTCTCCCTCTAATCATCTCCAGCTTAATCACAGGGTTGTC
 AGGCCTGGATGCTAAGGCTAGTGGCCGCTTGGGCACGAGAGCCATGGTGTATTACATGTCCACGACCATC
 ATTGCTGCAGTACTGGGGGTCAATCTGGTCTTGGCTATCCATCCAGGCAATCCCAAGCTCAAGAAGCAGC
 TGGGGCCTGGGAAGAAGATGATGAAGTGTCACGCTGGATGCCCTTCTGGACCTTATTGCAATCTCTTT
 CCTGAAAACCTTGTC CAAGCCTGCTTTCAACAGATTCAACAGTGACGAAGAAAGTCTGGTTGCACCA
 CCGCCAGACGAGGAGGCCAACGCAACCAGCGCTGAAGTCTCTCTGTTGAACGAGACTGTGACTGAGGTGC
 CGGAGGAGACTAAGATGGTTATCAAGAAGGGCCTGGAGTTCAAGGATGGGATGAACGTCTTAGGTCTGAT
 AGGGTTTTTTCATTGCTTTTGGCATCGCTATGGGGAAGATGGGAGATCAGGCCAAGCTGATGGTGGATTTT
 TTCAACATTTTGAATGAGATTGTAATGAAGTTAGTGATCATGATCATGTGGTACTCTCCCTGGGTATCG
 CCTGCCTGATCTGTGGAAAGATCATTGCAATCAAGGACTTGAAGTGGTTGCTAGGCAACTGGGGATGTA
 CATGGTAACAGTGATCATAGGCCTCATCATCCACGGGGGCATCTTCTCCCTTTGATTTACTTTGTAGTG
 ACCAGGAAAAACCCCTTCTCCCTTTTGGCTGGCATTTTCCAAGCTTGGATCACTGCCCTGGGCACCGCTT
 CCAGTGCTGGAACTTTGCTGTACCTTTCTGTTGCCTGGAAGAAAATCTGGGGATTGATAAGCGTGTGAC
 TAGATTCTGCTCTTCTGTTGGAGCAACCATTAAATGAGTGGTACAGCCCTTTATGAAGCGGTGGCCGCC
 ATCTTTATAGCCCAAATGAATGGTGTGCTCTGGATGGAGGACAGATTGTGACTGTAAGCCTCACAGCCA
 CCCTGGCAAGCGTCGGCGCGGCCAGTATCCCAAGTGCCGGGCTGGTCAACATGCTCCTCATTCTGACAGC
 CGTGGGCCTGCCAACAGAGGACATCAGCTTGCTGGTGGCTGTGGACTGGCTGCTGGACAGGATGAGAACT
 TCAGTCAATGTTGTGGGTGACTCTTTTGGGGCTGGGATAGTCTATCACCTCTCCAAGTCTGAGCTGGATA
 CCATTGACTCCAGCATCGAGTGCATGAAGATATTGAAATGACCAAGACTCAATCCATTTATGATGACAT
 GAAGAACCACAGGGAAGCAACTCTAATCAATGTGCTATGCTGCACACAACCTCTGTATAGTAGATGAA
 TGCAAGGTAACCTGGCAGCCAATGGAAAGTCAGCGACTGCAGTGTGAGGAAGAACCCTGGAAACGTG
 AGAAATAAGGATATGAGTCTCAGCAAATTCTTGAATAAACTCCCCAGCGT

>gi|31543627|ref|NM_004172.2| Homo sapiens solute carrier family 1
 (glial high affinity glutamate transporter), member 3 (SLC1A3), mRNA
 AGCGGGGTAGTAACCTTGCACTTTCAGAGCACATGCACACTGTCAGGGCTAGCCTGCCTGCTTACGCGCGC
 TGGCGGATTGTTGCTCCGTTGTACCTGCTGGGGAATTACCTCGTTACTGCTCGATATCTTCCACCCCTTA
 CAAAATCAGAAAAGTTGTGTTTTCTAATACCAAGAGGAGGTTTGGCTTTCTGTGGGTGATTCCCAGACA
 CTGAAGTGCAAGAAGAGACCCTCTAGAAAAGTAAAATATGACTAAAAGCAATGGAGAAGAGCCCAAGA
 TGGGGGGCAGGATGGAGAGATTCCAGCAGGGAGTCAGTAAACGCACACTTTTGGCCAAGAAGAAAGTGCA
 GAACATTACAAAGGAGGATGTTAAAAGTTACCTGTTTCGGAATGCTTTTGTGCTGCTCACAGTCACCGCT
 GTCATTGTGGGTACAATCCTTGGATTTACCCTCCGACCATACAGAATGAGCTACCGGGAAGTCAAGTACT
 TCTCCTTTCTGGGGAACCTTGTATGAGGATGTTACAGATGCTGGTCTTACCATTATCATCTCCAGTCT
 TGTACAGGAATGGCGGCGCTAGATAGTAAGGCATCAGGGAAGATGGGAATGCGAGCTGTAGTCTATTAT
 ATGACTACCACCATCATTGCTGTGGTGATTGGCATAATCATTGTATCATCATCATCCTTGGGAAGGGCA

CAAAGGAAAACATGCACAGAGAAGGCAAAATTGTACGAGTGACAGCTGCAGATGCCTTCCTGGACTTGAT
 CAGGAACATGTTCCCTCCAAATCTGGTAGAAGCCTGCTTTAAACAGTTTAAACCAACTATGAGAAGAGA
 AGCTTTAAAGTGCCCATCCAGGCAACGAAACGCTTGTGGGTGCTGTGATAAACAATGTGTCTGAGGCCA
 TGGAGACTCTTACCCGAATCACAGAGGAGCTGGTCCCAGTTCCAGGATCTGTGAATGGAGTCAATGCCCT
 GGGTCTAGTTGTCTTCTCCATGTGCTTCGGTTTTGTGATTGGAAACATGAAGGAACAGGGGCAGGCCCTG
 AGAGAGTTCTTTGATTCTCTTAACGAAGCCATCATGAGACTGGTAGCAGTAATAATGTGGTATGCCCCCG
 TGGGTATTCTCTTCTGATTGCTGGGAAGATTGTGGAGATGGAAGACATGGGTGTGATTGGGGGCGAGCT
 TGCCATGTACACCGTGACTGTCATTGTTGGCTTACTCATTACGCAGTCATCGTCTTGCCACTCCTCTAC
 TTCTTGGTAACACGAAAAACCCCTGGGTTTTATTGGAGGGTTGCTGCAAGCACTCATCACCGCTCTGG
 GGACCTCTTCAAGTTCTGCCACCCATACCATCACCCTCAAGTGCTTGAAGAGAACAATGGCGTGGACAA
 GCGCGTCACAGATTCTGTGCTCCCCGTAGGAGCCACCATTAAACATGGATGGGACTGCCCTCTATGAGGCT
 TTGGCTGCCATTTTCATTGCTCAAGTTAACAACCTTTGAAGTGAAGTTCGGACAAATTATTACAATCAGCA
 TCACAGCCACAGCTGCCAGTATTGGGGCAGCTGGAATTCCTCAGGCGGGCTGGTCACTATGGTCATTGT
 GCTGACATCTGTGCGGCTGCCACTGACGACATCACGCTCATCATCGCGGTGGACTGGTTCTTGGATCGC
 CTCCGGACCACCACAAACGTACTGGGAGACTCCCTGGGAGCTGGGATTGTGGAGCACTTGTACGACATG
 AACTGAAGAACAGAGATGTTGAAATGGGTAACCTCAGTGATTGAAGAGAATGAAATGAAGAAACCATATCA
 ACTGATTGCACAGGACAATGAACTGAGAAACCCATCGACAGTGAAACCAAGATGTAGACTAACATAAAG
 AAACACTTCTTGAGCACCAGGTGTTAAAAACCATTATAAAATCTTTCCATCTCATTACAGCTCATTTCGC
 TCAGCAAGCCCGTCATCTTCCCTTTCCCTCCCTCTGATAAGACTGGAATAAGTCTCCAAAACACAAG
 GGAGGATTTTGGGTGGCCAAAGTGTACAATTTTCATCCACAATTGAAATTTTTAAATCATTTTCATGTTA
 GTCTTACCGAATAAGGTACCAAGATCACAATAAGTGTGATCAGATCTTACAAGTTTATGTGGCACACAA
 TCCTATAAATGTGATTTTTTTTATATAAGTTAAAGAGACAAATAGTAGGCTAAAAACATTTTAAATCAAC
 TTTTGAAATTTAAAAATCTTTTCAAGATACAATTCAGTTTTAGTTTTCAAAATGTTAACAACCTGAATTACA
 ACCGGTTATCAGTTGGACAGTAAGATTTTATCCCTTCTCTCTGACTGGTATACCTATTTTCATTAGTAG
 CTAGGTGCACATATACATCTAGCACAGCTGTGAGGACAGACAGAAGGCAAAGTTTCCATGTGGCCTTGAG
 CAAGTCCCATCTCACCTCTAGGCCTCAGTGTCTCATCTATAAAATGAGGGACTTCCCTAGAAGTCTTCA
 TGGTCTCTTCCAGCCAGACATCTGTGATGTATGAAAGCACCTGCCCTCTGTTTTCCCTCAGAACACC
 CTGTACCATCCAGTGGACACGAGGCTTCAGAAAAGACACTTCAATGGGAGTGAACATTTCTCAATAGG
 ACAGGATGGCTGTGTGTGGTGGTACCAGGTCTGTGAGCAAAGTGCAGGTATGCAAGTCCGAGGACAG
 GAGGCCATTCCAGGAGTGGGATTATTCATCAAACTCTTTGCCAGTTTCATCCCAATGGGGGAAGTATTCC
 CTTCTTTCTACTCTGGGAAGAATGTCTCTGCCACTCCTCAACTGATGATAGACTTCGAAAACAGATGA
 GAAGACTAGCAGCTAGCAAGGGTGTCTGTAGTCACACTGTGGAACACTAAAGAGCTAGGAAAGAGTTGAG
 CACAGGCAACATTACAAACAAAGGATTTGAAAACACCAAGAGTACAGGTCTTCTTTAAGGAAGAATAAAA
 AAGAAGAGGTTCAATTTTCTGGCTTTTTTTTTCACCTGAAACACTTTTTCTCGAGTCCAAAATCATTTCC
 CCCGTGAAGTCTGCTTACCAAAACATAAGACGACTTATATATTTGAAAGAAGTCAAATGAATGAGCTCTC
 TAATAGAAGTCCATGAGTTGAGTGGGTATTTCTTATTTGAAAGTGTTTTTCTTTAATCAAAGTCCCTTAG
 AATGAGGGAAACAAAATATTTATTTGTTTTGGAATCCCACTTATCAAATCATTCAAAACTTTCAGCTGGA
 GTGGGGTTTTGCTTTTTGTTTTGTTGTGTCCATAAGAGAAATGGTAGAAGATGAATCAGTATGAAGACACT
 GTCAATGAGGTATGAGAAAAACAGCAGGGGCATTAGTTTCAGGCAAGGCAGCTCCCAGGTTTAGAGATT
 AATTTTTACCCCTAAGGAATATCCAGTCAAAGACGCTGAGTGGGAGCTGTCAGGCAGTAGCAGCTGTGT
 TTGAGTTTCTGGCTGAAAATGGTGAAGAATGGACTTAATCATGCTAACAACTGAAAAATCTAGACATAG
 ATCCTCTGATATACAATTAGAGATATTTTTATATAGACCCCAAGCATTCTGTGCATAAAAGTTAACATTA
 GGCTGTGGTGCAGTAACCATTTAATGTGAGGCTCTATTTTCGAAATACACTACAAATGTTAAAGTACGT
 GGCTGTCCCTCTTAAGACACTAGTAGAGCAAAGACTTAATCATATCAACTTAATTCGTTACACAATATGT
 GTTTTTTTAATATACTAACCATTTCTTATGGAAGGTCTGTGGGGAGCCCATCATCTCGCCAAGCCATC
 ACAGGCTCTGCATACACATGCACTCAGTGTGGACTGGGAAGCATTACTTTGTAGATGTATTTTCAATAAA
 GAAAAAATAGTTTTACATTAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAG

>gi|20070238|ref|NM_006671.2| Homo sapiens solute carrier family 1
 (glutamate transporter), member 7 (SLC1A7), mRNA
 GGCACGAGGCTGGTGTTTAGCAACTCCGACCACCTGCCTGCTGAGGGGCTAGAGCCCTCAGCCCAGACCC
 TGTGCCCCCGGCCGGGCTCTCATGCGTGGAATGGTGTGTGCCCCCTTGCCAGCAGGCCAGGCTCACCATG
 GTGCCGCATGCCATCTTGGCACGGGGGAGGGACGTGTGCAGGCGGAATGGACTCCTCATCCTGTCTGTGC
 TGTCTGTCTATCGTGGGCTGCCTCCTCGGCTTCTTCTTGGAGACCCGGCGCTCTCACCACAGGAAATTAG
 TTACTTCCAGTTCCCTGGAGAGCTCCTGATGAGGATGCTGAAGATGATGATCCTGCCACTGGTGGTCTCC
 AGCTTGATGTCCGACTTGCCCTCCCTGGATGCCAAGACCTCTAGCCGCTGGGCGTCTCACCCTGGCGT
 ACTACCTGTGGACCACCTTCATGGCTGTCTATCGTGGGCATCTTCATGGTCTCCATCATCCACCCAGGCAG

CGCGGCCAGAGAAGGAGACCACGGAGCAGAGTGGGAAGCCCATCATGAGCTCAGCCGATGCCCTGTTGGAC
 CTCATCCGGAACATGTTCCAGCCAACCTAGTAGAAGCCACATTCAAACAGTACCGACCAAGACCACCC
 CAGTTGTCAAGTCCCCCAAGGTGGCACCAGAGGAGGCCCTCCTCGCGGATCCTCATCTACGGGGTCCA
 GGAGGAGAATGGCTCCCATGTGCAGAACTTCGCCCCGGACCTGACCCCGCCGCCCGAGGTCTGTTACAAG
 TCAGAGCCGGGACACAGCGATGGCATGAATGTGCTGGGCATCGTCTTCTTCTCTGCCACCATGGGCATCA
 TGCTGGGCGCATGGGTGACAGCGGGGCCCCCTGGTCAGCTTCTGCCAGTGCCTCAATGAGTCGGTCAT
 GAAGATCGTGGCGGTGGCTGTGTGGTATTTCCCCCTTCGGCATTTGTGTTCTTCATTGCGGGTAAGATCCTG
 GAGATGGACGACCCAGGGCCGTCGGCAAGAAGCTGGGCTTCTACTCAGTCACCGTGGTGTGCGGGCTGG
 TGCTCCACGGGCTCTTTATCCTGCCCTGCTCTACTTCTTCATCACCAAGAAGAATCCCATCGTCTTCAT
 CCGCGGCATCCTGCAGGCTCTGCTCATCGCGCTGGCCACCTCCTCCAGCTCAGCCACACTGCCCATCACC
 TTCAAGTGCCGTGCTGGAGAACAACCACATCGACCGCGCATCGCTCGCTTCGTGCTGCCCGTGGGTGCCA
 CCATCAACATGGACGGCACTGCGCTCTACGAGGCTGTGGCCGCCATCTTCATCGCCAGGTCAACAACATA
 CGAGCTGGACTTTGGCCAGATCATCACCATCAGTATGCAGCCACTGCAGCCAGCATTTGGGCGAGCTGGC
 ATCCCCAGGCCGGCTCGTCAACCATGGTCATCGTGTCACTTCCGTGGGACTGCCACCGATGACATCA
 CCCTCATCATTTGCCGTTGACTGGGCTCTGGACCGTTTCCGCACCATGATTAACGTGTGGGTGATGCGCT
 GGCAGCGGGGATCATGGCCCATATATGTCGGAAGGATTTTGCCCGGGACACAGGCACCGAGAACTGCTG
 CCCTGCGAGACCAAGCCAGTGAGCTCCAGGAGATCGTGGCAGCCAGCAGAATGGCTGTGTGAAGAGTG
 TAGCCGAGGCTCCGAGCTCACCTGGGCCCCACCTGCCCCACCACGTCCCCGTTCAAGTGGAGCGGGA
 TGAGGAGCTGCCCCTGCGAGTCTGAACCACTGCACCATCCAGATCAGCGAGCTGGAGACCAATGTCTGA
 GCCTGCGGAGCTGCAGGGCAGGCGAGGCTCCAGGGCAGGGTCTGAGGCAGGAACCTCGACTCTCCAA
 CCCTCCTGAGCAGCCGGCAGGGCCAGGATCACACATTTCTTCTCACCTTGAGAGGCTGGAATTAACCC
 GCTTGACGGAAAAATGATCTCAGAGAAGGGAAAGGCTGCATGGGGGAGCCCCATCCAGGGAGTGATGGGC
 CCGGCATTTGCTGAGGCCCCGCTGTGACAGTTTCCCCGGTGTGAGCCCGGTGAGGGCGGCAGGCAGGGGT
 TATCCGGCCCCACTTTCTGGATGACAGACTTGAGGCTCTGAGAGCTGAAAACACTTGTCCAAGGCTCTAC
 GTTAAGGTCAAGACACTAACTCAAATCTTTCAAGCCCCGCTCTCCTCTTGGAGGACAGGGCAGCTTGCA
 GCTGTGTCCAGGCCAGGCCACCCCATACAGGTGGCCTCAGCCACACAGTTCTCCCCAAGGGGAGCA
 GCCAGGGCCAAGCCCCGCTGCCTTCCCCAGGCCACAGTGCCTCAGTCTCCTGTCTGCCACGTGTCTT
 TTGCAAAGCTCCTTGGATGTGGAGACAGATGCTTTACTAGAGCTGAAAGGCCCTTGCACATCATCAGG
 CCAACCTCCCATGGAATAGGTAGGCAAGCCAGGACTCCGGGAAGGAGGTGCAGCCAGGATGCTCTGGTGG
 AGCTGCCGATGGGGCCCTGGTGTGCAAACTCCCCAAAGGCCTGTGCGTCCAAGTGGAGTCAGGTTTTCTA
 TTCTTTCTGTGTTTGCAAATTCAGTGTTAACTAAATAAAGGTATTTTGTTTTTCAAAAAAAAAAAAAAA
 AAA

>gi|21314631|ref|NM_003038.2| Homo sapiens solute carrier family 1
 (glutamate/neutral amino acid transporter), member 4 (SLC1A4), mRNA
 CCCGACTCTGCGCCTCTCCTCGCCTTTCTCGCACCTGCTCCTGCGCCAGGCCGCGAGACCCCCGGGGCG
 GCTTCCCAGAACCTGCGGAGCACAACTGGCCGACCGACCCATTTCATTGGGAACCCCGTCTTTTGGCAGAG
 CCCACGTCCCCCTGCCACCTCTAGCTCGGAGCGGCGTGTAGCGCCATGGAGAAGAGCAACGAGACCAACGG
 CTACCTTGACAGCGCTCAGGCGGGGCTGCGGCCGGGCCCCGAGCTCCGGGGACCGCGGGCGGACGCGCA
 CGGCGTTGCGCGGGCTTCTTGC GGCGCCAAGCGCTGGTGTGCTCACC GTGTCCGGGGTGTGCGGGGCG
 CGGGCTTGGGCGCGGCGTGTGCGGGGCTCAGCCTGAGCCGCACGCAGGTCACCTACCTGGCCTTCCCCGG
 CGAGATGCTGCTCCGCATGCTGCGCATGATCATCTGCCGCTGGTGGTCTGCAGCCTGGTGTGCGGGCGCC
 GCCTCCCCTCGATGCCAGCTGCCCTCGGGCGTCTGGGCGGCATCGCTGTGCGCTACTTTGGCCTCACACAC
 TGAGTGCCCTCGGCGCTCGCCGTGGCCTTGGCGTTTCATCATCAAGCCAGGATCCGGTGGCGAGACCCTTCA
 GTCCAGCGACCTGGGGCTGGAGGACTCGGGGCTCCTCCTGTCCCCAAAGAGACGGTGGACTCTTTCTCTC
 GACCTGGCCAGAAACCTGTTTCCCTCCAATCTTGTGGTTGCAGCTTTCCGTACGTATGCAACCGATTATA
 AAGTCGTGACCCAGAACAGCAGCTCTGGAAATGTAACCCATGAAAAGATCCCCATAGGCAC TGAGATAGA
 AGGGATGAACATTTTAGGATTGGTCTGTTTGTCTGGTGTTAGGAGTGGCCTTAAAGAACTAGGCTCC
 GAAGGAGAAGACCTCATCCGTTTCTTCAATTCCCTCAACGAGGCGACGATGGTGTGGTGTCTGGATT
 TGTGGTACGTACCTGTGGGCATCATGTTCTTGTGGAAGCAAGATCGTGGAAATGAAAGACATCATCGT
 GCTGGTGACCAGCCTGGGGAAATACATCTTGCATCTATATTGGGCCATGTTATTCATGGAGGAATTGTT
 CTGCCACTTATTTATTTTGTTTTACACGAAAAAACCCATTTCAGATTCTCTCTGGGCCCTCTCGCCCCAT
 TTGCGACAGCATTTGCTACCTGCTCAGCTCAGCGACCTTCCCTCTATGATGAAGTGCATTGAAGAGAA
 CAATGGTGTGGACAAGAGGATCAGCAGGTTTATTCTCCCCATCGGGGCCACCGTGAACATGGACGGAGCA
 GCCATCTTCCAGTGTGTGGCCGCGGTGTTTCATTGCGCAACTCAACAACGTAGAGCTCAACGCAGGACAGA
 TTTTACCATTCTAGTGACTGCCACAGCGTCCAGTGTGGAGCAGCAGGCGTGCCAGCTGGAGGGGTCTCT
 CACCATTGCCATTATCTGGAGGCCATTGGGCTGCCTACTCATGACCTGCCTCTGATCCTGGCTGTGGAC

TGGATTGTGGACCGGACCACCACGGTGGTGAATGTGGAAGGGGATGCCCTGGGTGCAGGCATTCTCCACC
 ACCTGAATCAGAAGGCAACAAAGAAAGGCGAGCAGGAACCTTGCTGAGGTGAAAGTGGAAGCCATCCCCAA
 CTGCAAGTCTGAGGAGGAGACATCGCCCCCTGGTGACACACCAGAACCCCGCTGGCCCCGTGGCCAGTGCC
 CCAGAAC TGGAATCCAAGGAGTCGGTTCTGTGATGGGGCTGGGCTTTGGGCTTGCCTGCCAGCAGTGATG
 TCCCACCCCTGTTACCCAGCCGCCAGTCATGGACACAGGGCACTGCCCTTGCCAACTTTTACCCTCCCAA
 GCAATGCTTTGGCCCAGTCGCTGGCCTGAGGCTTACCTCTCGGCACTGGCATTGGGCTCCCCAGCCGGA
 CTGGTTACCAAGGACAAGGACACTCTGACATTCGGCTTGATCCATGTCCAGGTGCAACTGTGTGTACACC
 AGGGATCTGTTTGGAAACAACCCCTTGAGCTGCCAGGCTCAAGAAATCATGGACTCACAGGGTCTGTGT
 GGTACATCTTGGAATAATGCAGATGTATTTCACTCTCCCCGGTCAGCTCTGCATCAGGTGTTTTCTGA
 GCAAACCAAGGGGGTTTATAGTCACTGTGCGCATTCGCTCGAGTTGCAGTAATTGAAAAATGCTCAAAT
 TCTTAGCCATGGCTGGCCTTTGCTGAGCTGGGACTCAGGTGTTTAAAGAGTTTGTGCTATAGCTAGGTGT
 GGATAGCTTCTGATCCCTGGGTTCGAGGAGACTGCAGGTGCCGCACATTGTCAAGTTAGAAATACTCCAG
 GTGGGTGTTAGCACTGTGGTGGTCTCTGGTCCACAGCCTTAGGTAAACAACCTAGATTCTGAGGTCAAAG
 AAAAAAGGAGAGGGAATGCAGCCTTGTGGGGGAGAAGCGGGGAGAGGGTTCTCTAATCTAATCAGGACA
 GGACAGGTTTCACATACAATGTCCAGTTTCGCATCCAGCCCTGGGGCACTTTTCTGCTTCTTCCAGA
 GGCCTGGGCCTCTGATAACACTTTGGCTTTTTCTCCATTCACGCTGATTTGGCAAAAGGCCAGAGATGGG
 CCTCCTTCCCTGGGGAGGTGTGATGTAGTTATCACATTCAGGACCCTTGTTGATTTATCATCTATTATTT
 GAATTCACCTGGACACTCTGTAAATGCTGCACTGCAGCAAAAAACAAACCACCACCACCCAGAGAAAA
 CCATGTACTAATTGGAGTGGGGTACCCCCATTCACAGGTTCCAGGTCCCCTGGCTTTGGCTGATTTCAA
 AATATAGAGCCCTTCTTGCCAGTACATCCAAGTTTAAATATATCAGCGAAATGGTCCATGTTTTTCCAA
 TTACCTGCTGACACGGTTCTAAGCTAAGTGAAGGGGAAGATCTGAGAGCGTGCTGTTTGTGGCTGTTGAT
 GCATATTCGTGATGTAACAGGTCTTGGGGCTCACTTTACCCCATTTGTAAATGGGGCTAATGTACCT
 GCCTCTTACCTACCTCAGAGGGATTTGGTGAAGCAAACCTGTTAATCTTCGAAAACGACCATTTCACCTCT
 TGGATATCAAGTGCTAACCCAGTATGTTCTTCTTTTTATGTAAGGGACAGCTTTGGAGAAAGGACTGCT
 CTGTGGAGCAGAGTCCCTTCTGCTGGTGAGGACAGCATTTCTGAGCAGGGCTTTGTTCTCTATGTGCATT
 AGGACTTTTATCATGCCCTTGTCTGTGTGTAGTTACTTGACAGCATCAAATGCCGCCTCTTCTTAATGT
 CCTTCAAGTTTTCATGAAC TAGCAACCCACCTTCCACCATTGTTCTGGGGCCTGATTTTGTCTGACT
 CCCAGACCCAGCCACTGTTTCTGCCACCCTGTAAACAGGCCATTAAAGCTCCCAGTGTTTCAGCCTCCTTC
 ACTCCCTTGTTTTCCCTGTTGCTATGTGTCACTGGGCCCTACAGACAGGGGCACACGCTTATGGATGTG
 TGTACCATTGAGATGAGAATGGGTAGATGGAACGGAGACCATCAAGCCACACCCCTTCTTAAACTGGG
 GACATGAGCCTGAGCAGAAAGGGTGAAGAAGAGCCATGGGACACAGAGTTGACCCAGCCAGGGGAAAGC
 CCAGCTCTCTTTAAACCAGCTAAGCCATTCCAGTCTCCTGTGAAGCCAAAAGGGACCAGGAACCGTGCAA
 AGGAAACTGGAACTTTTCCCCGCTGGGTAGAGCATGTTGCTGATACTCTTCTGTTTTCAAGGGAAACAA
 TCACATTGTTTGATTCCAAATGGTAAATGAACACTCACTATTCTTCAGGCTTCAGTAAATCTTTTTTCT
 TCCTTCATATATATATACACAACACACACACATATGTATATCTATACACACATGTGTGTTGTGTATAT
 GCATGTGTGTGTGTGTGCGTGTGTGTATAGTTTTAGCTCCAAGCCAAGCAAGTTTGTGTTTGGATAGAGG
 GGAAC TTAAC TATTAAC TACAAGTTGTATGTCTGTGGTATCTTGATTTTCCCATTTCTAAAGATGAATTT
 CACAAAGCCATAAAGCGTGAAATTAGAGCTGGACTTAAGACTCATTGGCCGACCATCCTGTGTCTCTGGCC
 TGGCCCTGCAGTAAGAAGCGTGTCTGGGTCTGGAGAAGGGTGCTTCCGAGAGTGTGCAGGTGGCCCTTCC
 CCTTGGAGGCGAGAAGAGAGAATGTGCTGTCTATCTTCTGGTTTTTCACTCCACAGAGTCGGTAGACCAG
 GGGTTACGTGACTGGGGAAATCTCACATCTCCTTGTCTGAAAACATTTCCCTGCTGTTCTCTTTCTAA
 CATGTTGTGGTAAATCTGTTTCACTACTGCTCATCTGACTGTTTTGTACATGTGACAATTGCCTTAAAC
 CTAGCACAGTCTCAGAAATGAATACCGTGTTTCCACTGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
 AAAAAAAAAA

>gi|4827011|ref|NM_005071.1| Homo sapiens solute carrier family 1 (high
 affinity aspartate/glutamate transporter), member 6 (SLC1A6), mRNA
 GATAGACCATGAGCAGCCATGGCAACAGCCTGTTCTTTCGGGAGAGCGGCCAGCGGCTGGGCCGGGTGGG
 CTGGCTGCAGCGGCTGCAGGAAAGCCTGCAGCAGAGAGCACTGCGCACGCGCTGCGCCTGCAGACCATG
 ACCCTCGAGCACGTGCTGCGCTTCTGCGCCGAAACGCCTTCATTCTGCTGACGGTCAGCGCCGTGGTCA
 TTGGGGTCAGCCTGGCCTTTGCCCTGCGCCCATATCAGCTCACCTACCGCCAGATCAAGTACTTCTCTTT
 TCCTGGAGAGCTTCTGATGAGGATGCTGCAGATGCTGGTGTACCTCTCATTTGCTCCAGCCTGGTCACA
 GGTATGGCATCCCTGGACAACAAGGCCACGGGGCGGATGGGGATGCGGGCAGCTGTGTACTACATGGTGA
 CCACCATCATCGCGGTCTTCATCGGCATCCTCATGGTCACCATCATCCATCCCGGGAAGGGCTCCAAGGA
 GGGGCTGCACCGGGAGGGCCGGATCGAGACCATCCCCACAGCTGATGCCCTTCATGGACCTGATCAGAAAT
 ATGTTTCCACCAAACCTTGTGGAGGCCTGCTTCAAACAGTTCAAGACGCAGTACAGCACGAGGGTGGTAA
 CCAGGACCATGGTGAGGACAGAGAACGGGTCTGAGCCGGGTGCCCTCCATGCCCTCTCCATTCCTCAGTGA

ACGCGGAGGAGCCGGGCGCGCCTGCCACGCAAAACTACCGGGCTGGCAGGGCGGGCGGGCGGGTGCAGCAT
CCCGGGTGGCGGCGGCAACGGCGGTGGTGACGGCGGCGACTGCAGCGGCCGGCTCTCACCTCTCCCCTGT
GCACCCGCATCTCGCCGCGCCGCCGAGCAGCCAGCAGTCCCCGGGTGCCCCAGCCACGCGCGCACGGCC
GAGCCACGCGACAATAGCGGCGACAGCCATGGGGAAACCGGCGAGGAAAGGATGCGAGTGGAAAGCGCTT
CCTGAAGAATAACTGGGTGTTGCTGTCCACCGTGGCCGCGGTGGTGCTAGGCATTACCACAGGAGTCTTG
GTTTCGAGAAACACAGCAACCTCTCAACTCTAGAGAAATTCTACTTTGCTTTTCCCTGGAGAAATTCTAATGC
GGATGCTGAAACTCATCATTTTGCATTAAATTATATCCAGCATGATTACAGGTGTTGCTGCAC TGGATT
CAACGTATCCGGA AAAAATTGGTGTGCGCGCTGTCTGTATTATTTCTGTACCAC TCTCATTGCTGTTATT
CTAGGTATTGTGCTGGTGGTGAGCATCAAGCCTGGTGTCAACCAGAAAGTGGGTGAAATTGCGAGGACAG
CGCAGCACCCCTGAAGTCAGTAGCAGGTGCAATGTTAGATCTCATCAGGAATATGTTCCCTGAGAATCT
TGCCAGGCCGTGTTTTCAGCAGTACAAAACTAAGCGTGAAGAAGTGAAGCC TCCAGCAGATCCAGAGATG
AACATCGCAAGAGATCCTTTCAGAGCTGTATGACAAC TGAATTTCCAAGAACA AAAACAAAGGAATACA
AAATTGTTGGCATGTATT CAGATGGCATAAACGTCCTGGGCTTGATTGCTTTTGCCTTGCTTTTGACT
TGTCATTGGAAAAATGGGAGAAAAGGGACAAATTCTGGTGGATTTC TTTCAATGCTTTGAGTGATGCAACC
ATGAAAATC GTTCAGATCATCATGTGTTATATGCCACTAGGTATTTTGTTCCTGATTGCTGGGAAGATCA
TAGAAGTTGAAGACTGGGAAATATTCGCAAGCTGGGCCTTTACATGGCCACAGTCC TGACTGGGCTTGC
AATCCACTCCATTGTAATTCTCCCGCTGATATATTT CATATGTCGTACGAAAGAACCCTTTCCGATTGCGC
ATGGGAATGGCCCAGGCTCTCCTGACAGCTCTCATGATCTCTTCCAGTTCAGCAACACTGCCTGTCACCT
TCCGCTGTGCTGAAGAAAATAACCAGGTGGACAAGAGGATCAC TCGATTCTGTGTTACCCGTTGGTGC AAC
AATCAACATGGATGGGACTGCGCTCTATGAAGCAGTGGCAGCGGTGTTTATTGCACAGTTGAATGACCTG
GACTTGGGCATTGGGCAGATCATCACCATCAGTATCACGGCCACATCTGCCAGCATCGGAGCTGCTGGCG
TGCCCCAGGCTGGCCTGGTGAGCATGGTGATTGTGCTGAGTGGCCGTGGGCC TGGCCGCCGAGGATGTCAC
CTGTGATCATTTGCTGTCTGACTGGCTCTGGACCGGTTCAGGACCATGGTCAACGTCC TTTGGTGATGCTTTT
CGGACGGGCATTGTGGAAAAGCTCTCCAAGAAGGAGCTGGAGCAGATGGATGTTTCATCTGAAGTCAACA
TTGTGAATCCCTTTGCCTTGGAATCCACAATCCTTGACAACGAAGACTCAGACACCAAGAAGCTTATGT
CAATGGAGGCTTTGCAGTAGACAAGTCTGACACCATCTCATTCACCCAGACCTCACAGTTCTAGGGCCCC
TGGCTGCAGATGACTGGAAACAAGGAAGGACATTTCCGTGAGAGTCA TCTCAAAACACTGCTTAAAGGAAAA
GAGAAACACTAATGGCCAAGTGACATTTGATTGATATACAGACC TCCAGATTATTTTCTATATTTGGA
TTCACAGCCTTTGCGCTCTGGGTTTTGGGATTGGGTGTGGGGTAAGTTGAAGGGAAATCAATTTAAAGG
AAAGTTCTATTATCTGGGTTTTAGAAATTCTATAAGAGACAAAGTTTGGAGTACATAAAGTAATAACTG
TTAGAATTAGGTAATGGATATGAAAGAGAAAATGCTTTCTCATGCATAGACAAGTGTTTTGGGTTTTTAA
AAAAAATATTCTGTCAATTGGTTACAAATTTTTTACTCAGGCTTTCTATTGGCATGGATTTCCCTTTGACCTC
TCACTTTTTTTATAAATTATAATGCATCTAAACCACCTGTCCCCAGTTAATGTGCCAAAAATGTCAATTTTT
AACTTATCTCCAGCCAATTTCAAGAAAACAGACCAGCATAGTTCTGCAATAACAGTTT TTAAGATGGGCA
TAGGGTTTTGGAAGAAAAGAGAGAAGGATATTTTTTTCAATGTACTGTATTGGGACGCTGGTAACTGTTAAC
CCAGTGTTCAAGCATAGAGCATATATATATATATGATATATGATATTTATTTTTCATATAATTTGCCAGA
CAGAGATCAGAATTGAACCGTCAATGTGAAATAAAGAGTTCTCCTTG TACTTGTAATAAACCACGATTC
CAACCCAGGTCTGCTTTTGGGGCTTATCAGAAC TCTTTCTAAGGAGCATAAGATGAGAAATCATGTTGT

TCGATCGTTTCACATCTGTATATCAGCTCTAAAGCAGAGATGTATTATGGTGATACTCCAAGGTGGCATA
GCCATTTCATTTACAACTTCCAGATTTGAGCTGCCTGGAGGGAATCCATATCAGCTCTGCATAAGATTATA
TACAAAGCTGTCACTCACAAAAGGCTGGATGTGCTTTTCATCCAACCTGGAAGGCTTTATTCTTCCAAGTTC
ATTCACTACTCAAAGAGGCCAGTACTTTGCCATCCTTGCACCTTTCTGTTATCAGGGCCCAAATAACAGTGG
CAAGCTACCAACTAAGTTGTATTTTAATAAAGATTCCATGGGTTGAACAAGCCACGTTGCAGAAAAAGAG
CTTCCCCTAACCTGGGTGTGTCAGAGTAAATCCCACGACATAAGCTGGTATCACTGGTTTCGGGGGAAAT
AGTTCCATTCTATGACTCTTGTCTCCTCCTCCAGGAGGACTGTTCTAACTAGTAATCTTGGCCCTATTCA
TTACATCCTCTGCTTGTCAATTCTGCTAATTTATGAAGATAGTTTATTATAGTCTGTACTTCAGTTCTCAT
CTTGTAATAATAGCTTAACATAAACTTGTACTTACACTGAAATCCAAAATAGTCATGTTTCTGCAGTATT
CTGTAGCCAACTTAAACCTGTGCTTTCATGTTTAAAGAAATGAGAAATTGTGCCAAAGATAGCAGAAGAGT
AGATAAGTGCTCAGTATTGACGACCTACATCTGAAATCTACAACATAATGATACTGAATTGTTATGTAA
CATCATAAATAGTAAATAATGATTCAATGTGAATTTTAAATGCAAATATTGCTATTGTTTATAGGAAAT
AAATCTAAATATAAACGAAAAAAAAAAAAAAAAA

>gi|5032092|ref|NM_005628.1| Homo sapiens solute carrier family 1
(neutral amino acid transporter), member 5 (SLC1A5), mRNA
GTAACCGCTACTCCCGACACCAGACCACCGCTTCCGTACACAGGGGCCCGCATCCACCCCTCCCGGAC
CTAAGAGCCTGGGTCCCTGTTCCTGGAGGTCCGCTTCCCGGCCCCCAGATTCTGGCATCCAGCCCTCA
GTGTCCAAGACCCAGGCAGCCCGGGTCCCGGCTCCCGGATCCAGGCGTCCGGGATCTGCGCCACCAGAA
CCTAGCCTCCTGCAGACCTCCGCCATCTGGGGGCACTCAACCTCCTGGAGCCAAGGGCCCCACGTCCAC
CCAGAGAACTCTCGTATTCCAGCTCCTAGGGCCAAGGAACCCGGGCGCTCCGAACCTCCAGCTTTCGG
ACATCTGGCACACGGGGCAGAGCAGAGAAGCTCAGCGCCAGCCTGGGGAATTTAAACACTCCAGCTTCC
AAGAGCCAAGGAACCTTCAGTGCTGTGAACCTACAACCTTAAGGAGCCCTCCAAAGTTCCAGTCTCCAGGT
GCTGTTACTCAACTCAGTCTTAGGAACGTCCGGTCTGGGAAGGAGCCCAAGCGCTCCAGCCAGCTTCC
AGGCGCTAAGAAACCCCGGTGCTTCCCATCATGGTGGCCGATCCTCCTCGAGACTCCAAGGGGCTCGCAG
CGGCGGACCCACCGCCAACGGGGGCTGGCGCTGGCCTCCATCGAGGACCAAGGCGCGGACAGGCGG
CTACTGCGGTTCCTGGGACAGGTGCGCGCTGCTGCTTCGAGCCAACCTGCTTGTGCTGCTGACAGTGTG
GCCGTGGTGGCCGGCTGGCGCTGGGACTGGGGGTGTCGGGGGCCGGGGGTGCGCTGGCGTTGGGCCCCG
AGCGCTTGAGCGCCTTCGTCTTCCCGGGCAGCTGCTGCTGCGTCTGCTGCGGATGATCATCTTGCCGCT
GGTGGTGTGCAGCTTGATCGGCGGCGCCGAGCTGGACCCGGCGCGCTCGGCCGTCTGGGCGCTGG
GCGCTGCTCTTTTTCCTGGTCACCACGCTGCTGGCGTCCGGCTCGGAGTGGGCTTGGCGCTGGCTCTGC
AGCCGGGCGCGCCTCCGCCGATCAACGCCTCCGTGGGAGCCGCGGGCAGTGCCGAAAATGCCCCAG
CAAGGAGGTGCTCGATTCTGTTCTGGATCTTGCGAGAAATATCTTCCCTTCCAACCTGGTGTGACGAGCC
TTTCGCTCATACTCTACCACCTATGAAGAGAGGAATATCACCGGAACCAGGGTGAAGGTGCCCCGTGGGGC
AGGAGGTGGAGGGGATGAACATCTGGGCTTGGTAGTGTGGCCATCGTCTTTGGTGTGGCGCTGCGGAA
GCTGGGGCCTGAAGGGGAGCTGCTTATCCGCTTCTTCAACTCCTTCAATGAGGCCACCATGGTTCTGGTC
TCCTGGATCATGTGGTACGCCCCCTGTGGGCATCATGTTCTGGTGGCTGGCAAGATCGTGGAGATGGAGG
ATGTGGGTTTACTCTTTGCCCGCCTTGGAAGTACATTCTGTGCTGCCTGCTGGGTCACGCCATCCATGG
GCTCCTGGTACTGCCCCCTCATCTACTTCTCTTACCCGCAAAAACCCCTACCGCTTCTGTGGGGCATC
GTGACGCCGCTGGCCACTGCCTTTGGGACCTCTTCCAGTTCCGCCACGCTGCCGCTGATGATGAAGTGG
TGGAGGAGAATAATGGCGTGGCCAAGCACATCAGCCGTTTCATCCTGCCCATCGGCGCCACCGTCAACAT
GGACGGTGCCGCGCTCTTCCAGTGCGTGGCCGAGTGTTCATTGCACAGCTCAGCCAGCAGTCTTGGAC
TTTCGTAAAGATCATCACCATCTGGTCACGGCCACAGCTCCAGCGTGGGGGACGCGGCATCCCTGCTG
GAGGTGTCCTCACTCTGGCCATCATCTCGAAGCAGTCAACCTCCCGGTGACCATATCTCCTTGATCCT
GGCTGTGGACTGGCTAGTCGACCGGTCCTGTACCGTCTCAATGTAGAAGGTGACGCTCTGGGGGCAGGA
CTCCTCCAAAATTATGTGGACCGTACGGAGTCGAGAAGCACAGAGCCTGAGTTGATACAAGTGAAGAGTG
AGCTGCCCCCTGGATCCGCTGCCAGTCCCCACTGAGGAAGGAAACCCCTCCTCAAACACTATCGGGGGCC
CGCAGGGGATGCCACGGTCGCCTCTGAGAAGGAATCAGTCATGTAAACCCCGGGAGGGACCTTCCCTGCC
CTGCTGGGGGTGCTCTTTGGACACTGGATTATGAGGAATGGATAAATGGATGAGCTAGGGCTCTGGGGGT
CTGCCCTGCACACTCTGGGGAGCCAGGGGCCCCAGCACCTCCAGGACAGGAGATCTGGGATGCCCTGGCTG
CTGGAGTACATGTGTTTACAAGGGTTACTCTCAAACCCCCAGTTCTCACTCATGTCCCCAACCTCAAGG
CTAGAAAACAGCAAGATGGAGAAATAATGTTCTGCTGCGTCCCCACCGTGACCTGCCCTGGCCTCCCTGT
CTCAGGGAGCAGGTACAGGTCACCATGGGGAATTCAGCCCCACCTGGGGGGATGTTACAACACCATGC
TGGTTATTTTGGCGGCTGTAGTTGTGGGGGGATGTGTGTGTGCACGTGTGTGTGTGTGTGTGTGTGTG
TGTGTGTGTGTCTGTGACCTCCTGTCCCCATGGTACGTCCCACCTGTCCCAGATCCCCATTTCCCTC
CACAATAACAGAAACACTCCCAGGGACTCTGGGGAGAGGCTGAGGACAAATACCCTGCTGCTCACTCCAGAG
GACATTTTTTTTAGCAATAAAATTGAGTGTCAACTATTTTAAAAAAAAAAAAAAAAA

>gi|4507046|ref|NM_003045.1| Homo sapiens solute carrier family 7 (cationic amino acid transporter, y+ system), member 1 (SLC7A1), mRNA
 CGATCCTGCCGGAGCCCCGCCGCCGCCGGCTTGGATTCTGAAACCTTCTTGTATCCCTCCTGAGACATC
 TTTGCTGCAAGATCGAGGCTGTCTCTGGTGAGAAAGTGGTGAGGCTTCCCGTCATATTCAGCTCTGAA
 CAGCAACATGGGGTGCAAAGTCTCTGCTCAACATTGGGCGAGCAGATGCTGCGGCGGAAGGTGGTGGACTGT
 AGCCGGGAGGAGACGCGGCTGTCTCGCTGCCGTGAACACTTTTGATCTGGTGGCCCTCGGGGTGGGCAGCA
 CACTGGGTGCTGGTGTCTACGTCTGGCTGGAGCTGTGGCCCGTGAGAATGCAGGCCCTGCCATTGTCTAT
 CTCCTTCCTGATCGCTGCGCTGGCCTCAGTGCTGGCTGGCTGTGCTATGGCGAGTTTGGTGTCTGGGTC
 CCCAAGACGGGCTCAGCTTACCTCTACAGCTATGTACCGCTTGGAGAGCTCTGGGCCCTTCATACCGGCT
 GGAACCTTAATCCTCTCCTACATCATCGGTACTTCAAGCGTAGCGAGGGCCTGGAGCGCCACCTTCGACGA
 GCTGATAGGCAGACCATCGGGGAGTTCTCACGGACACACATGACTCTGAACGCCCCCGGCTGTGGCT
 GAAAACCCCGACATATTTCGAGTGATCATAATTCTCATCTTGACAGGACTTTTAACCTCTGGTGTGAAAG
 AGTCGGCCATGGTCAACAAAATATTCACTTGTATTAACTGCTCTGGTCTGGGCTTCATAATGGTGTGAGG
 ATTTGTGAAAGGATCGGTTAAAACTGGCAGCTCACGGAGGAGGATTTTGGGAACACATCAGGCCGCTCTC
 TGTTTGAACAATGACACAAAAGAAGGGAAGCCCGTGTTGGTGGATTTCATGCCCTTCGGGTCTCTGGTG
 TCCTGTCTGGGGGAGCGACTTGCTTCTATGCCCTTCGTGGGCTTTGACTGCATCGCCACCACAGGTGAAGA
 GGTGAAGAACCCACAGAAGGCCATCCCCGTGGGGATCGTGGCGTCCCTCTTGATCTGCTTCATCGCTAC
 TTTGGGGTGTCTGGCTGCCCTCACGCTCATGATGCCCTACTTCTGCCTGGACAATAACAGCCCCCTGCCCG
 ACGCCTTTAAGCACGTGGGCTGGGAAGGTGCCAAGTACGAGTGGCCGTGGGCTCCCTCTGCGCTCTTTC
 CGCCAGTCTTCTAGGTTCATGTTTCCCATGCTTCCGCTGGGTTATCTATGCCATGGCTGAGGATGGACTGCTA
 TTTAAATTCCTTAGCCAACGTCAATGATAGGACCAAAACACCAATAATCGCCACATTAGCCTCGGGTGCCG
 TTGCTGCTGTGATGGCTTCCTCTTTGACCTGAAGGACTTGGTGGACCTCATGTCCATTGGCACCTCTCCT
 GGCTTACTCGTTGGTGGCTGCCTGTGTGTTGGTCTTACGGTACCAGCCAGAGCAGCCTAACCTGGTATAC
 CAGATGGCCAGTACTTCCGACGAGTAGATCCAGCAGACCAAAATGAATTGGCAAGCACCAATGATTTCCC
 AGCTGGGGTTTTTACCAGAGGCAGAGATGTTCTCTTTGAAAACCATACTCTACCCAAAACATGGAGCC
 TTCCAAAATCTCTGGGCTAATTGTGAACATTTCAACCAGCCTTATAGCTGTTCTCATCATCACCTTCTGC
 ATTTGTGACCGTCTTTGGAAGGGAGGCTCTCAACAAAGGGCGCTGTGGGCAGTCTTTCTGCTCGCAGGGT
 CTGCCCTCCCTCTGTGCCGTGGTCAAGGGCGCTCATGAGGAGCCCGAGAGCAAGCAAGCTCTCATTT
 TAAGGTTCCCTTCCTGCCAGTGCTCCCCATCCTGAGCATCTTCGTGAACGTCTATCTCATGATGCAGCTG
 GACCAGGGCACCTGGGTCCGGTTTGCTGTGTGGATGCTGATAGGCTTCATCATCTACTTTGGCTATGGCC
 TGTGGCACAGCGAGGAGGCGTCCCTGGATGCCGACCAAGCAAGGACTCCTGACGGCAACTTGGACCAGTG
 CAAGTGACGCACAGCCCCGCCCCCGGAGGTGGCAGCAGCCCCGAGGGACGCCCCAGAGGACCGGGAGG
 CACCCACCCCTCCCCACAGTGCAACAGAAACCACCTGCGTCCACACCCTCACTGCA

>gi|4507048|ref|NM_003046.1| Homo sapiens solute carrier family 7 (cationic amino acid transporter, y+ system), member 2 (SLC7A2), mRNA
 GAATTCGGGCTCTCAAAATTTCTATAGAATCAAGATAGAACCTTTAGATGTCACACCAGAAACTAGCAA
 CTGGAATGAAGATAGAAACAAGTGGTTATAACTCAGACAACTAATTTGTGCGAGGGTTTATTGGAACACC
 TGCCCCACCGGTTTGCGACANAAGTTTCTCCTGTGCGCTTCGTGACAGTCAGAATGATTCCTTGCAGAG
 CCGCGCTGACCTTTGCCCCGATGTCTGATCCGGAGAAAAATCGTGACCTGGACAGTCTAGAAGACACCAA
 ATTATGCCGCTGCTTATCCACCATGGACCTCATTGCCCTGGGCGTTGGAAGCACCTTTGGGGCCGGGGTT
 TATGTCCTCGCTGGGGAGGTGGCCAAGGCAGACTCGGGCCCCAGCATCGTGGTGTCCTTCCTCATTTGCTG
 CCCTGGCTTCAGTGATGGCTGGCCTCTGCTATGCCGAATTTGGGGCCCGTGTTCCTCAAGACGGGGTCTGC
 ATATTTGTACACCTACGTGACTGTGCGAGAGCTGTGGGCCCTTCATCACTGGCTGGAATCTCATTTTATCG
 TATGTGATAGGTACATCAAGTGTTGCAAGAGCCTGGAGTGGCACCTTTGATGAACCTCTTAGCAAAACAGA
 TTGGTCAAGTTTTGAGGACATACTTCAGAATGAATTACACTGGTCTTGCAGAATATCCCGATTTTTTTCG
 TGTGTGCTTATATTACTTCTAGCAGGTCTTTTGTCTTTTGGAGTAAAAGAGTCTGCTTGGGTGAATAAAA
 GTCTTCACAGCTGTTAATATTCTCGTCCTTCTGTTTGTGATGGTTGCTGGGTTTGTGAAAGGAAATGTGG
 CAAACTGGAAGATTAGTGAAGAGTTTCTCAAAAATATATCAGCAAGTGCCAGAGAGCCACCTTCTGAAAA
 CGGAACAAGTATCTATGGGGCTGGTGGCTTTATGCCTTATGGCTTTACGGGAACGTTGGCTGGTGTGCA
 ACTTGCTTTTATGCCTTTGTGGGATTTGACTGCATTGCAACAACTGGTGAAGAAGTTCGGAATCCCCAGA
 AAGCTATTCCCATTGGAATTGTGACGTCTTTGCTTGTGTTGCTTTATGGCCTATTTTGGGGTCTCTGCAGC
 TTTAACACTTATGATGCCGTAACCTCCTCGATGAAAAAGCCCCCTTCTGTAGCGTTTGAATATGTG
 GGATGGGGTCTGCCAAATATGTCGTGCGAGCTGGTTCCTCTGCGCCTTGTCACCAAGTCTTCTGGGCT
 CTATGTTTCTTTACCCCGAATCTGTTTGGCATGGCCCGGGATGGCTTACTGTTTAGATTTCTTGCCAG
 AGTGAGTAAGAGGCAGTACCAGTTGCTGCCACGTTGACTGCAGGGGTCAATTTCTGCTTTGATGGCCTTT

CTGTTTGACCTGAAGGCGCTTGTGGACATGATGTCCATTGGCACAACATCATGGCCTACTCTCTGGTGGCAG
 CCTGTGTTCTCATCCTCAGGTACCAGCCTGGCTTATCTTACGACCAGCCAAATGTTCTCTGAGAAAGA
 TGGTCTGGGATCGTCTCCAGGGTAACCTCGAAGAGTGAGTCCCAGGTCACCATGCTGCAGAGACAGGGC
 TTCAGCATGCGGACCCCTCTTCTGCCCCCTCCCTTCTGCCAACACAGCAGTCAGCTTCTCTCGTGAGCTTTC
 TGGTAGGATTCTTAGCTTTCCCTCGTGTGGGCTGAGTGTCTTGACCACTTACGGAGTTCATGCCATCAC
 CAGGCTGGAGGCCTGGAGCCTCGCTCTCCCTACGCTGTTTCTTGTCTCTTCTGTTGCCATCGTTCTCACC
 ATCTGGAGGCAGCCCCAGAATCAGCAAAAAGTAGCCTTCATGGTTCATTCTTACCATTTTTTGCCAGCGT
 TCAGCATCTTGGTGAACATTTACTTGATGGTCCAGTTAAGTGCAGACACTTGGGTGAGATTTCAGCATTTG
 GATGGCAATTGGCTTCCCTGATTTACTTTTCTTATGGCATTAGACACAGCCTGGAGGGTCATCTGAGAGAT
 GAAAACAATGAAGAAGATGCTTATCAGACAACGTTTCATGCAGCAGCAGAAGAAAAATCTGCCATTCAAG
 CAAATGACCATCACCCAAAGAAATCTCAGTTCACCTTTCATATTCCATGAAAAGACAAGTGAATTCTAACA
 CTTGCAGGAGCAGAT

>gi|31543640|ref|NM_032803.3| Homo sapiens solute carrier family 7
 (cationic amino acid transporter, y+ system), member 3 (SLC7A3), mRNA
 GCCTGAGCGGCCGAACCTCGGCAGCTCCAACCCAACTCGGCTTAACTCCGCCTACCGAGCCAGTCCAAG
 ACTCTGTGCTCCCTAGGTTTGCAACAGCTCTCTGATCATCTTCTTCAATTCTGCTAGGATGCCGTGGCA
 AGCATTTTCGAGATTTGGTCAAAAGCTGGTACGCAGACGTACACTGGAGTCAGGCATGGCTGAGACTCGC
 CTTGCCAGATGCCAAGCACCTGGATTAGTGGCCCTGGGTGTGGGCAGCACATTGGGTGCAGGCGTGT
 ATGTCTTAGCTGGCGAGGTGGCCAAAGATAAAGCAGGGCCATCCATTGTGATCTGCTTTTTGGTGGCTGC
 CCTGTCTTCTGTGTTGGCTGGGCTGTGCTATGCGGAGTTTGGTGGCCGGGTTCCCGTTCTGGTTCGGCA
 TATCTCTACAGCTATGTCACTGTGGGTGAACCTCTGGGCCTTACCCTGGCTGGAACCTCATCCTCTCCT
 ATGTCAATTGGTACAGCCAGTGTGGCCCGGGCCTGGAGCTCTGCTTTTGACAACCTGATTGGGAACCAT
 CTC TAAGACTCTGCAGGGGTCCATTGCACTGCACGTGCCCCATGTCTTGCAGAATATCCAGATTTCTTT
 GCTTTGGGCCCTCGTGTGCTGCTCACTGGATTGTTGGCTCTCGGGGCTAGTGAGTCGGCCCTGGTTACCA
 GAGTGTTCACAGGCGTGAACCTTTTGGTCTTGGGTTCGTATGATCTCTGGCTTCGTTAAGGGGGACGT
 GCACAACCTGGAAGCTACAGAAGAGGACTACGAATTGGCCATGGCTGAACCTCAATGACACCTATAGCTTG
 GGTCTCTGGGCTCTGGAGGATTGTGCTTTTCGGCTTCGAGGAATTCTCCGTGGAGCAGCGACCTGTT
 TCTATGCATTTGTTGGTTTCGACTGTATTGCTACCCTGGAGAGAAGCCAGAAATCCCAAGCGTTCAT
 CCCGATGGGCATTGTGATCTCACTGTCTGTCTGCTTTTGGCGTATTTTGTGTCTCTTCTGCACCTACC
 CTGATGATGCCCTTACTACCAGCTTCAGCCTGAGAGCCCTTTGCCTGAGGCATTTCTCTACATTGGATGGG
 CTCCTGCCCGCTATGTTGTGGCTGTTGGCTCCCTCTGTGCTCTTTCTACCAGCCTCCTGGGCTCCATGTT
 CCCCATGCCCTCGGGTATCTACGCGATGGCAGAGGATGGCCTCCTGTTCGTGTACTTGCTCGGATCCAC
 ACCGGCACACGCACCCCAATCATAGCCACCGTGGTCTCTGGCATTATTGCAGATTTCATGGCATTCCTCT
 TCAAACTCACTGATCTTGTGGACCTCATGTCAATTGGGACCCTGCTTGCTTACTCCCTGGTGTGATTTG
 GTTCTCATCCTCAGGTATCAACCTGATCAGGAGACAAAGACTGGGGAAGAAGTGGAGTTGCAGGAGGAG
 GCAATAACTACTGAATCAGAGAAGTTGACCCTATGGGGACTATTTTCCCACTCAACTCCATCCCCACTC
 CACTCTCTGGCCAAATTTGTCTATGTTTGTTCCTCATTGCTTGCTGTCTGCTGACTGCTCTTTGCCTGGT
 GCTGGCCAGTGGTCAGTTCCATTGCTTTCTGGAGACCTGGTGTGGACTGCAGTGGTTGTGTGCTCCTG
 CTGCTCATTATTGGGATCATTGTGGTCACTCTGGAGACAGCCACAGAGCTCCACTCCCCCTCACTTTAAGG
 TGCCTGCTTTGCCTCTCCTCCCACTAATGAGCATCTTTGTGAATATTTACCTTATGATGCAGATGACAGC
 TGGTACCTGGGCCCCGATTTGGGGTCTGGATGCTGATTGGCTTTGCTATCTACTTCGGCTATGGGATCCAG
 CACAGCCTGGAAGAGATTAAGAGTAACCAACCTCAGCAAGTCTAGAGCCAAACTGTAGACCTTGATC
 CCGGCACTCTCTATGTCCACTCAGTTTGACATCGTACACCTAAATGCTGTCTGGTCCCCCTGCACAATAA
 TGGAGAGTACTCTGACCCAGTGACAGCTAGCCCTCCCTGTGATGGTGGTGGTGGTGGTGGTGGTGGTGGT
 TCTGTACGATGTGAAGGATGTGTCTTTGCTATTTCTTGTCTATTTTAACCCGTCTGCTTCTAAATGATGT
 CTAGCTGCTTACCACTTTAAAAAATGATATTAAAAAGAAAGTAGAAAAATAAAAA
 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

>gi|29789099|ref|NM_018837.1| Homo sapiens similar to glucosamine-6-
 sulfatases (SULF2), mRNA
 GGGCCATTTCTGGACAACAGCTGCTATTTTCACTTGAGCCCAAGTTAATTTCTCGGGGAGTTCTCGGGCG
 CGCACAGGCAGCTCGGTTTGCCCTGCGATTGAGCTGCGGGTCGCGGCCGGCGCCGGCTCTCCAATGGCA
 AATGTGTGTGGCTGGAGGCGAGCGCGAGGCTTTCGGCAAAGGCAGTCGAGTGTTCGAGACCGGGGCGAG
 TCCTGTGAAAGCAGATAAAAGAAAACATTTATTAACGTGTCATTACGAGGGGAGCGCCCGGGCGGGCTG
 TCGCACTCCCCGCGGAACATTTGGCTCCCTCCAGCTCCTAGAGAGGAGAAGAAGAAAGCGGAAAAGAGGC
 AGATTACGTCGTTTCCAGCCAAGTGGACCTGATCGATGGCCCTCCTGAATTTATCACGATATTTGATTT

ATTAGCGATGCCCCCTGGTTTGTGTGTTACGCACACACACGTGCACACAAGGCTCTGGCTCGCTTCCC TC
 CCTCGTTTTCCAGCTCCTGGGCGAATCCCACATCTGTTTCAACTCTCCGCCGAGGGCGAGCAGGAGCGAGA
 GTGTGTCTGAATCTGCGAGTGAAGAGGGACGAGGGAAAAGAAACAAAGCCACAGACGCAACTTGAGACTCC
 CGCATCCCAAAGAAGCACCAGATCAGCAAAAAAGAAAGATGGGCCCCCGAGCCTCGTGCTGTGCTTGC
 TGTCCGCAACTGTGTTCTCCCTGCTGGGTGGAAGCTCGGCCCTCCTGTGCGCACCACCGCCTGAAAGGCAG
 GTTTCAGAGGGACCGCAGGAACATCCGCCCAACATCATCTGGTGCTGACGGACGACCAGGATGTGGAG
 CTGGGTTCATGCGAGGTGATGAACAAGACCCGGCGCATCATGGAGCAGGGCGGGGCGCACTTCATCAACG
 CCTTCGTGACCACACCCATGTGCTGCCCCTCACGCTCCTCCATCCTCACC GGCAAGTACGTCCACAACCA
 CAACACCTACACCAACAATGAGAACTGCTCCTCGCCCTCCTGGCAGGCACAGCACGAGAGCCGCACCTTT
 GCCGTGTACTCAATAGCACTGGCTACCGGACAGCTTCTTCGGGAAGTATCTTAATGAATACAACGGCT
 CCTACGTGCCACCCGGCTGGAAGGAGTGGGTCTGGACTCCTTAAAAACTCCC GCTTTTATAACTACACGCT
 GTGTCTGGAACGGGGTGAAAGAGAAGCACGGCTCCGACTACTCCAAGGATTACCTCACAGACCTCATCACC
 AATGACAGCGTGAGCTTCTTCCGCACGTCCAAGAAGATGTACCCGCACAGGCCAGTCCATGGTCACTCA
 GCCATGCAGCCCCCACGGCCCTGAGGATTACAGCCCCACAATATTACAGCCTCTTCCCAAACGCATCTCA
 GCACATCAGCCGAGCTACAACCTACGCGCCCAACCCGGACAAAACACTGGATCATGCGCTACACGGGGCCC
 ATGAAGCCCATCCACATGGAATTACCAACATGCTCCAGCGGAAGCGCTTGACAGCCCTCATGTGGTGG
 ACGACTCCATGGAGACGATTTACAACATGCTGGTTGAGACGGGCGAGCTGGACAACACGTACATCGTATA
 CACCGCCGACCACGGTTACCACATCGGCCAGTTTGGCCTGGTGAAAGGGAAATCCATGCCATATGAGTTT
 GACATCAGGGTCCC GTTCTACGTGAGGGGCCCCAACGTGGAAGCCGGCTGTCTGAATCCCCACATCGTCC
 TCAACATTGACCTGGCCCCCACCATCTTGACATTGACGGCCTGGACATACCTGCGGATATGGACGGGAA
 ATCCATCCTCAAGCTGCTGGACACGGAGCGGCCGGTGAAATCGGTTTCACTTGAAAAAGAAGATGAGGGTC
 TGGCGGGACTCCTTCTTGGTGAGAGAGGCAAGCTGCTACACAAGAGAGACAATGACAAGGTGGACGCC
 AGGAGGAGAACTTTCTGCCCAAGTACCAGCGTGTGAAGGACCTGTGTGTCAGCGTGTGAGTACCAGACGGC
 GTGTGAGCAGCTGGGACAGAAGTGGCAGTGTGTGGAGGACGCCACGGGGAAGCTGAAGCTGCATAAGTGC
 AAGGGCCCCATGCGGCTGGGCGGCAGCAGAGCCCTCTCCAACCTCGTGCCCAAGTACTACGGGCAGGGCA
 GCGAGGCTTGACCTGTGACAGCGGGGACTACAAGCTCAGCCTGGCCGGACGCCGAAAAAAC TCTTCAA
 GAAGAAGTACAAGGCCAGCTATGTCCGCAGTCGCTCCATCCGCTCAGTGGCCATCAGGTTGGACGGCAGG
 GTGTACCACTAGGCCTGGGTGATGCCGCCAGCCCCGAAACCTCACC AAGCGGCATGGCCAGGGGGCC
 CTGAGGACCAAGATGACAAGGATGGTGGGGACTTCAGTGGCACTGGAGGCCCTTCCGACTACTCAGCCGC
 CAACCCCATTTAAAGTGACACATCGGTGCTACATCCTAGAGAACGACACAGTCCAGTGTGACTTGGACCTG
 TACAAGTCCCTGCAGGCCTGGAAGACCAAGCTGCACATCGACCACGAGATTGAAACCCCTGCAGAACA
 AAATTAAGAACCTGAGGGAAGTCCGAGGTCACTGAAGAAAAGCGGCCAGAAGAATGTGACTGTACAA
 AATCAGCTACCACACCCAGCACAAAGGCCGCTCAAGCACAGAGGCTCCAGTCTGCATCCTTTTCAGGAAG
 GGCCTGCAAGAGAAGGACAAGGTGTGGCTGTTGCGGGAGCAGAAGCGCAAGAAGAAACTCCGCAAGCTGC
 TCAAGCGCTGCAGAACAACGACACGTGCAGCATGCCAGGCCTCACGTGCTTACCCACGACAACCAGCA
 CTGGCAGACGGCGCCCTTCTGAGACTGGGGCCTTTCTGTGCC TGCACCAGCGCCAACAATAACACGTAC
 TGGTGATGAGGACCATCAATGAGACTACAATTTCTCTTCTGTGAATTTGCAACTGGCTTCC TAGAGT
 ACTTTGATCTCAACACAGACCCCCTACCAGCTGATGAATGCAGTGAACACACTGGACAGGGATGTCTCAA
 CCAGCTACACGTACAGCTCATGGAGCTGAGGAGCTGCAAGGGTTACAAGCAGTGTAAACCCCGGACTCGA
 AACATGGACCTGGGACTTAAAGATGGAGGAAGCTATGAGCAATACAGGCAGTTTCAGCGTCGAAAGTGGC
 CAGAAATGAAGAGACCTTCTTCCAAATCACTGGGACAACGTGGGAAGGCTGGGAAGGTTAAGAAACAAC
 AGAGGTGGACCTCCAAAAACATAGAGGCATCACCTGACTGCACAGGCAATGAAAAACCATGTGGGTGATT
 TCCAGCAGACCTGTGCTATTGGCCAGGAGGCCTGAGAAAGCAAGCAGCACTCTCAGTCAACATGACAGA
 TTCTGGAGGATAAACAGCAGGAGCAGAGATAAATTCAGGAAGTCCATTTTGGCCCTGCTTTTGGCTTTGG
 ATTATACCTCACCAGCTGCACAAAATGCATTTTTTTCGTATCAAAAAGTCAACCTAACCTCCCCCAGAA
 GCTCACAAAGGAAAACGGAGAGAGCGAGCGAGAGAGATTTCTTGGAAATTTCTCCCAAGGGCGAAAGTC
 ATTGGAATTTTAAATCATAGGGGAAAAGCAGTCTGTTCTAAATCCTCTTATTCTTTTGGTTTGTGACA
 AAGAAGGAACTAAGAAGCAGGACAGAGGCAACGTGGAGAGGCTGAAAACAGTGCAGAGACGTTTGACAAT
 GAGTCAGTAGCACAAAAGAGATGACATTTACCTAGCATATAAACCCCTGGTTGCCCTCTGAAGAAACTGCCT
 TCATTGTATATATGTGACTATTTACATGTAATCAACATGGGAACCTTTAGGGGAACCTAATAAGAAATCC
 CAATTTTCAGGAGTGGTGGTGTCAATAAACGCTCTGTGGCCAGTGTAAAAGAAAAAAAAAAAAAAAAATGTG
 GACATTTCTGTTCTGTCCAGATACCATTTCTCCTAGTATTTCTTTGTTATGTCCAGAACTGATGTTTT
 TTTTTTAAGGTACTGAAAAGAAATGAAGTTGATGTATGTCCCAAGTTTGTATGAACTGTATTTGTAAAA
 AAAATTTTGTAGTTTAAAGTATTGTATACAGTGTTCAAAACCCAGCCATGACCAGCAGTTGGTATGAA
 GAACCTTTGACATTTTGTAAAAGGCCATTTCTTGGGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
 AAAAAAAAA

```

>gi|29789063|ref|NM_015170.1| Homo sapiens sulfatase 1 (SULF1), mRNA
GGAGTTCTCAGACCTCCAGTTTCAGCCCTGCCCTCAGCCTCCAATCCGTAAGAGACACCCAGCCCCAGCA
ATTGGATTGGGCAGCCCGTCTTGACACACCACTGTGCTGAGTGCTTGAGGACGTGTTTCAACAGATGGTT
GGGGTTAGTGTGTGTCATCACATTTCGAGTGGGGATTAAGAGAAGGAAGGCTGCCTTGCTGGAGCTGTGTG
GTCTTCTCCAAGTGAGAGTCGCAGGCAATAGAACTACTTTTGCTTTTGGAGGAAAAGGAGGAATTCATTTT
CAGCAGACACAAGAAAAGCAGTTTTTTTTTTCAGGGATTCTTCACTTCTCTTGAACAAGGAACCTCACTCAG
AGACTAACACAAAGGAAGTAATTTCTTACCTGGTCATTATTTAGTCTACAATAAGTTTCATCCTTCTTCAG
TGTGACCAGTAAATCTTCCCATAC TCTTGAAGAGAGCATAATTGGAATGGAGAGGTGGTGCTGACGGCC
ACCCACCATCATCTAAAGAAGATAAACTTGGCAAATGACATGCAGGTCTTCAAGGCAGAATAATTGCAG
AAAATCTTCAAAGGACCCATCTGCAGATGTCTGAATACCTCTGAGAATAGAGATTGATTATTCAACCA
GGATACCTAATTCAAGAACTCCAGAAATCAGGAGACGGAGACATTTTGTGCTAGTTTGTCAACATTGGACCA
AATACAATGAAGTATTCTTGCTGTGCTCTGGTTTGGCTGTGCTTGGGCACAGAATTGCTGGGAAGCCCTCT
GTTTCGACTGTGAGATCCCCGAGGTTTCAGAGGACGGATACAGCAGGAACGAAAAAACATCCGACCCAACAT
TATTTCTGTGCC TACCGATGATCAAGATGTGGAGCTGGGGTCCCTGCAAGTCATGAACAAAACGAGAAAG
ATTATGGAACATGGGGGGGCCACCTTCATCAATGCCTTTGTGACTACACCCATGTGCTGCCCCGTACGGT
CCTCCATGCTCACC GGGAAGTATGTGCACAATCACAATGTCTACACCAACAACGAGAAGTGTCTTCCCC
CTCGTGGCAGGCCATGCATGAGCCTCGGACTTTTGTGCTGATATCTTAACAACACTGGCTACAGAACAGCC
TTTTTTGGAAAATACCTCAATGAATATAATGGCAGCTACATCCCCCTGGGTGGCGAGAATGGCTTGAT
TAATCAAGAATTCCTGCTTCTATAATTACACTGTTTGTGCGCAATGGCATCAAAGAAAAGCATGGATTGGA
TTATGCAAAGGACTACTTCACAGACTTAATCACTAACGAGAGCATTAACTTCTCAAATGTCTAAGAGA
ATGTATCCCCATAGGCCCGTTATGATGGTGATCAGCCACGCTGCGCCCCACGGCCCCGAGGACTCAGCCC
CACAGTTTTCTAAACTGTACCCCAATGCTTCCCAACACATAACTCCTAGTTATAACTATGCACCAATAT
GGATAAACACTGGATTATGCAGTACACAGGACCAATGCTGCCCATCCACATGGAATTTACAAACATTTCTA
CAGCGCAAAAGGCTCCAGACTTTGATGTGCTAGTGATGATTCTGTGGAGAGGCTGTATAACATGCTCGTGG
AGACGGGGGAGCTGGAGAATACTTACATCATTTACACCGCCGACCATGGTTACCATATTGGGCAGTTTGG
ACTGGTCAAGGGGAAATCCATGCCATATGACTTTGATATTCTGTGTGCCTTTTTTTTATTCTGTGGTCCAAGT
GTAGAACCAGGATCAATAGTCCCACAGATCGTCTCAACATTGACTTGGCCCCACGATCCTGGATATTG
CTGGGCTCGACACACCTCCTGATGTGGACGGCAAGTCTGTCTCAAACCTCTGGACCCAGAAAAGCCAGG
TAACAGGTTTTCGAACAAAACAAGAGGCCAAAATTTGGCGTGATACATTCTAGTGGAAAGAGGCCAAATTT
CTACGTAAGAAGGAAGAATCCAGCAAGAATATCCAACAGTCAAATCACTTGCCCAATATGAACGGGTCA
AAGAACTATGCCAGCAGGCCAGGTACCAGACAGCCTGTGAACAACCGGGGCAGAAGTGGCAATGCATTGA
GGATACATCTGGCAAGCTTTCGAATTCACAAGTGTAAGGACCCAGTGACCTGCTCACAGTCCGGCAGAGC
ACGCGGAACCTCTACGCTCGCGGCTTCCATGACAAAGACAAAGAGTGCAGTTGTAGGGAGTCTGGTTACC
GTGCCAGCAGAAGCCAAAGAAAGAGTCAACGGCAATTCTTGAGAAACCAGGGGACTCCAAAGTACAAGCC
CAGATTTGTCCATACTCGGCAGACACGTTCTTGTCCGTGCAATTTGAAGGTGAAATATATGACATAAAT
CTGGAAGAAGAAGAAGAAATTGCAAGTGTGCAACCAAGAAACATTGCTAAGCGTCATGATGAAGGCCACA
AGGGGCCAAGAGATCTCCAGGCTTCCAGTGGTGGCAACAGGGGCAGGATGCTGGCAGATAGCAGCAACGC
CGTGGGCCCACCTACCACTGTCCGAGTGACACACAAGTGTTTTATTCTTCCCAATGACTCTATCCATTGT
GAGAGAGAAGTGTACCAATCGGCCAGAGCGTGAAGGACCATAAGGCATACATTGACAAAGAGATTGAAG
CTCTGCAAGATAAAATTAAGAATTTAAGAGAAGTGAGAGGACATCTGAAGAGAAGGAAGCCTGAGGAATG
TAGCTGCAGTAAACAAAGCTATTACAATAAAGAGAAAGGTGTAAAAAAGCAAGAGAAATTAAGAGCCAT
CTTCACCCATTC AAGGAGGCTGCTCAGGAAGTAGATAGCAAACCTGCAACTTTTCAAGGAGAACAACCGTA
GGAGGAAGAAGAGAGGAAGAGAAGAGACGGCAGAGGAAGGGGAAGAGTGCAGCCTGCCTGGCCTCAC
TTGCTTCACGCATGACAACAACCACTGGCAGACAGCCCCGTTCTGGAACCTGGGATCTTTCTGTGCTTGC
ACGAGTTCTAACAATAACACCTACTGGTGTGTTGCGTACAGTTAATGAGACGCATAATTTCTTTCTGTG
AGTTTGCTACTGGCTTTTTTGGAGTATTTTGATATGAATACAGATCCTTATCAGCTCACAAATACAGTGCA
CACGGTAGAACGAGGCATTTTGAATCAGCTACACGTACAAC TAATGGAGCTCAGAAGCTGTCAAGGATAT
AAGCAGTGCAACCCAAGACCTAAGAATCTTGATGTTGGAATAAAGATGGAGGAAGCTATGACCTACACA
GAGGACAGTTATGGGATGGATGGGAAGGTTAATCAGCCCCGTC TCACTGCAGACATCAACTGGCAAGGCC
TAGAGGAGCTACACAGTGTGAATGAAAACATCTATGAGTACAGACAAAAC TACAGACTTAGTCTGGTGGA
CTGGACTAATTACTTGAAGGATTTAGATAGAGTATTTGCAC TGCTGAAGAGTCACTATGAGCAAAAATAAA
ACAAATAAGACTCAAAC TGCTCAAAGTGACGGGTTCTTGGTTGTCTCTGCTGAGCACGCTGTGTCAATGG
AGATGGCCTCTGCTGACTCAGATGAAGACCCAAGGCATAAGGTTGGGAAAACACCTCATTTGACCTTGCC
AGCTGACCTTCAAACCTGCAATTTGAACCGACCAACATTAAGTCCAGAGAGTAAACTTGAATGGAATAAC
GACATTCCAGAAGTTAATCATTGTAATTCGAACTGGAGAAAAACCGAAAAATGGACGGGGCATGAAG
AGACTAATCATCTGGAACCGATTTCAGTGGCGATGGCATGACAGAGCTAGAGCTCGGGCCCAGCCCCAG
GCTGCAGCCCATTTCGAGGCACCCGAAAGAACTTCCCCAGTATGGTGGTCC TGGAAAGGACATTTTTGAA

```


GATCAACTATATCTTCCTGTGCATTCCGATGGAATTTTCAGTTCATCAGATGTTCCACCATGGCCACCGCAG
AACACCGAAGTAATTCAGCATAGCGGGGAAGATGTTGACCAAGGTGGAGAAGAATCACGAAAAGGAGAA
GTCACAGCACCTAGAAGGCAGCGCTCCTCTTCACTCTCCTCTGATTAGATGAACTGTTACCTTACCCT
AAACACAGTATTTCTTTTAACTTTTTATTTGTAACTAATAAAGGTAATCACAGCCACCAACATTCCA
AGCTACCCTGGGTACCTTTGTGCAGTAGAAGCTAGTGAGCATGTGAGCAAGCGGTGTGCACACGGGAGACT
CATCGTTATAATTTACTATCTGCCAAGAGTAGAAAGAAAGGCTGGGGATATTTGGGTGGCTTGGTTTTG
ATTTTTTGCTTGTGTGTGTGTGTGTACTAAAACAGTATTATCTTTTGAATATCGTAGGGACATAAGTAT
ATACATGTTATCCAATCAAGATGGCTAGAATGGTGCCTTTCTGAGTGTCTAAACCTTGACACCCCTGGTA
AATCTTTCAACACACTTCCACTGCCTGCGTAATGAAGTTTGTATTCAATTTTAAACCTGGAATTTTCA
ATGCCGTCAATTTTCAGTTAGATGATTTTGCACCTTTGAGATTAAAATGCCATGTCTATTTGATTAGTCTTA
TTTTTTTATTTTACAGGCTTATCAGTCTCACTGTGGCTGTCTATTGTGACAAAAGTCAAATAAACCCCCA
AGGACGACACACAGTATGGATCAGATATTGTTTGACATTAAGCTTTTGCCAGAAAATGTTGCATGTGTTT
TACCTCGACTTGTCTAAATCGATTAGCAGAAAGGCATGGCTAATAATGTTGGTGGTGAATAAATAAAT
AAGTAAACAAAATGAAGATTGCCTGCTCTCTGTGCCTAGCCTCAAAGCGTTCATCATACATCATACCT
TTAAGATTGCTATATTTTGGGTATTTTCTTGACAGGAGAAAAGATCTAAAGATCTTTTATTTTCATCT
TTTTTGGTTTTCTTGGCATGACTAAGAAGCTTAAATGTTGATAAAATATGACTAGTTTGAATTTACACC
AAGAAGTCTCAATAAAAGAAAATCATGAATGCTCCACAATTTCAACATACCACAAGAGAAGTTAATTTTC
TTAACATTGTGTTCTATGATTATTTGTAAGACCTTCACCAAGTTCTGATATCTTTTAAAGACATAGTTCA
AAATTGCTTTTGAATACTGTATTTCTTGAATAATCCTTGTGTGTATTAGGTTTTTAAATACCAGCTAA
AGGATTACCTCACTGAGTCATCAGTACCCTCCTATTAGCTCCCAAGATGATGTGTTTTGCTTACCCT
AAGAGAGGTTTTCTTCTTATTTTAGATAATTCAAGTGCTTAGATAAATTATGTTTTCTTTAAGTGTTTA
TGGTAACTCTTTTAAAGAAAATTTAATATGTTATAGCTGAATCTTTTGGTAACTTTAAATCTTTATCA
TAGACTCTGTACATATGTTCAAAATAGCTGCTTGCCTGATGTGTGTATCATCGGTGGGATGACAGAAACA
ACATATTTATGATCATGAATAATGTGCTTTGTAAAAAGATTTCAAGTTATTAGGAAGCATACTCTGTTTT
TTAATCATGTATAATATCCATGATACTTTTATAGAACAATTTCTGGCTTCAGGAAAGTCTAGAAGCAATA
TTTCTTCAAATAAAAGGTGTTTAAACTTT

>gi|27597095|ref|NM_173355.1| Homo sapiens uridine phosphorylase-2
(UPase-2), mRNA

GCCGCGCGGCACGGGTGAGTGCCAGGGAATTCACAGTGCTGCCATGCTGGCCCCAGGCTGTGAGTTGG
ACCCAGACCAAGAAGTGGTGAGGACAAGGCCTGAAGATGTGCTGCTTCCCATTCACCTCCACCATGAT
TGTAAGTGTCTGAGGCCCTCCAGCCATGCTTCTGTACAGCCTGTGGAAGTGTGACTTTTACATAGTA
GAGAGAATGGCTTCAGTTATACCTGCCTCCAATAGGTCCATGAGATCTGACAGGAATACATATGTTGGAA
AAAGGTTTGTTCAGGTTAAAAATCCTTACTTGGATTTGATGGATGAAGACATTCTTATCACTTGGATTT
GGGAACAAAACACACAACCTACCAGCAATGTTTGGAGATGTAAAGTTTGTCTGTGTCGGTGGGAGCCCC
AACAGAATGAAAGCATTTGCACTGTTTATGCACAAGGAGCTCGGGTTTGAGGAAGCTGAAGAAGACATAA
AAGACATCTGTGCTGGGACAGACAGATACTGTATGTACAAAACCGGGCTGTGCTCGCCATCAGTCACGG
CATGGGCATCCCCCTCCATTTCTATTATGCTTCATGAACCTCATCAAATTACTCCACCATGCACGGTGTGCTGC
GATGTCACCATTTATAGAATCGGTACATCAGGGGGAATAGGGATTGCACCAGGGACTGTTGTAATAACGG
ATATAGCTGTAGACTCCTTCTTTAAGCCCCGGTTTGAACAGGTCATTTTGGACAACATTGTCACCCGAAG
TACTGAACCTGGACAAAGAACTGTCTGAAGAACTGTTCAACTGTAGCAAAGAAATCCCCAACTTCCCAACC
CTCGTTGGACATACAAATGTGTACCTATGATTTTTATGAAGGCCAAGGCCGACTAGATGGAGCACTGTGCT
CCTTTTCCAGAGAAAAAAGTTAGACTACTTGAAGAGAGCATTTAAAGCTGGTGTGAGGAATATTGAAAT
GGAATCTACAGTGTTCAGCTATGTGTGGACTCTGTGGTCTAAAAGCTGCTGTGGTCTGTGTGACACTT
CTCGACAGACTCGACTGTGATCAGATCAACTTGCTCATGATGTCTGGTGGAGTACCAGCAACGGCCTC
AGCTCCTAATCTCCAACTTCATCAGACGGCGGCTTGGACTTTGTGACTAGACGCTCTAACTGGGCAGCCC
AACCCTCCCCGCAAGTTTGTAGCTCAAGTTGTAATGTGAAAGTCATATTTTATTTGTGGCATTTTATA
TAGTTCTCATCCACATGCTAAATGGAAAGACTTTATGAAATCCTTCTCTCTTAAAAAGGAATTTATTGTA
AAAGAATACTCACACTAAATTAATTTCAAATTTCAATTTTAGAATAAGTTAACTAAATCAGTCTAATAATT
AAAATTTTAAACTCCTGGATTATGACATTTGGAGTTTCATATGCAGCATTAATTAAGCTCACATCAAAA
TAGACCAGCCATTTGCACATAGTTACCAGTCACTCTGTTTCTGAAACCAATCCAGAAATTCATGTTAGAA
CATTGTGTCAGGCATTTCTAGGTATTGGACAAGTGACAGAACTGATCATCCATATTCAAAATATTTACAGA
AAAAAGTGTGAGTGAATGTAGGTCTATATGGAAAAGTTACCATTAGGAAAATATGGCCATACTGCTTTT
TTACCCCTGCACTATTTCACTGACACAGAAATAGGATTCGAGATTTACATATGCCTGTTTGTGATTTTA
TCTCAGAATATAACAACTCTGGATTACATTTTCTCTCAACTGAAAACCTCAGTAATGCAAAAGAAAGGC
CTTCCTTTTCTTCAGAAATCAAGGTGACTAAATTCCTTCTAGCCATTTTCTTGTGAAAAAAGTGTT
TTGAGGAGGCTGTATTTTTTATTCATTTCAATATTAACCAATAGGCTGAGAAATTCATAGCAATTGAA

TTAAGACTTGAGTTCAAGATGTTAAAGGAATCCTTTTGTATGACTTGTCCAGACCGGCCCTATTGCCCTAC
 CCGTTTGATTCTCTGGGAGGCATGGTGCTCCTCCTGGCTGCACATGCTCTCTGTGAAGTGCATCCTT
 GGGTTCCTCTCAATTTGACATACTTGGGAGTCGGAATGACTGGATGGGGGATGGGTAGAGGATGACTGGC
 ACACTTGTATAGTATGTAATGTTGTGTACATATTATTTATATGGTTGTGGCTCAAATAAAGTTTAATTCC
 AACACAAAAAAAAAAAAAAAAA

>gi|31742506|ref|NM_003364.2| Homo sapiens uridine phosphorylase (UP),
 transcript variant 1, mRNA

GGTCAGCTGAGTTCGCCGGCCAGGGCAGGCGGGGCCCCGAGCCTAGCGGTAACCCCCGGGCAGGGCGGGG
 CCGCTCGCAGACTCCATATGAGATTACCTCGCAGGTGGTTCCCTCATTTCGAGTGCCTCCGGCGCACAGAC
 CCGCGCCCCGCGTCTGCGAGCCTCCCGAGAGCCGTCCCTTCGTCCGGCCCTGGAGCATTGCGTTTGTTCG
 CCGGTGTGCGCAGTGCAGGATGGCGCCGCGGGTGTAGCGGCTCTCTGCGCAGGCCGAGTGGGCCAGAGA
 AGCGAGGAACTCCGCAGCTCGTCGACACGTCTCGTCTCCTGTCCCAATTCAGGGCTTGGTGAGGTGACTC
 GCGGTGCGGGTGACTCGCCGGCAGGACACTGCCTGGAACGCCTGGAGCGCCTCCCACTGCAGACGTCTG
 TCCGCCTCCAGCCGCTCTCCTCTGACGGGTCTGCCTCAGTTGGCGGAATGGCGGCCACGGGAGCCAATG
 CAGAGAAAGCTGAAAGTCACAATGATTGCCCCGTCAGACTTTTAAATCCAAACATAGCAAAATGAAAGA
 AGATATTCTCTATCATTCAATCTCACCCTAGCAGACACAATTTCCAGCCTTGGTTTGGAGATGTGAAG
 TTTGTGTGTGTGGTGGAGCCCTCCCGGATGAAAGCCTTCATCAGGTGCGTTGGTGCAGAGCTGGGCC
 TTGACTGCCCAGGTAGAGACTATCCAACATCTGTGCGGGAAGTACCCTATGCCATGTATAAAGTAGG
 ACCGGTGTCTGTCTGTGATCATGGTATGGGCATTCTTCTATCTCAATCATGTTGCATGAGCTCATAAAG
 CTGCTGTACTATGCCCCGGTGTCTCAACGTCACTATCATCCGCATTGGCACTTCTGGTGGGATAGGTCTGG
 AGCCCCGCACTGTGGTCATAACAGAGCAGGCAGTGGATACCTGCTTCAAGGCAGAGTTTGAGCAGATTGT
 CCTGGGGAAGCGGGTCATCCGGAACCGACCTTAACAAGAAGCTGGTGCAGGAGCTGTTGCTGTGTTCT
 GCAGAGCTGAGCGAGTTCACCACAGTGGTGGGGAACACCATGTGCACCTTGGACTTCTATGAAGGGCAAG
 GCCGTCTGGATGGGGCTCTCTGTCTTACACGGAGAAGGACAAGCAGGCGTATCTGGAGGCAGCCTATGC
 AGCCGGCGTCCGCAATATCGAGATGGAGTCTCCTGGTGTGTTGCCGCCATGTGCAGCGCCTGCGGCCCTCAA
 GCGGCCGTGGTGTGTGTACCCCTCCTGAACCCGCTGGAAGGGGACCAGATCAGCAGCCCTCGCAATGTGC
 TCAGCTGATACCAGAGGCGCAGCGGCTGGTGAGCTACTTCAATCAAGAAGAACTGAGCAAGGCTG
 AGCGCTGCCCTGCACCTCCGCAGACCTGCTGTGATGACTTGCCATTAAAGCATTGTCCAAAATCCCTG
 TTGTGTGGACTTTGAGCACACTTTACACAAGAATCTAGAAAATCAGATCGCGATTAAAGAGACAGAGAATC
 TTGGATTAAACGCATGGGAGATGTTCTTCTTTTGAAGTTTCAATTGGAGCATTTTCAATGATGTTAGCCT
 GATTTGGGGTTTCTTCAAGAACATTCTACCAAATTTTGTACTATTTCTAGGGAAATTTTTCAGACTTTA
 AAATTCTAATGGTAGTCAGATTTCTATGTCACTAAACAAGAAATCTGACAATAGTGCCAGGAACTAATTT
 CCTGATACATTAAAAAATTCATGCAAAAAAAAAAAAAAAAAA

>gi|20127508|ref|NM_006834.2| Homo sapiens RAB32, member RAS oncogene
 family (RAB32), mRNA

GGGGGCCGCGAGCACTGGCGGGTTCTGGGTCTGTGACCGGTCAGGCGGCGTCAGCGGGCGCGGCGGAGG
 GCTGGCCGGCCTCGGGGGAGTTTCCGCGGCGCGCGGGGGCGCGGCGGAGAGCGCGAGGCCGGGACCCCG
 GCCAGACTCGGAGTCGAGGCGCGCCGACAGCCGACGCTCATGGCGGGCGGAGGAGCCGGGGACCCCG
 GCCTGGGGGGCGGCCGCGCCCGAGCGCCGAGACCCGCGAGCACCTCTTCAAGGTGCTGGTGATCGGCCGA
 GCTTGGCGTGGGCAAGACCAGCATCATCAAGCGCTACGTCCACCAGCTCTTCTCCAGCACTACCGGGCC
 ACCATCGGGGTGGACTTCGCCC TCAAGGTCTCAACTGGGACAGCAGGACTCTGGTGCGCCCTGCAGCTGT
 GGGACATCGCGGGCAGGAGCGATTGGGCAACATGACCCGAGTATAC TACAAGGAAGCTGTTGGTGCTTT
 TGTAGTCTTTGATATATCAAGAAGTTCCACATTTGAGGCAGTCTTAAAAATGGAAAAGTGATCTGGATAGT
 AAAGTTTATCTTCCAAATGGCAGCCCTATCCCTGCTGTCTCTTGGCTAACAAATGTGACCAGAACAAGG
 ACAGTAGCCAGAGTCTTCCAGGTGGACCAATTTGCAAGAAGCATGGCTTTGCCGGATGGTTTGAAAC
 CTCTGCAAGGATAACATAAATAGAGGAAGCTGCCCCGTTCTTAGTGAGAGAAGATTCTTGTAACCAC
 CAAAGCTTTCTAATGAAGAAAACGATGTGGACAAAATTAAGCTAGATCAAGAGACCTTGAGAGCAGAGA
 ACAAATCCCAGTGTGTGATATATGGCTTCTGCTTCTTGTGTGTGCCTCAGCTCTGAAGAAGTTCCT
 GAGAATGGGTACAGATGTATGTTAGCTGGGAGTCTTCCACATGTGGCACTTCAAAAGGCAGCACCAC
 TGGGCGCCTGCACTTATTTGAAAATGGAACCTTTGGGAGAAGTATCCCTGCTAGTGGCTCTGTAACCTAAC
 AGATGACAATTAGGCTTTTGTGATTGTTGCCATCATATGGAAGATAATGTTTACATCTTTTAAACATTT
 TTATATGACAATTCCTCAGGATTTGGTAAGGCTTCCAAGTTGTAGCTTTTAGTGTAAGTGCTGGGGTGGT
 AATAAAATGTTACCTGCAAAAAAAAAAAAAAAAAA

>gi|11641236|ref|NM_022337.1| Homo sapiens RAB38, member RAS oncogene family (RAB38), mRNA
 GGCTGCGCTTCCCTGGTCAGGCACGGCAGTCTGGCCGCGCCAGGATGCAGGCCCCGCACAAGGAGCA
 CCTGTACAAGTTGCTGGTGATTGGCGACCTGGGCGTGGGGAAGACCAGTATCATCAAGCGCTACGTGCAC
 CAGAACTTCTCCTCGCACTACCGGGCCACAATCGGCGTGGACTTCGCGCTCAAGGTGCTCCACTGGGACC
 CGGAGACTGTGGTGCGCTGCAGCTCTGGGATATCGCAGGTCAAGAAAGATTTGGAACATGACGAGGGT
 CTATTACCGAGAAGCTATGGGTGCATTTATTGTCTTCGATGTCACCAGGCCAGCCACATTTGAAGCAGTG
 GCAAAGTGGAAAAATGATTTGGACTCCAAGTTAAGTCTCCCTAATGGCAAACCGGTTTCAGTGGTTTTGT
 TGGCCAACAAATGTGACCAGGGGAAGGATGTGCTCATGAACAATGGCCTCAAGATGGACCAGTTCTGCAA
 GGAGCACGGTTTCGTAGGATGGTTTGAACATCAGCAAAGGAAAAATATAAACATTGATGAAGCCTCCAGA
 TGCCTGGTGAACACATACTTGCAAATGAGTGTGACCTAATGGAGTCTATTGAGCCGGACGTCGTGAAGC
 CCCATCTCACATCAACCAAGGTGGCAGCTGCTCTGGCTGTGCCAAATCCTAGTAGGCCACTTTGCTGGT
 GTCTGGTAGGAATGACCTCATTTGTTCCACAAATTGTGCCTCTATTTTTTACCATTTTGGGTAAACGTCAGG
 ATAGATATACCACATGTGGCAAGCCAAAGATCTATGCCTCTGTTTTTTCAATGAGAGAGAAATAGCAAAT
 GTTCTTTCTATGCTTTCCTCACCATCATCACAGTGTTTACAAACTTTTGAAAATATTTAGTCTGTTACAA
 ACTTCTGTCTAGTGTGACCAAAATCCTGCAGGGCCACAGTCGGCACTGTTATTTGCTTCTTTTAATCA
 GCAAAGGCCTCAAGTCTTAAAAATAAAGGGGAGAAGAACAACAACTAGCTGTCAAGTCAAGGACTGGCTTTC
 ACCTTGCCCTGGTGTCTTTTCCAGATTCAATATATTCTCTGATGGCCTGACAGGCCTATTAAGTAGAT
 GTGATATTTTCTTCCAAGATGACCTCCATTCTCGGCAGACCTAAGAGTTGCCTCTGAGTTAGCTCTTTGG
 AATCGTGAACACAGGTGTGCTATATTGTCTTGTCTTAAGTGTCACTTGCCATGGCCTGAATGTTGGCTT
 AACTGAATATTGTATGAAAAGACATGCCTCCATATGTGCCTTTCTGTTAGCTCTCTTTGACTCAAGCTGT
 GGGGCTCCTCTATACATGCTATACATGTAATATATATATATATATTTTTTGCAAGTGAACAATAAACAT
 TAAAAGATAAAA

>gi|4506374|ref|NM_003929.1| Homo sapiens RAB7, member RAS oncogene family-like 1 (RAB7L1), mRNA
 CCACACTTCCCGCTCCCTAAAACGCACACCCCGCTAGCCATGGGCAGCCGCGACCACCTGTTCAAAGTG
 CTGGTGGTGGGGGACCCGACAGTGGGCAAGACGTCGCTGGTGCAGCGATATTCCCAGGACAGCTTCAGCA
 AACACTACAAGTCCACGGTGGGAGTGGATTTTGCTCTGAAGTTCTCCAGTGGTCTGACTACGAGATAGT
 GCGGCTTCAGCTGTGGGATATTGCAGGGCAGGAGCGCTTCACCTCTATGACACGATTGTATTATCGGGAT
 GCCTCTGCCTGTGTTATTATGTTTGACGTTACCAATGCCACTACCTTCAGCAACAGCCAGAGGTGGAAAC
 AGGACCTAGACAGCAAGCTCACACTACCCAATGGAGAGCCGGTGGCCTGCCTGCTCTTGGCCAACAAGTG
 TGATCTGTCCCTTGGGCAGTGAGCCGGGACCAGATTGACCGGTTTCAGTAAAGAGAACGGTTTCACAGGT
 TGGACAGAAACATCAGTCAAGGAGAACAAAAATATTAATGAGGCTATGAGAGTCTCATTGAAAAGATGA
 TGAGAAATTCACAGAAGATATCATGTCTTTGTCCACCAAGGGGACTACATCAATCTACAAACCAAGTC
 CTCAGCTGGTCTCTGCTAGTAGTGTGTTGGCTTATTTTCCATCCCAGTTCCTGGGAGGTCTTTTAAGTC
 TCTTCCCTTTGGTTGGCCACCTGACCATTTTATTAAGTACATTTGAATTGTCTCCTGACTACTGTCCAGT
 AAGGAGGCCCATTGTCACTTAGAAAAGACACCTGGAACCCATGTGCATTTCTGCATCTCCTGGATTAGCC
 TTTCACATGTTGCTGACTCACATTAGTGCCAGTTAGTGCCCTTCGGTGTAAAGATCTTCTCATCAGCCCTCA
 ATTTGTGATCCGGAATTTTGTGAGAAGGATTAGAAATCAGCACCTGCGTTTTAGAGATCATAATTCAC
 CTACTTCTGAGCTTATTTTCCATTTGATATTCATTGATATCATGACTTCCAATTGAGAGGAAAATGAGA
 TCAAAATGTCATTTCCCAAATTTCTTGTAGGCCGTTGTTTCAGATTCCTTCTGTCTTGGAAATGTAAACATC
 TGATTCTGGAATGCAGAAGGAGGGGCTGGGCATATGTTGGATTTTGGCTACTAGAAGTGTCCTCAGAAAGT
 CACTGTATTTTGAACCTTCTAACGTCATAAATAAGTTTCTTGTCTTGGCATCAAGAATAGTCAAGTT
 TTTTGGCCGGGCATGGTGGCTCATGCCTGTAATCCCAGCACTTGGGGAGGCCAAGGCAGGCGGATCACAT
 GAGGCCAGGAATTCGAGACCAACCTGGTCAGCATGGCAAACCCCGTCTCTACTAAAAGTACAAAAATTA
 GCCAGGCGTGATGGCACGTGTCTGTAATCCCAGCTACTCTGGAGACTGAGGTGGGAGAATCGCTTGAGAC
 TGGGAGGCAGAGGTTGCAGTGAACCGAGATCATGCCACCGCACTTCAGCCTGGGTGACAGAGAAGGACTC
 CGTCTCAAAAAAAGAAAAAGAATAGTCATTTTAACTACCTATCTCATGCAATGAAAGCATTTTC
 TTCCACAAAGAGCTTAATCCTCATGATAGGATTGCCTAGTGTCTCCCATTTGCAGGTTTCTGGGTTGATG
 TCTTAATGCATAATACTGCAAGTGACATCAGCTGGCTGTGATGCTTCGAAATAGGTCTGCTCCTCACAGC
 TTTGGGAATCTGAATGGAAGAAGAAAAGAGAGAAGTTAACAACCTCCACTGGGGCACTTTGTGAACATG
 TAGGCACTTAGTCATAGGAAACATATTATGTGCAGGTCTTAGCCTGGGGTAGGAAAGTAGATAGACAGAA
 AATCATTAGGTAATTTAAGTACTAAATTGGGCAGGGCTTTTTAGTATCAAATCACTACTAGACCGTTTAA
 TTTGTTAAATTATCTCTAGGATGGTGATTTATAACCTACCCAAAGTTATCGATATTCTTACTAAACTCTG
 AGGCCTGAAGTTCTGTGATAGACCTTAAATAAGTGTCTAAGTCAGTGGTTCCCAAATCTGGCTGGTCGG
 GAATACCTGGGAAGTTTGTAAATTTTTTAAAAATGTTTAAAGATTTTGGGTCTGAGCCAGGCGTGG

TGGCTCACACCTGTAATCCCAGCACTTTGGGAGGCTGAGGCAGGTGGATCGCCTGAGGTCAGGAGTTCAA
 GATCAACCTGGCCAACATACTGAAACCCCGTCTCTACTAAAAATAAGAAAAATTAGCTGGGCGTGGTGGC
 GGGCACCTGTAATCCCAGCTACTTTGGGAGGCTGAGGCAGGAGAATCACTTGAACCTGGGAGTTAGAGGTT
 GCAGTGAGCTGAGATCACACCATTCGCTTCAGCCTGGGCAACAAGAGTGAAACTCCATCTCC

>gi|8923041|ref|NM_017633.1| Homo sapiens chromosome 6 open reading
 frame 37 (C6orf37), mRNA

ACAACCTGCTAAAGCTCCAGAGACACGAGCGTGTGTGGCAGCAAGAGCCGCCAGTTCGGGACCACCGCAGC
 TGGGGTGGCAGCGCGCAGGAGGGGTCTCGGGGGCGGAGTGGTGAGCGCAGGCGGCAGGGGTCTGGGAAA
 GACGAAGTCGCTATTTGCTGTCTGAGCGCGCTCGAGCTCTCGGAAGTGTGCGCGCTCTCGGTTTCGCT
 CTCGCTCGCTGCGCTCTAGAAAGGGCGGCCGCCCTCCAGGACTGACCAGGGCCAAGTGGCGCTCGCGGG
 CACTACATGGCGGAGGGTGAAGGGTACTTCGCCATGTCTGAGGACGAGCTGGCCTGCAGCCCTACATCC
 CCTAGGCGGCGACTTCGGCGGCGGCGACTTCGGCGGCGGCGACTTCGGCGGCGGCGACTTCGGCGGCGG
 CGACTTCGGCGGTGGCGGCAGCTTCGGTGGGCATTGCTTGGAATATTGCGAAAGCCCTACGGCGCACTGC
 AATGTGCTGAACCTGGGAGCAAGTGCAGCGCTGGACGGCATCTGAGTGAGACCATTCGGATTACGGGC
 GCGGCAACTTCCCCACGCTCGAGCTGCAGCCGAGCCTGATCGTGAAGGTGGTGCGGCGGCGCTGGCCGA
 GAAGCGCATTTGGCGTCCGCGACGTGCGCCTCAACGGCTCGGCAGCCAGCCATGTCCTGCACCAGGACAGC
 GGCTTGGGCTACAAGGACCTGGACCTCATCTTCTGCGCCGACCTGCGCGGGGAAGGGGAGTTTCAGACTG
 TGAAGGACGTCCTGTGCTGGACTGCCGTGTTGGACTTCTTACCCGAGGGGGTGAACAAAGAGAAGATCACACC
 ACTCACGCTCAAGGAAGCTTATGTGCAGAAAATGGTTAAAGTGTGCAATGACTCTGACCGATGGAGTCTT
 ATATCCCTGTCAAACAACAGTGGCAAAAATGTGGAAGTGAATTTGTGGATTCCCTCCGGAGGCAGTTTG
 AATTCAGTGTAGATTCTTTTCAAATCAAATTAGACTCTCTTCTGCTCTTTTATGAATGTTTCAGAGAACCC
 AATGACTGAGACATTTACCCCAATAATCGGGGAGAGCGTCTATGGCGATTTCCAGGAAGCCTTTGAT
 CACCTTTGTAACAAGATCATTGCCACCAGGAACCCAGAGGAAATCCGAGGGGGAGGCCTGCTTAAGTACT
 GCAACCTCTTGGTGAAGGGCTTTAGGCCCCGCTCTGATGAAATCAAGGCCCTTCAAAGGTACATGTGTTT
 CAGGTTTTTCATCGACTTCTCAGACATTGGAGAGCAGCAGAGAAAACCTGGAGTCCATTTTGCAGAACCA
 TTTGTGGGATTTGGAAGACCGCAAGTATGAGTATCTCATGACCCCTTCATGGAGTGGTAAATGAGAGCAG
 TGTGCCTGATGGGACATGAAAGAAGACAGACTTTAAACCTTATCACCATGCTGGCTATCCGGGTGTTAGC
 TGACCAAAAATGTCATTCTTAATGTGGCTAATGTCACTTGCTATTACCAGCCAGCCCCCTATGTAGCAGAT
 GCCAACTTTAGCAATTACTACATTGCACAGGTTACGCCAGTATTACGTGCCAGCAACAGACCTACTCCA
 CTTGGCTACCTTGCAATTAAGAATCATTTAAAAATGTCCTGTGGGGAAGCCATTTTCAGACAAGACAGGAG
 AGAAAAAAAAAAAAAAAAAAAAA

>gi|8923191|ref|NM_017709.1| Homo sapiens hypothetical protein FLJ20202
 (FLJ20202), mRNA

TCTCTCACTCGGTGCCGCTGCCTAGGGGCTGTAGAGGTGCGCGCCGCTCCTGCTGGGGCTTGCCACGCCA
 AGGACCTGCCTCTGTGCGCTCCTCTTCTATTGCCAGTTTCCCCAGCCAGAACATCCCCGTAAGATGGCA
 GAGGAGAGCAGCTGTACCAGGGATTGCATGTCTTCAGCGTGCTCAACTGGGATCAGGTTAGCCGGCTGC
 ATGAGGTCTCACTGAAGTTGTACCTATCCACGGACGAGGCAACTTTCCAACCTTGGAGATAACTCTGAA
 GGACATCGTCCAGACCGTCCGCAGTCGGCTGGAGGAGGCAGGCATCAAAGTGCACGACGTCCGGCTGAAT
 GGCTCCGCAGCTGGCCACGTTTGGTCAAAGATAATGGCTTTGGGCTGCAAAGACCTGGACCTAATCTTCC
 ATGTGGCTCTTCCAACAGAGGCAGAATTTAGCTGGTTAGAGATGTGGTTCTGTGTTCCCTTCTGAACCT
 CCTGCCAGAGGGTGTGAACAAGCTCAAAATCAGTCAGTCAGTCTGAAGGAGGCATATGTGCAGAAGCTA
 GTGAAGGTTTGCACGGACACTGTCCGCTGGAGCCTGATCTCCCTCTCCAACAAGAACGGGAAGAAGCTGG
 AGCTGAAGTTTGTGCGACTCCATTCGGCGTCAGTTTGAAGTTAGTGTGGACTCTTTCCAATCATCTGGA
 TTCTTTGCTTTTCTTCTATGACTGTTCCAATAATCCATCTCTGAGCACTTCCACCCACCGTGATTGGG
 GAGAGCATGTACGGGGACTTTGAGGAAGCTTTTGACCATCTGCAGAACAGACTGATCGCCACCAAGAACC
 CAGAAGAAATCAGAGGCGGGGGACTTCTCAAGTACAGCAACCTTCTTGTGCGGGACTTCAGGCCACAGA
 CCAGGAAGAAATCAAACTCTAGAGCGCTACATGTGCTCCAGGTTCTTCATCGACTTCCCGGACATCCTT
 GAACAGCAGAGGAAGTTGGAGACTTACCTTCAAAACCACTTCGCTGAAGAAGAGAGAAGCAAGTACGACT
 ACCTCATGATCCTTCGAGGGTGGTGAACGAGAGCACCGTGTGCTCATGGGGCATGAACGCAGGCAGAC
 TCTGAACCTCATCTCCCTCCTGGCCTTGCCTGTGCTGGCGGAACAAAACATCATCCCCAGTGCCACCAAC
 GTCACCTGTTACTACCAGCCGGCCCCCTTACGTAGTGATGGCAACTTCAGCAACTACTACGTTGCCCATC
 CTCCAGTCACCTACAGCCAGCCTTACCCTACCTGGCTGCCCTGTAACCTTGAAGCTGAGGGTTTC
 CACAGTGGGAACCCCAATAGGGCTAGGGCTCTCAGGTAGGGGAGCCTCCTTCTAGATGTAGGCATTTGGC
 TTTTAAAGGGGAACCTCAGCTCTGATTCTGCTTTTTTTTTTTTTTTTCTTTGTGTACCCATTGGAATGGG
 TCTACAGTGTATCATGAGCCAACCTCAAAGGACCCGTATTACAGTGCCACGTTGGAAAACGCTACAGGA

AGCATGACCTATCCACATCTTTCCAAGATAGACACTAACATGTCATGTCCCAAACATTAGCACGTGGGGG
 TTGAGCTCTGTGCAGTAATCGAGATTGGGAGAAATTTGGGCAGCGCGTGAGAAGTGCTAAGCTACTTGT
 TCTCACTTGAGCCCCGGGTAGGCTGTGTTGGCCCTCAC TTGGGATTCTCAGCAGTTACATGAAAGTTGTGC
 TGATAATCTCTTCTCTGTACCAATTTTAGTCAGGCAGAAAATGGTAAACATGAGGGTGCTCTTGTGACT
 TAATTTTTGTTC AAGGGACTAAATTGCTTATGTTTATTCCTGT CAGCGGAGTGGAGAATGTCATTTCATC
 AATAAACCAAAGCCAATAGCTGGGAGAAATTGAGATCTGGTTGAAAGTGGTTTATGGTTTACATGCTGTACT
 ATCTTGAGGAATTGCCGAGATATTGCTGAGGGGAAAAAAAAAATGACCTTTTCTTGAAATGTAAC TTGAAAA
 CAAAATAAAATGTGGAACATAAAAAAAAAAAAAAAAAAAAAAAAAA

>gi|16418426|ref|NM_052943.1| Homo sapiens hypothetical protein
 MGC16491 (MGC16491), mRNA

GGCACGAGGGTTAGTCGTGCGCGTCTGAGGTGTTCCCTGGCTTTGTCTCGCCGTCGTTGCCGCCCCGTC
 CTGCC TCGTGCCCCCGCCTTGCCCCCTGCCAGCCGCTCTCCCTGTCTCCTCCCTTAACCACCCCC
 ACGGTTTCGCGGTCGGGGCCCGGGGCCGGGCGGATGATGCCGTCGGAGAGCGGAGCTGAGCGCAGGGA
 CCGGGCGGCTGCTCAGGTGGGGACGGCTGCGGCCACGGCGGTGGCCACGGCAGCCCCGGCAGGCGGCGGC
 CCCGACCCGGAGGCTTATCGGCTTCCCCGACGGCACCTGAGTGGGCTGAGCTGGCCACAGGTGAAGC
 GACTGGACGCTCTTCTGAGCGAGCCGATTCCTTTCACGGGCGCGCAACTTCCACGCTGAGCGTGCA
 GCCCCGGCAGATCGTGAGGTGGTCCGAGCACCCTGGAGGAGCAGGGACTACATGTGCACAGTGTGCGG
 CTGCATGGTTACGCTGCCAGCCACGTGCTGCACCTGAGAGTGGCCTGGGC TACAAGGATCTGGACCTGG
 TGTTCGGGTGGACCTGCGCAGTGAGGCATCCTTCCAGCTGACCAAGGCAGTGGTGTGCGCTGCCCTACT
 AGACTTCCCTGCCGCGCGGTGTGAGCCGGGCCAAGATCACGCCACTGACACTCAAGGAGGCATACGTGCAG
 AAGCTGGTGAAAGTGTGCACAGACTCGGACCGCTGGAGCCTCATCTCACTGTCCAACAAGAGCGGCAAGA
 ACGTGAGCTCAAGTTTGTGGACTCGGTGAGACGCCAGTTTGAATTCAGCATAGACTCCTTCCAGATCAT
 CCTGGACTCCCTGTTGCTCTTTGGCCAGTGCTCGTCCACTCCCATGTCTGAGGCCTTCCACCCAACGGTC
 ACAGGCGAAAGCCTGTACGGGGACTTCACCGAGGCCCTGGAGCACCTGCGGCACCGTGTATCGCCACGC
 CGAGTCCCGAGGAGATCCGAGGTGGTGGCCTCCTCAAGTACTGCCACCTCCTGGTGCGGGGCTTCCGGCC
 CCGGCCACGACCGATGTGCGGCCCTGACGCGCTACATGTGCTCCCGCTTCTTCATCGACTTTCCAGAC
 CTGGTGGAGCAGCGGCGCACCTTAGAGCGCTACCTGGAGGCCCACTTCGGTGGGGCAGATGCAGCCCGCC
 GTTACGCCTGCCCTGGTGACACTGCACCGGGTGGTCAACGAGAGCACCGTGTGCC TCATGAACCACGAGCG
 CCGCCAGACGCTGGACCTCATTGCCGCACTGGCGCTGCAGGCAC TGGCTGAGCAGGGCCAGCTGCCACT
 GCCGCCCTGGCCTGGCGCCCTCCAGGCACTGACGGGGTGTGCGCAGCCACTGTCAATTACTACGTGACCC
 CCGTGCAACCTCTCCTGGCTCACGCCTATCCACCTGGCTGCC TTGTAAC TGACTCAGACCTGGCCAGA
 AGGGAAGGGACTGGGCC TCACGGGTGGGGTGGGGCTCCAAGAGTGTGTGGAGGACATGACCAGAGGCG
 CAGAATGTGCCAGGAGGCCAGCACTGCAGGGTTGGGCC TTTGATTGAACAGACCAGACTTTCCCGAGCA
 CGAGGCCCTGTGGGCTTAATGCCAGCCTGGGGCTCTCAGCAGGAGGACCTTTGGTTATTGCATCACAC
 TTTTGAGACCAGCATGCCGGTGGGGTGCAGGCGCCATTTGTCTTGGAAGTCAATTTCTTCAGGAGACAAA
 GCAAGTTTGGGGTGGTAGCCTTAATGCCAGCACCTTGGATTGATTCCCATGGTCTAAGAGGGTCTTTT
 GACTAAGGCCAGCCTGTGGTGATCCAGCAGCTTTGGAGGGCCTGCAGCCACACACTGACTGATACATG
 GTATGTTACAGGACTTGTCACTTTGGATCAGGCTGGGCCACTACAGGGCTGATGGGCCA ACTTGGGTA
 ATCACTGGGAGGAGGAACTGAGGCCCGAGGCAGATACCTGGGTTTGGGGAGGGCTGGGTTTAGCCCGG
 TTCTCTGAACTACATACCCACGCTGTGAAGCCTTCTCTTGAGGGGTGAGGTGGGCAGTATACAGGAG
 CCAAGCAGCTCCAGGGTGGGTCGAGCCAGGCCCTTCCAGGGTGC CACTCCTTTCCCCCTGTGTTGCA
 AGGAGCCCTCTGCCTGGGAGCTTGTCTTTGGAAGGTGCTGAGCCTGAATTTGCCACAGTCAAGTCAAC
 GTTCCACCCCACTGGCTTGGTTACAGGGCTCTACATGGGGTAGGGGAGCCATAC TCCACCCCTTGC
 GTATTCCCTGGGTTTGTGATATTTGCACTCTTAACACCTGCCAATAAAGACGGTCTACACTGAAAAA
 AAAAAAAAAAAAAA

>gi|22749286|ref|NM_152630.1| Homo sapiens hypothetical protein
 MGC26999 (MGC26999), mRNA

AGCGGGAATGTGTGGAGATCAGGTGATGCAGAACTCTGGTAAAGGATGTTAGTCTTCAGAGTAAGACTGT
 TAAATTACAGCAAAGTCTACTATCAGAAGACACGTTTTTCTGGTTGGTGTATTACTTTATTCAGTGATAAT
 ATGAAATTACTATCTTCACTGAAGGATCTACTGAATAATCATTTTAACTCAACTATATTGCATTCTTGA
 AAGCAATCCTTTCAATTGATCTACTGACTTCACAATGTCTGAAATCAGATTACCAATCTCACTTGGGAT
 CAAGTTATAACACTGGATCAAGTGT TAGATGAAGTAATTTCAATTCATGGAAAGGGGAATTTCCCCACAA
 TGGAGGTA AAAACAAAAGACATATTCATGTTGTGAAAGATCAACTCATAGGGCAAGGAATTATTGTTAA
 AGATGCCAGATTGAATGGTTCCGTAGCAAGTTACATAC TTGCAAGCCACAATGGAATCAGCTATAAGGAT
 CTGGACGTTATTTTGGTGTGAGCTTCCAGGTAACGAAGAAATTCAGGTGTTAAAGATGCAGTTCTAG

ACTGTCTACTTGACTTTTTACCAAAAGATGTAAAGAAGGAAAAGCTCTCCCCAGATATCATGAAAGACGC
TTACGTACAGAAATTGGTCAAGGTTTGCAATGGGCATGATTGTTGGAGTCTTATCTCCCTTTCAAATAAC
ACTGGGAAGAATTTAGAACTAAAATTTGTGAGTTCACTCAGACGGCAGTTTGAATTTAGTGTAGATTCCCT
TTCAAATTTGTTTTGGATCCCATGTTAGACTTCTACAGTGACAAAAATGCCAAGCTAACCAAAGAATCCCTA
TCCTGTTGTGGTAGCTGAAAGCATGTATGGAGACTTCCAGGAAGCAATGACACATTTGCAACACAAGCTC
ATATGTACCAGGAAACCTGAAGAGATTAGAGGTGGTGGCCTTCTGAAGTACTGCAGCTTGCTGGTTCATG
GCTTCAAGCCAGCCTGTATGTCAGAAATCAAAAACCTAGAACGTTATATGTGCTCTAGATTCTTTATTGA
TTTTCTCATATAGAAGAACAGCAAAAGAAAATTGAATCATACCTCCACAACCATTTTCATAGGTGAAGGA
ATGACCAAGTATGACTACCTTATGACCTTGCATGGAGTTGTGAATGAAAGCACTGTTTGCCTCATGAGTT
ATGAAAGAAGACAGATTCTCCACCTGATCACCATGATGGCTTTGAAAGTACTTGGAGAATAAATATTCT
ACCCAATACACAAAAGGTAACCTTGCTTTTATCAGCCTGCTCCGTACTTTGCAGCTGAGGCAAGGTACCTT
ATTTATGTAATACCTGAGCCACCCCCCGTTAGCTTCCAGCCATACCACCCACTGCACCTTCGTGGATCAA
ATGGTATGAGTTAAAAAATACACATGCAACCATAGAAACTTTGATTAAATTAAACACCATTTAAAAGCAA
GTTTCCAAATGTAAAATTCAAGATCTGTTTTTATTTTGTACAGTTACCAATTATTTTCAATTACAGTTG
ATGGAATAGTAGTTACCAACTATTTTGTATCAATTAAGCCATCATAGTAAATACAAATATCTATAAACCAA
CCACTTTAAATGTTTTTCAAACGTTATTGACTAGCTATAATAACTTTCAAATGTTGTGTTCCCTTGAAG
TGATTTTCAAATCTTTTGACTACTTGTGACTTTCAAATTTACTGTGACAAATATATTAGAGATGCCTGCA
TCTTTGACTATAACATAAAAGGACAGATTCATGTTTTAAATTAAGATGTACAGTGAAGTATCAAATTTT
TTATATTACAGTCAGTTCCCTTTGTTACATTTAGTTCTTTTTTCTATTGAACAATACCATCAATAAATGTT
CAGTACTTAACAGAAATGCCTAATTTCAAAGCAACAGATTCAGAAGACATTAAACCCCTGGACTTTTCA
AGCATTTTTTTTTTGACAATTAATTTGGGTGGGATACAAAATCCTACTATAGTTTAAAGGAATACTGGAA
AAAAATGGTTCTGGAAAGCCCTGGACATTAGTAATTTGTCTATATAATAAAACATTTAGTTAATTTG
TGGATTTGAATAACAATACCATACAGCTGTGCAACTACTACTGAGATCAGATACAGGGTTTTTGGCCCT
CAAAATAAGTTAAAACGAATGAAAATGACCAGCTCTGAATGTGAAAGCTTCTATCATCATCTACTACAA
AGAGACTCTAAATGGCAAACAGGAAAAAACAGGCAGAGGTTTATAGCCATAGCATTACCATTTTATGG
CTTATTTGAAAAACATCTTTATGTTGAGAAACATTCCATTTTCAGTAAGTTAAAAATTTTCATTGTAAC
AAATGAAGCAGGTTTTTCATTTTGCTTTACCAAGATCAATTTCTAACATGCCACTCAAGTGCCCTCTC
GTGTGAATTTTTTGCGTAACACTATAATTTATTCAACAACACTGTATACCTTTACAGTATATCTAAATATATA
CTTACTACATCGAGTATACCTGTAGAAAAATCTACCAGTGGGATTAAAAATGTATTTTCTTCCCATAAATG
AAAATAATTCATGACCAATATTACCTGAAGTGTGAGAAGAGGAGCTGTCTTAAATTAATATGGATCAGGC
CATTGGAACCTTCAGTATTTCCACTTGGTTATGATTTTTTGTGAGTTTCCATTTTCAGTTTTTATTGACC
ATCAGTTTTTCTATCACAAATGACCTGACAGTTGTATCCCGATAATTTGCCAGTTATAGACTGTGCTTA
TCTTCACTGTATTACAATATTGGACAGAAGTACACCAATACGGCATTTTCAAATAACCAATATTTCTGA
GATGTTTTATACCACAGCACAAGTGGCCATCAGTATATTTTTAAAGAATTATATATCACAGAACCTTGAG
GCTTTAAACACTATGGACTTACCCTTCTTGGACAAAGAATTTGTAATTACAATAAAATATTTAGAAAT
GAAAGGCCAAAAA

>gi|22212711|gb|AAM94374.1|AF497517_1 adenylate cyclase type V [Homo sapiens]
TNAQDQFLKQLVSNVLIFSCTNIVGVCTHYPAEVSQRQAFQETRECIQARLHSQRENQQQERLLLSVLP
RHVAMEMKADINAKQEDMMFHKIYIQKHDNVSILFADIEGFTSLASQCTAQELVMTLNELFARFDKLAEE
NHCLRIKILGDCYYCVSGLPEARADHAHCCVEMGMDMIEAISLVREVTGVNVNMRVGIHSGRVHCGVLGL
RKWQFDVWSNDVTLANHMEAGGKAGRIHITKATLNYLNGDYEVPEPGCGGERNAYLKEHSIETFLILRCTQ
KRKEEKAMIAKMNRQRTNSIGHNPPHWGAERPFYNHLGGNQVSKEMKRMGFEDPKDKNAQESANPEDEV
EFLGRAIDARSIDRLRSEHVRKFLTFREPDLKKYSQVDDRFGAYVACASLVFLFICFVQITIVPHSI
FMLSFYLTCSLLLTLVFVSVIYSCVKLFPSPLQTLRKIVRSKMNSTLVGVFTITLVFLAAAVNMFTCN
SRDLLGLAQEHNISASQVNACHVAESAVNYSLGDEQGFCSPPWPCNFPEYFTYSVLLSLLACSVFLQI
SCIGKLVLMLAIELIYVLIVEVPGVTLFDNADLLVTANAIDFFNNGTSQCPEHATKVLAKVVTPIIISVF
VLALYLHAQQVESTARLDFLWKLQATEEKEEMEELQAYNRRLHNILPKDVAAHFLARERRNDELYYQSC
ECVAVMFASIANFSEFYVELEANNEGVECLRLLENEIADFDEIISEDREFRQLEKIKITIGSTYMAASGLND
STYDKVGKTHIKALADFAMKLMQMKYINEHSFNNFQMKIGLNIGPVVAGVIGARKPQYDIWGNVTNVAS
RMDSTGVPDRIQVTTDMYQVLAANTYQLECRGVVKVKGKGEMMTYFLNGGPPLS

>gi|10181096|ref|NP_056085.1| adenylate cyclase 6 isoform a [Homo sapiens]
MSWFSGLLVPKVDERKTAWGERNGQKRSRRRGTRAGGFCTPRYMSCLRDAEPPSPTPAGPPRCPWQDDAF

IRRGPGKGKELGLRAVALGFEDTEVTTTAGGTAEVAPDAVPRSGRSCWRRVLVQVFQSKQFRSAKLERLY
 QRYFFQMNSSLTLLMAVLVLLTAVLLAFHAAPARPPAYVALLACAAALFVGLMVVNCNRHSFRQDSMWV
 VSYVVLGILAAVQVGGALAADPRSPSAGLWCPVFFVYIAYTLLPIRMRAAVLSGLGLSTLHLILAWQLNR
 GDAFLWKQLGANVLLFLCTNVIGICTHYPAEVSQRQAFQETRGIYIARLHLQHENRQQRLLLSVLPQHV
 AMEMKEDINTKKEDMMFHKIYIQKHDNVSILFADI EGFTSLASQCTAQELVMTLNELFARFDKLAENHC
 LRIKILGDCYYCVSGLPEARADHAHCCVEMGVDMI EAI SLVREVTGVNVNMRVGIHSGRVHCGVLGLRKW
 QFDVWSNDVTLANHMEAGGRAGRIHITRATLQYLNQDYEVPEGRGGERNAYLKEQHI ETFLILGASQKRK
 EEKAMLAKLQRTTRANSMELMPRWVPDRAFSRTKDSKAFRQMGIDDSKDNRGTDALNPEDEVDEFSLR
 AIDARSIDQLRKDHVRRFLLTFQREDLEKKYSRKVDPRFGAYVACALLVFCFCFIQLLI FPHSTLMLGI
 YASIFLLLLITVLICAVYSCGSLFPKALQRLSRISVRSRAHSTAVGIFSVLLVFTSAIANMFTCNHTPIR
 SCAARMLNLT PADITACHLQQLNYSGLDAPLCEGTMPCTCSFPEYFIGNMLLSLASSVFLHIS SIGKLA
 MIFVLGLIYLVLLLLGPPATIFDNYDLLGVHGLASSNETFDGLDCPAAGRVALKYMTPVILLVFAALY
 LHAQQVESTARLDLFWKLQATGEKEEMEELQAYNRRLHNLIPKDVAAHFLARERRNDELYYQSCECVAV
 MFASIANFSEFYVELEANNEGVECLRLNEI IADFDEI ISEERFRQLEKIKTIGSTYMAASGLNASTYDQ
 VGRSHITALADYAMRLMEQMKHINEHSFNNFQMKIGLNMGPVAVGIGARKPQYDIWGNTVNVSSRMDST
 GVPDRIQVTTDLQVLAAGYQLECRGVVVKVGKGEMTTYFLNGGPSS

>gi|4557257|ref|NP_001106.1| adenylate cyclase 8; Adenylyl cyclase-8,
 brain [Homo sapiens]
 MELSDVRCLTGSEELYTIHPTTPAGDGRSASRPQRLWLQTAVRHITTEQRFIHRGGSGSGSGSGSGKASD
 PAGGGPNHHPQLSGDSALPLYSGLPGERAHSTCGTKVFPERSGSGSASGSGGGGDLGLHLDCAPSNSD
 FFLNGGYSYRGVIFPTLRNSFKSRDLERLYQRYFLGQRRKSEVVMNVLDVLTCLLTLVLHL SLASAPMDP
 LKGILLGFFTGI EVVICALVVVRKDTTSHTYLQYSGVVTWVAMTTQILAAGLGYGLLGDGIGYVLTFLFA
 TYSMLPLPLTWAILAGLGTSLLQVILQVVIPLRAVISINQVVAQAVLFMCMTAGIFISYLSDRARQAF
 LETRRCVEARLRLETENQRQERLVS LPRFVLEMINDMTNEVEHLQH QFHRIYIHYENVSI LADV
 KGFTNLSTTLQAQELVRMLNELFARFDRLAHEHHCLRIKILGDCYYCVSGLPEPRQDHAHCCVEMGLSMI
 KTIRYVRSRTKHDVDMRIGIHSGSVLCGVGLRKWQFDVWSWDVDIANKLES GGI PGRIHISKATLDCLN
 GDYNVEEGHGKERNEFLRKHNIETYLIKQPEDSLLSLPEDIVKESVSSSDRRNSGATFTEGSWSPEL PFD
 NIVGKQNTLAALTRNSINLLPNHLAQAALHVQSGPEEINKRI EHTIDLRS GDKLRREHIKPFSLMFKDSSL
 EHKYSQMRDEVFKSNLVCAFI VLLFITAIQSLLPSSRVMPTIQFSILIMLHSAVLITTAEDYKCLPLI
 LRKTCCWINETY LARNV IIFASILINFLGAILNILWCDFDKSIPLKNLTFNSSAVFTDICSYPEYFVFTG
 VLAMVTCAVFLRLNSVLKLA VLLIMIAIYALLTETVYAGLFLRYDNLNHSGEDFLGTKEVSLLLMAMFL
 AVFYHGQQLEYTARLDLWRVQAKEEINEMKELREHNENMLRNI LPSHVARHFLEKDRDNEELYQSQYDA
 VGVMFASIPGFADFSQTEMNQVECLRLNEI IADFDELLGEDRFQDIEKIKTIGSTYMAVSGLSPEK
 QQCEDKWGHLCALADFSALTESIQEINKHSFNNFELRIGISHGSVAVGIGAKKPQYDIWGKTVNLASR
 MDSTGVSGRIQVPEETYLILKDQGFADYRGEIYVKGISEQEGKIKTYFLLGRVQPNPFI LPPRRLPGQY
 SLAAVVLGLVQSLNRQRQKQLLNENNTGIIKGHYNRRTLLSPSGTEPGAQAEGTDKSDLP

>gi|7661962|ref|NP_055585.1| centaurin, gamma 1; centaurin gamma1;
 phosphoinositide 3-kinase enhancer; Arf GAP with GTP-binding protein-
 like, ANK repeat and PH domains 2; GTP-binding and GTPase activating
 protein 2 [Homo sapiens]
 MHAQRQFVVAAVRAEVRREHVAQALNRLRKLAEVDDPELQDSIQASLDSIREAVINSQEWTLRSRIPE
 LRLGLVLDARS GKSSLIHRFLTGSYQVLEKTESEYKKEMLVDGQTHLVLIREEAGAPDAKFSGWADAVI
 FVFSLEDENSFQAVSRLHGLSSLRGEGRGGLALAVGTQDRI SASSPRVVG DARARALCADMKRCSYSE
 TCATYGLNVDRVFQEVAKVVTLRKQQQLLAACKSLPSSPSHSAASTPVAGQASNGGHTSDYSSSLPSSP
 NVGHRELRAEAAAVAGLSTPGSLHRAAKRRTSLFANRRGSDSEKRSLSRGETTGSGRAIPIKQSFL LKR
 SGNLSLNKEWKKKYVTLSSNGFLLYHPSINDYIHSTHGKEMDLLRTTVKVPGRPPRAISAFGPSASINGL
 VKDMSTVQMGEGL EATTPMPSPSPSPSSLPQPPDQTSKHL LKPDRLARALSTDCTPSGDLSPLSREPPP
 SPMVKKQRRKKLTTPSKTEGSAGQAEENFEFLIVSSGTQWHFEAASFEERDAWVQAI ESQILASLQCC
 ESSKVKLR TDSQSEAVAIQAIRNAKGNSICVDCGAPNPTWASLNLGALICIECSGIHRNLGTHLSRVRSL
 DLDDWPRELTLVLTAIGNDTANRVWESDTRGRAPSRDSSREERESWIRAKYEQLLFLAPLSTSEEPLGR
 QLWAAVQAQDVATVLLLLAHARHGPLDTSVEDPQLRSPHLAAELAHVVITQLLLWYGADVAARDAQGR
 ALFYARQAGSQLCADILLQHGCPEGGSAATTPSAATTPSITATPSPRRRSSAASVGRADAPVALV

>gi|7662484|ref|NP_055729.1| centaurin, gamma 2; Arf GAP with GTP-binding protein-like, ANK repeat and PH domains 1; GTP-binding and GTPase-activating protein 1 [Homo sapiens]
 MNYQQQLANSAAIRAEIQRFESVHPNIYSIYELLERVEEPVLQNQIREHVIAIEDAFVNSQEWTLRSRVP
 ELKVGIVGNLASGKSALVHRYLTGTYVQEESEPGRFKKEIVVDGQSYLLIRDEGGPPEAQFAMWVDAV
 IFVFSLEDEISFQTVYHYYSRMANYRNTSEIPLVLVGTQDAISSANPRVIDDARARKLSNDLKRCTYYET
 CATYGLNVERVFQDVAQKIVATRKKQQLSIGPCKSLPNSPSHSSVCSAQVSAVHISQTSNGGGSLSDYSS
 SVPSTPSISQKELRIDVPPTANTPTPVKQSKRRSNLFTSRKGSDDPKKEKGLSRAISIGSGRAIPIKQ
 GMLLKRSGKSLNKEWKKKYVTLCDNGVLTYHPSLHDYMQNVHGKEIDLRTTVKVPGRPPRATSACAPI
 SSPKTNGLSKDMSSLHISPNSDTGLGDSVCSPPSISSTTSPKLDPPPSPHANRKKHRRKKSTSNFKADGL
 SGTAEQEENFEFIIVSLTGQTHFEATTYEERDAWVQAIESQILASLQSCSSKNKSRLTQSSEAMALQ
 SIRNMRGNSHCVCDETQNPWASLNLGALMCIECSGIHRNLGTHLSRVSLDLDDWPIELIKVMSSIGNE
 LANSVWEESQGRTKPSVDSSTREEKERWIRAKYEQLFLAPLPCTELSLGQHLLRATADEDLRTAILLLA
 HGSRDEVNETCGEGDGRALHLACRKGNNVLAQQLLIWYGVDTVARDAGNTALAYARQASSQECIDVLLQ
 YGCPDERFVLMATPNLSRRNNNRNNSGRVPTII

>gi|16799069|ref|NP_114152.2| centaurin, gamma 3; MRIP-1 protein [Homo sapiens]
 MNFQAGGGQSPQQQQLAGGPPQQFALSNSAAIRAEIQRFESVHPNIYAIYDLIERIEDLALQNQIREHV
 ISIEDSFVNSQEWTLRSRVPKVGIVGNLSSGKSALVHRYLTGTYVQEESEPGRFKKEIVVDGQSYLL
 LRDEGGPPELQFAAWVDAVVFVFSLEDEISFQTVYNYFLRLCSFRNASEVPMVLVGTQDAISSANPRVI
 DDSRARKLSTDLKRCTYYETCATYGLNVERVFQDVAQKVVALRKKQQLAIGPCKSLPNSPSHSAVSAASI
 PAVHINQATNGGSAFSDYSSVPSTPSISQRELRIETIAASSTPTPIRKQSKRRSNIFTSRKGADLDRE
 KKAEECKVDSIGSGRAIPIKQGILLKRSGKSLNKEWKKKYVTLCDNGLLTYHPSLHDYMQNIHGKEIDL
 RTTVKVPGRKLRPRATPATAPGTS PRANGLSVERSNTQLGGGTGAPHSASSASLHSEPLSSSAWAGPRPE
 GLHQRSQSVSSADQWSEATSLPPGMQHPASGPAEVLSSSPKLDPPPSPHSNRKKHRRKKSTGTTPRPDGP
 SSATEEAEESEFVVSLSLTGQTHFEASTAEERELWVQSVQAQILASLQGCRAKDKTRLGNQNAALAVQ
 AVRTVRGNSFCIDCAPNPDWASLNLGALMCIECSGIHRHLGAHLRVSLDLDDWPELLAVMTAMGNA
 LANSVWEGALGGYSKPGPDACREEKERWIRAKYEQLFLAPLPSSDVPLGQQLLRAVVEDDLRLVMLLA
 HGSKEEVNETYGDGDGRALHLSSAMANVFTQLLIWYGVDVRSRDARGLTPLAYARRAGSQECADILIQ
 HGCPGEGCGLAPTPNREPANGTNPSAELHRSPSL

>gi|5453840|ref|NP_006377.1| DEAD (Asp-Glu-Ala-Asp) box polypeptide 17 isoform 1; probable RNA-dependent helicase p72 [Homo sapiens]
 MRGGGFGDRDRDRGFGARGGGGLPPKKFGNPGERLRKKKWDLSELPKFEKNFYVEHPEVARLTPYEV
 DELRRKKEITVRGGDVCPKPVFAFHANFPQYVMDVLMQHFTEPTPIQCQGFPLALSGRDMVGIAQTGS
 GKTLAYLLPAIVHINHQPYLERGDGPICLVLAPTRELAQQVQVQVADYGYKCSRLKSTCIYGGAPKGPQIR
 DLERGVEICIAATPGRLIDFLESGKTNLRRCTYLVLEADRMLDMGFEPQIRKIVDQIRPDRQTLMWASATW
 PKEVRQLAEDFLDYTQINVGNELSANHNILQIVDVCMSEKDHKLIQLMEEIMAEKENKTIIFVETKR
 RCDLRLTRMRDGPWAMCIHGDKSQPERDWVLNEFRSGKAPILIATDVASRGLDVEDVKFVINYDYPNSS
 EDYVHRIGRTARSTNKGTYTFFTPGNLQKARELIKVLEEANQAINPKLMQLVDHRRGGGGGGGGRSRYRT
 TSSANNPNLMYQDECRRRLRGVKDGGRRDSASYRDRSETDRAGYANGSGYSPNSAFGAQAGQYTYGQGT
 YGAAAYGTSSYTAQEYAGTYGASSTTSTGRSSQSSSQFSGIGRSGQQPQPLMSQQFAQPPGATNMIGY
 MGQTAYQYPPPPPPPPSRK

>gi|4758138|ref|NP_004387.1| DEAD (Asp-Glu-Ala-Asp) box polypeptide 5; DEAD box-5 [Homo sapiens]
 MSGYSSDRDRGRDRGFGAPRFGGSRAGPLSGKKFGNPGEKLVKKKWNLDLPKFEKNFYQEHPLARRTA
 QEVETYYRSKEITVRGHNCPPVLFNFYEANFPANVMDVIARQNFTEPTAIQAQGWVVALSGLDMVGVAQT
 GSGKTLAYLLPAIVHINHQPFLERGDGPICLVLAPTRELAQQVQVQVAAEYCRACRLKSTCIYGGAPKGPQ
 IRDLERGVEICIAATPGRLIDFLECGKTNLRRCTYLVLEADRMLDMGFEPQIRKIVDQIRPDRQTLMWASATW
 TWPKEVRQLAEDFLKDYIHINIGALELSANHNILQIVDVCHDVEKDEKLIRLMEEIMSEKENKTIIFVET
 KRRCDLRLTRMRDGPWAMCIHGDKSQPERDWVLNEFKHGKAPILIATDVASRGLDVEDVKFVINYDYPN
 SSEDIHHRIGRTARSTNKGTYTFFTPNNIKQVSDILSVLREANQAINPKLLQLVEDRGSGRSGRSGRGMK
 DRRDRYSAGKRGGFNTFRDRENYDRGYSLLKRDFFGAKTQNGVYSAANYTNGSFGSNFVSAGIQTSFRT
 GNPTGTYQNGYDSTQQYGSNVPNMHNGMNQAYAYPATAAAPMIGYPMPTGYSQ

>gi|8922581|ref|NP_060643.1| hypothetical protein FLJ10665 [Homo sapiens]
 MKAFGPPHEGPLQGLVASRIETYGGRHRASAQSTTGRLYPGRYPVLDPSSRRRLQQYVFPFARGSGQARGLS
 PMRLRDPEPEKRHGGHVGAGLLHSPKLKELTKAHELEVLRLHTFSMFGMPRLPPEDRRHWEIGEGGDSGLT
 IEKSWRELVPGHKEMSQELCHQQEALWELLTTELIYVRKLKIMTDLLAAGLLNLQVRGLLMEVSAETLFG
 NVPSLIRTHRSFWDEVLPPTLEETRASGQPLDPIGLQSGFLTFGQRFHPYVQYCLRVKQTMAYAREQQET
 NPLFHAFVQWCEKHKRSRQMLCDLLIKPHQRITKYPLLLHAVLKRSPEARAQEALNAMIEAVESFLRHI
 NGQVRQGEQESLAAAAQRIGPYEVLEPPSDEVEKNLRPFSTLDTSPMLGVASEHTRQLLLEGPVRVKE
 GREGKLDVYLFLFSDVLLVTKPQRKADKAKVIRPPLMLDKLVCQPLRDPNSFLLIHLTEFQCVSSALLVH
 CPSPTDRAQWLEKTQQAQALQKLKAEYVQKRELLTLYRDQDRESPSTRPSTPSLEGSQSSAEGRTPE
 FSTIIPHLVVTEDTDEDAPLVDDTSDSGYGTLPPTPTGSRSPLSRLRQALRRDPRLTFSTLELRDIP
 LRPHPPDPQAPQRRSAPELPEGILKGGSLPQEDPPTWSEEDGASERGNVVVETLHRARLRGQLPSSPTH
 ADSAGESPWESSGEEEEEGPLFLKAGHTSLRPMRAEDMLREIREELASQRIEGAEEPRDSRPRKLTRAQL
 QMRGPHIIQLDTPLSASEV

>gi|29728516|ref|XP_030970.8| similar to hypothetical protein FLJ10665 [Homo sapiens]
 MNSVLTKHGSPPRSWSLCSGTDDQSPAEEKGLRCQNPACMDKGRAAKVCHHADCCQLHRRGPNLCEAC
 DSKFHSTMHYDGHVRFDLPPQGSVLARNVSTRSCPPRTSPAVDLEEEEEESSVDGKGRKSTGLKLSKKK
 ARRRHTDDPSKECFTLKFDLNDIETEIVPAMKKKSLGEVLLPVFERKGIALGKVDIYLDQSNTPSLSTF
 EAYRFGGHYLRVKAPAKPGDEGKVEQGMKDSKSLSLPILRPAGTGPPALERVDAQSRRESLDILAPGRRR
 KNMSEFLGEASIPGQEPPTPSSCSLPSGSSGSTNTGDSWKNRAASRFSGFFSSGPSTSAFGREVDKMEQL
 EGKLHTYSLFGLPRLPRGLRFDHDSWEEYDEDEDENACLRLEDSSWRELIDGHEKLTRRQCHQQEAVWE
 LLHTEASYIRKLRIINFLCCLLNQESGLLCEVEAERLFSNIPEIAQLHRRLLWASVMAVLEKARRTR
 ALLQPGDFLKGFKMFSGSLFKPYIRYCMEEEGCMEYMRGLLRDNDLFRAYITWAEKHPQCQLKLSMDLAK
 PHQRLTKYPLLLKSVLRKTEEPRAKEAVVAMIGSVERFIHVNACMRQRQERQLAAVVSRI DAYEVVES
 SSDEVKLLKEFLHLDLTAPIPGASPEETRQLLLEGLSRMKEGKDSKMDVYCFLLTDLVLVTKAVKKAER
 TRVIRPPLLVDKIVCELRDPGSLFLIYLNEFHSAVGAYTFQASGQALCRGWDTIYNAQNQLQQLRAQE
 PPGSQQLPLQSLLEEEEEEQEEEEEEEEEGEDSGTSAASSPTIMRKSSGSPDSQHCASDGSTETLAMVV
 VEPGDTLSSPEFDSGPFSSQSDETSLSTTASSATPTSELLPLGPVDGRSCSMDSAYGTLSPTSLQDFVAP
 GPMAELVPRAPEPRVPSPPPSPRLRRRTPVQLLSCPPHLLKSKSEASLLQLLAGAGTHGTPSAPSRLS
 ELCLAVPAPGIRTQGSPEAGPSWDCRGAPSPGSGPGLVGCLAGEPAGSHRKRCGLPSGASPRVQPEFP
 PGVSAQHRKLTLAQLYRIRTTLLLNSTLTASEV

>gi|8923496|ref|NP_060334.1| hypothetical protein FLJ20530 [Homo sapiens]
 MAAFLKNVCLGLEDLQYVFMISSHELFITLLKDEERKLLVDQMRKRSRPNLNCIKPVTsfyDIPASASVN
 IGQLEHQLILSVDPWIRIQLIELHGMTSERQFWTVSNKWEVPSVYSGVILGIKDNLTLDLVYILMAKGL
 HCSTVKDFSHAKQLFAACLELVTEFSPKLQVMLNEMLLLDIHTHEAGTGQAGERPPSDLISRVRGYLEM
 RLPDIPLRQVIAEECVAFMLNWRENEYLTQVPAFLQSNPYVKLGQLLAATCKELPGPKESRRTAKDLW
 EVVVQICSVSSQHKRGNDGRVSLIKQRESTLGIMYRSELLSFIKKLREPLVLTIIISLFAKLHNVREDIV
 NDITAEHISIWPSIPNLQSVDFEAVAITVKELVRYTSLINPNHSLWLIQADIYFATNQYSAALHYLQ
 AGAVCSDFFNKAVPPDVYTDQVIKRMKCCSLNCHTQVAILCQFLREIDYKTAFKSLQEQNSHDAMDSY
 YDIWVDVTILEYLTYLHHRGETDKRQIAIKAIGQTELNASNPEEVLQLAAQRRKKKFLQAMAKLYF

>gi|4557601|ref|NP_000798.1| gamma-aminobutyric acid A receptor, alpha 2 precursor [Homo sapiens]
 MKTKLNIYNIEFLLFVFLVWDPARLVLANIQEDEAKNNITIFTRILDRLLDGYDNRLRPLGLGDSITEVFT
 NIYVTSFGPVSDTDMEYTIDVFRQWKDERLKFKGPMNLRNLNLMASKIWTPTDFHNGKKSVAHNMT
 MPNKLRLRIQDDGTLTYMRLTVQAECPMHLEDPFMDAHSCLPKFGSYAYTTSEVTYIWTYNASDSVQVAP
 DGSRLNQYDLLGQSIGKETIKSSTGEYTMHTAHFLKRKIGYFVIQTYLPCIMTVILSQQVFWLNRESVP
 ARTVFGVTTVLTMTTSLISARNSLPKVAYATAMDWFIACVAFVFSALIEFATVNYFTKRGWTWDGKSVV
 NDKKKEKASVMIQNAYAVAVANYAPNLSKDPVLSTISKSATTPEPNKKPENKPAEAKKTfNSVSKIDRM
 SRIVFPVLFGTFNLVYWTYLNREPVLGVSP

>gi|4503861|ref|NP_000801.1| gamma-aminobutyric acid (GABA) A receptor, alpha 5 precursor [Homo sapiens]

MDNGMFSGFIMIKNLLFCISMNLSHFSGFSQMPSTSSVKDETNDNITIFTRILDGLLDGYDNRLRPGLGE
 RITQVRTDIYVTSFGPVSDTEMEYTIDVFRQSWKDERLRFKGPMQRLPLNNLLASKIWTPTDFFHNGKK
 SIAHNMTTPNKLLRLEDDGTLTYTMRLTISAECPMQLEDFPMDAHACPLKFGSYAYPNSEVVYVWTNGST
 KSVVVAEDGSRLNQYHLMGQTVGTENISTSTGEYTIMTAHFHLKRKIGYFVIQTYLPCIMTVILSQVSFW
 LNRESVPARTVFGVTTVLTMTTSLISARNSLPKVAYATAMDWFIACVYAFVFSALIEFATVNYFTKRGWA
 WDGGKALEAAKIKKKREVILNKSTNAFTTGKMSHPNIPKEQTPAGTSNTTSVSVKPSSEKTSSEKKTYN
 SISKIDKMSRIVFPVLFGTFLVYWATYLNREPVIKGAASPK

>gi|4504019|ref|NP_000162.1| glycine receptor, alpha 1 [Homo sapiens]
 MYSFNTLRLYLPGAIVFFSLAASKEAEARSATKPMSPSDFLDKLMGRTSGYDARIRPNFKGPPVNVSCN
 IFINSFGSIAETMTDYRVNIFLRQQWNPRLAYNEYPDSDLDDPSMLDSIWKPDLFFANEKGAFHEIT
 TDNKLRLSRNGNVLYSIRITLTLACPMDLKNFPMQVQTCIMQLESFGYTMNDLIFEWQEQGAVQVADGL
 TLPQFILKEEKDLRYCTKHNTGKFTCIERFHLERQMGGYLIQMYIPSLILVILSWISFWINMDAAPAR
 VGLGITTVLTMTTQSSGSRASLPKVSIVKAIDIMAVCLLFVFSALLEYAAVNFVSRQHKELLRFRRKRR
 HHKEDEAGEGRNFNSAYGMGPACLAQKDGISVKGANNSNTTNPPAPSKSPEEMRKLFIQRAKKIDKISR
 IGFPMAFLIFNMFYWIYKIVRREDVHNQ

>gi|20127457|ref|NP_004121.2| glycogenin [Homo sapiens]
 MTDQAFVTLTNDAYAKGALVLGSSSLKQHRRTTTLVVLATPQVSDSMRKVLETVFDEVIMVDVLDSDGSA
 HLTLMKRPELGVTLTKLHCWSLTQYSKCVFMDADTLVLANIDDLFDREELSAAPDPGWPDCFNNGVVFYQ
 PSVETYNQLLHLASEQGSFDGGDQGIILNTFFSSWATTDIRKHLFFIYNLSSISIIYSYLPFAKVFASAKV
 VHFLGRVKPWNITYDPKTKSVKSEAHDPNMTHPEFLILWWNIFTTNVLPLLQQFGLVKDTCSYVNVLSDL
 VYTLAFSCGFCKEDVSGAISHLSLGEIPAMAQPFVSSEERKERWEQQQADYMGADSFNIDKRLDITYLQ

>gi|5453674|ref|NP_003909.1| glycogenin 2 [Homo sapiens]
 MSETEFHHGAQAGLELLRSSNSPTSASQSAGMTVTDQAFVTLATNDIYCQALVLGQSLRRHRLTRKLVV
 LITPQVSSLLRVILSKVDFEIVNLIDSADYIHLAFLKRPELGLTLTKLHCWTLTHYSKCVFLDADTLV
 LSNVDELFDGEFSAAPDPGWPDCFNNGVVFQPSLHTHKLQLHAMEHGSFDGADQGLNSSFNRWSTT
 DIHKHLFFIYNLSSNTMYTYSFAFKQFGSSAKVHFLGSMKPWNYKYNPQSGSVLEQGSVSSSQHQAFL
 HLWWTVYQNNVLPYKVSQAGEARASPGHTLCHSDVGGPCADSASGVGEPENSTPSAGVPCANSPLGSN
 QPAQGLPEPTQIVDETLSPGRRSEDMIACPETETPAVITCDPLSQSPQPADFTETETILQPAKVES
 VSSEETFEPSELPAEALRDPQLDALEVDLAVSVSISIEEKVKELSPEEERRKWEGRIDYMGKDAFA
 RIQEKLDRLQ

>gi|30147855|ref|XP_301113.1| similar to Glycogenin-1 [Homo sapiens]
 MTDQVFVTLTNDAYTKGALVLGSSSLKQHRRTTTLVVLATPQVSDSMRKVLETVFDEIIVDVLDSDGSA
 HLTLMKRPELAFKVFASAKVVHFLGQVKPWNITYDPKTKSVKNESHHPNVTHPEFLILWWNIFTTTLVLP
 LLQQFGLVKDTFSYVNVENVSGAISHLSLGGIPAMAQSVYPQKNRSGGNRARIWEQIPLTTSRGLT
 LTSSRNTAFSCGHIHFTSLVSDT

>gi|4757726|ref|NP_004632.1| myeloid/lymphoid or mixed-lineage leukemia
 (trithorax (Drosophil; ALL1 fused gene from chromosome 10;
 myeloid/lymphoid or mixed-lineage leukemia (trithorax (Drosophila)
 homolog); translocated to, 10 [Homo sapiens]
 MVSSDRPVSLEDEVSHSMKEMIGGCCVCSDERGAENPLVYCDGHGCSVAHVQACYGIVQVPTGPWFCKR
 CESQERAARVRCELCPHKDGAALKRTDNGGWAHVVCALYIPEVQFANVSTMEPIVLQSVPHDRYNTCTYIC
 DEQGRESKAATGACMTCNKHGCRQAFHVTCQAQFAGLLCEEENGDNDVQYCYCKYHFSKLKSKRGSNR
 SYDQSLSDSSSHSQDKHHEKEKKYKEKDKHKQKHKKQPEPSPALVPSLTVTTEKTYTSTSNNSISGLK
 RLEDTTARFTNANFQEVSAHTSSGKDVSETRGSEGKGGKSSAHSSGQRGRKPGGGRNPGTTVSAASPPFQ
 GFSFGTSGSVKSSSGSSVQSPQDFLSFTDSDLRNDYSYSHSQSSATKDVHKGESGSEGGVNSFSTLIGL
 PSTSAVTSQPKSFENSPGDLGNSSLPTAGYKRAQTSIEETVKEKKRKGKQSKHGPRPKGNKNQENV
 SHLSVSSASPTSSVASAAGSITSSSLQKSPDLLRNGSLQSLSVGSSPVGSEISMQYRHDGACPTTTFSEL
 LNAIHNDRGDSSTLTQELKFIGIYNSNDVAVSFNNVSGSGSSTPVSSSHLPQQSSGHLQQVQALSPSA
 VSSAAPAVATTQANTLSGSSLSQAPSHMYGNRSNSSMAALIAQSENNTQDQDLGDNRLVGRGSSPRGS
 LSPRSPVSSLQIRYDQPGNSSLENLPPVAASIEQLLERQWSEGGQFLLEQGTSDILGMLKSLHQLQVEN
 RRLEEQIKNLTAKKERLQLLNAQLSVFPPTITANPSPSHQIHTFSAQTAPTDTSLNSSKSPHIGNSFLPD
 NSLPVLNQDLTSSGQSTSSSSALSTPPPAGQSPAQQSGVSGVQVNGVTVGALASGMQPVSTIPAVSA

VGGIIGALPGNQLAINGIVGALNGVMQTPVTMSQNPTPLTHTTVPPNATHPMPATLTNSASGLGLLSDQQ
RQILIHQQQFQQLLNSQQLTPVHRHPHFTQLPPTHFSPSMEIMQVRK

>gi|5174577|ref|NP_005928.1| myeloid/lymphoid or mixed-lineage leukemia
(trithorax homolog, Drosophila); translocated to, 6; Myeloid/lymphoid
or mixed-lineage leukemia, translocated to, 6; myeloid/lymphoid or
mixed-lineage leukemia (trithorax (Drosophila) homolog); translocated
to, 6 [Homo sapiens]

MKEMVGGCCVCSDERGWAENPLVYCDGHACSVAVHQACYGIVQVPTGPFWRKCESQERAARVRC ELC PH
KDGALKRTDNGGWAHVVCALYIPEVQFANVLTMEPIVLQYVPHDRFNKTCYICEETGRESKAASGACMTC
NRHGCRQAFHVTCQAQMGALLCEEVLEVDNVKYCYCKYHFSKMKTSRHS SGGGGGAGGGGSGGGGS
GFISGRRSRASPSSTQQEKHPTHHERGQKKSRKDKERLQKHKKRPESPPSILTPPVPTADKVVSSSASS
SSHHEASTQETSESSRESKGGKSSSHSLSHKGGKLLSSGKGVSSFTSASSSSSSSSSSSGGPFQPAVSSLQ
SSPDFSAFPKLEQPEEDKYSKPTAPAPSAPPSPSAPEPPKADLFEQKVVFSGFGPIIMRFSTTTSSSGRAR
APSPGDYKSPHVTGSGASAGTHKMPALSATPVPADETPETGLKEKKHKASKRSRHGPGRPKGSRNKEGT
GGPAAPSLPSAQLAGFTATAASPFGGSLVSSGLGGLSSRTFGPSGSLPSLSLESPLLGAGIYTSNKDPI
SHSGGMLRAVCSTPLSSSLGPPGT SALPRLSRSPFTSTLPSSSASISTTQVFSLAGSTFSLPSTHIFGT
PMGAVNPLLSQAESSHTEPDLEDCSFRCRGTSPQESLSSMSPISSLPALFDQTASAPCGGGLDPAAPGT
TNMEQLLEKQGDGEAGVNIVEMLKALHALQKENQRLQEQILSLTAKKERLQILNVQLSVFPFALPAALPA
ANGPVPGPYGLPPQAGSSDSLSTSKSPPGKSSLGLDNLSTSSDPHSGCPSRSSSSLSFHSTPPPLPLL
QQSPATLPLALPGAPAPLPQPQNGLGRAPGAAGLGAMPAEGLLGGLAGSGGLPLNGLLGGNGAAAPN
PASLSQAGGAPTLQLPGCLNSLTEQQRHLLQQEQQLQQLQQLLASPQLTPEHQTVVYQMIQQIQQKREL
QRLQMAGGSQPLMASLLAGSSTPLLSAGTPGLLPTASAPLLPAGALVAPSLGNNTSLMAAAAAA AVAA
AGGPPVLTATQNPFLSLSGAEGSGGGPKGGTADKGASANQKEG

>gi|4505957|ref|NP_002688.1| POU domain, class 2, transcription factor
1; Octamer-binding transcription factor-1 [Homo sapiens]

MNNPSETSKPSMESGDGNTGTQTNGLDFQKQVPVVGGAISTAQAQAFGLHQLVQLAGTSLQAAAQSLNV
QSKSNEESGDSQQPSQPSQPSQPSQAAIPQTQLMLAGGQITGLTLTPAQQLLLQQAQAQALLAAVQQH
SASQQHSAAGATISASAATPMTQIPLSQPIQIAQDLQQLQQLQQQNLNLQQFVLVHPTTNLQPAQFIISQ
TPQGGQGLLQAQNLQTLQPLQSSQANLLQSQPSITLTSQPATPTRTIAATPIQTLQPSQSTPKRIDTPSLE
EPSDLEEELEQFAKTFKQRRIKLGFTQGDVGLAMGKLYGNDFSQTTISRFEALNLSFKNMCKLKPLLEKWL
NDAENLSSDSSLSPPSALNSPGIEGLSRRRKKRTSIETNIRVALEKSFLNQKPTSEEITMIADQLNMEK
EVIRVWFCNRRQKEKRINPPSSGGTSSSPIKAIFPSPTSLVATTPSLVTSSAATTLTVSPVLPLTSAVT
NLSVTGTSDDTSSNNTATVISTAPPASSAVTSPSLSPSPSASASTSEASSASETSTTQTTSTPLSSPLGTS
QVMVTASGLQTAAAAALQGAQLPANASLAAMAAAAGLNPSLMAQSQAAGGALLSLNPGTLGSLPAL
MSNSTLATIQALASGGSPLPITSLDATGNLVFANAGGAPNIVTAPLFLNPQNL SLLTSNPVSLVSAAAASA
GNSAPVASLHATSTSAESIQNSLFTVASASGAATTTTASKAQ

>gi|4505959|ref|NP_002689.1| POU domain, class 2, transcription factor
2 [Homo sapiens]

MVHSSMGAPEIRMSKPLEAEKQGLDSPSEHTDTERNGPDTNHQNPQNKTS PFSVSPTGPSTKIKAE DPSG
DSAPAAPLPQPAPQPHLPQAQLMLTGSQLAGDIQQLLQQLVLPVGHHLQPPAQFLLPQAQSQPGLLP
TPNLFQLPQQTQGALLTSQPRAGLPTQPPKCLEPPSHPEEPSDLEEELEQFARTFKQRRIKLGFTQGDVGL
AMGKLYGNDFSQTTISRFEALNLSFKNMCKLKPLLEKWL NDAETMSVDSSLPSPNQLSSPSLGFGLPGR
RRKKRTSIETNVRFALEKSFLANQKPTSEEILLIAEQHMEKEVIRVWFCNRRQKEKRINPCSAAPMLPS
PGKPASYS PHMVT PQGAGTLP LSQASSSLSTTVTTLS SAVGTLHPSRTAGGGGGGGGAAPPLNSIPSVT
PPPPATTNSTNPSPQGS HSAIGLSGLNPSTGPGLWWNPAPYQP

>gi|7657409|ref|NP_055167.1| POU transcription factor; likely ortholog
of mouse POU domain, class 2, transcription factor 3 [Homo sapiens]

MVNLESMHTDIKMSGDVADSTDARSTLSQVEPGNDRKGLDFNRQIKTEDLSDSLQQTLSHRPCHLSQGPA
MMSGNQMSGLNASPCQDMASLHPLQQLVLPVGHLLQSVSQFLLSQTQPGQQGLQPNLLPFPQQSGLLLPQ
TGPGLASQAFGHPLPGSSLEPHLEASQHLVPVKHLPSSGGADEPSDLEEELEKFAKTFKQRRIKLGFTQGD
VGLAMGKLYGNDFSQTTISRFEALNLSFKNMCKLKPLLEKWL NDAESSPDPSVSTPSSYPSLSEVFGR
KRKKRTSIETNIRLTLEKRFQDNPKPSSEEISMIAEQLSMEKEVVRVWFCNRRQKEKRINCPVATPIKPP

VYNSRLVSPSGSLGPLSVPPVHSTMPGTVTSSCSPGNNSRPSSPGSGLHASSPTASQNNNSKAAVNSASSF
NSSGSWYRWNHSTYLH

>gi|4506071|ref|NP_002731.1| protein kinase C, iota [Homo sapiens]
MSHTVAGGGSGDHSQVRVKAYYRGDIMTHFEPSISFEGLCNEVRDMCSFDNEQLFTMKWIDEEGDPCT
VSSQLELEEAFLRYELNKDSELLIHVFPCVPERPGMPCPGEDKSIYRRGARRWRKLYCANGHTFQAKRFN
RRAHCAICTDRIWGLGRQGYKCINCKLLVHKKCHKLVTEICGRHSLPQEPVMPMDQSSMHSDDHAQTVIPY
NPSSHESLDQVGEEKEAMNTRESGKASSSLGLQDFDLRVIGRGSYAKVLLVRLKKTDRYAMKVVKEL
VNDDDEDIDWVQTEKHVFEQASNHPFLVGLHSCFQTESRLFFVIEYVNGGDLFMHMQRQRKLP EEHARFYS
AEISLALNYLHERGIIYRDLKLDNVLLDSEGHKLTIDYGMCKEGLRPGDITSTFCGTPNYIAPEILRGED
YGFSDVWWALGVLFMFEMMAGRSPFDIVGSSDNPQNTEDYLFQVILEKQIRIPRSLSVKAASVLKSFLNK
DPKERLGCHPQTGFADIQGHPPFRNVDMMEQKQVVPFFKPNI SGEFGLDNFDSQFTNEPVQLTPDDDD
IVRKIDQSEFEGFEYINPLLSAEECV

>gi|10864650|ref|NP_002735.2| protein kinase C, zeta [Homo sapiens]
MPSRTDPKMEGSGGRVRLKAHYGGDIFITSVDAATTFEELCEEVRDMCRLHQHPLTLKWVDSEGDPC
SSQMELEEAFLRLARQCRDEGLIHVFPSTPEQPLPCPGEDKSIYRRGARRWRKLYRANGHLFQAKRFNR
RAYCGQCSERIWGLARQGYRCINCKLLVHKRCHGLVPLTCRKHMDSVMPSQEPVDDKNEDADLPSEETD
GIAYISSSRKHDSIKDDSEDLKPVIDGMDGKISQGLGLQDFDLIRVIGRGSYAKVLLVRLKKNQDIYAM
KVVKELVHDDDEDIDWVQTEKHVFEQASSNPFVGLHSCFQTTSRLFLVIEYVNGGDLFMHMQRQRKLP
EHARFYAAEICIALNFLHERGIIYRDLKLDNVLLDADGHIKLTIDYGMCKEGLRPGDITSTFCGTPNYIA
EILRGEEYGFSDVWWALGVLFMFEMMAGRSPFDIITDNPDMNTEDYLFQVILEKPIRIPRFLSVKASHVLK
GFLNKDPKERLGCRPQTGFSDIKSHAFFRSIDWDLLEKKQALPPFPQITDDYGLDNFDTQFTSEPVQLT
PDEDAIKRIDQSEFEGFEYINPLLLSTEESV

>gi|4759124|ref|NP_004162.1| solute carrier family 1, member 2;
H.sapiens mRNA for glutamate transporter; glutamate/aspartate
transporter II; excitatory amino acid transporter 2; glial high
affinity glutamate transporter [Homo sapiens]
MASTEGANNMPKQVEVRMPDHLGSEEPKHRHLGLRLCDKLGKNLLLTITVFGVILGAVCGGLRLASPI
HPDVVMLIAFPDILMRMLKMLILPLIISSLTGLSGLDAKASGRLGTRAMVYYMSTTIIAAVLGVILVL
AIHPGNPKLKKQLGPGKKNDVSSSLDAFLDLIRNLFPENLVQACFQQIQTVTKKVLVAPPPDEEANATSA
EVSLNETVTEVPEETKMVIKKGLEFKDGMNVGLIGFFIAFGIAMGKMGDQAKLMVDFFNILNEIVMKL
VIMIMWYSPLGIACLICGKIIAIDLEVVARQLGMYMVTVIIGLIIHGGIFLPLIYFVTRKNPFSLFA
IFQAWITALGTASSAGTLPVTFRCLEENLGIDKRVTFRVLPVGATINMDGTALYEAAVAFIAQMNGVVL
DGGQIVTVSLTATLASVGAASIPSAGLVMTLLILTAVGLPTEDISLLVAVDWLLDRMRTSVNVVGD
SFGAGIVYHLSKSELDTIDSQHRVHEDIEMTKTQSIYDDMKNHRESNSNQCVYAAHNSVIVDECKVT
LAANGKSADCSVEEEPWKREK

>gi|4759126|ref|NP_004163.1| solute carrier family 1 (glial high
affinity glutamate transporter), member 3 [Homo sapiens]
MTKSNGEPPKMGRMERFQQGVKRRTLLAKKKVQNTKEDVKSYLFRNAFVLLTVTAVIVGTILGFTLRP
YRMSYREVKYFSFPGELLMRMLQMLVPLIISSLVGTMAALDSKASGKMGMRAVYYMTTIIAVVIGII
IVIIHHPGKGTENMHREGKIVRVTAADAFDLIRNMFPPNLVEACFKQFKTNYEKRSFKVPIQANETLV
GAVINNVSEAMETLTRITEELVPVPGSVNGVNALGLVVFMCFGFVIGNMKEQQQALREFFDSLNEAIMR
LVAVIMWYAPVGILFLIAGKIVEMEDMGVIGQLAMYTVTIVIGLLIHAVIVLPLLYFLVTRKNPWVFIG
GLLQALITALGTSSSSATLPITFKCLEENNGVDKRVTRFVLPVGATINMDGTALYEALAAIFIAQVNNFE
LNFGQIITISITATAASIGAAGIPQAGLVMTMIVLTSVGLPTDDITLIIAVDWFLDRLRTTTNVLGDSL
GAGIVEHLRHELKNRDVEMGNSVIEENEMKKPYQLIAQDNETEKPIDSETKM

>gi|20070239|ref|NP_006662.2| solute carrier family 1 (glutamate
transporter), member 7; excitatory amino acid transporter 5 (retinal
glutamate transporter) [Homo sapiens]
MVPHAILARGRDVCRNGLLILSVLSVIVGCLLGFFLRTRRLSPQEISYFQFPGELLMRMLKMMILPLVV
SSLMSGLASLDAKTSSRLGVLTVAYYLWTTFMVAVIVGIFMVSIIHPGSAAQKETTEQSGKPISSADALL
DLIRNMFANLVEATFKQYRTKTPVVKSPKVAPEEAPPRRILYGVQEENGSHVQNFALDLTPPEVVY
KSEPGTSDGMNVLGIVFFSATMGIMLGRMGDSGAPLVSFQCCLNESVMKIVAVAVWYFPFGIVFLIAGKI

LEMDDPRAVGKKLGFSVTVVCGLVLHGLF ILPLLYFFITKKNPIVFIRGILQALLIALATSSSSATLPI
TFKCLLENNHIDRRIARFVLPVGATINMDGTALYEAVAAIFIAQVNNYELDFGQIITISITATAASIGAA
GIPQAGLVTMVIVLTSVGLPTDDITLIIAVDWALDRFRMTINVLGDALAAGIMAHICRKDFARDTGTEKL
LPCETKPVSLQEIVAAQQNGCVKSVAEASELTGPTCPHHVPVQVERDEELPAASLNHCTIQISELETNV

>gi|21314632|ref|NP_003029.2| solute carrier family 1, member 4;
glutamate/neutral amino acid transporter;
alanine/serine/cysteine/threonine transporter [Homo sapiens]
MEKSNETNGYLDQAQAGPAAGPGAPGTAAGRARRCAGFLRRQALVLLTVSGVLGAGLGAALRGLSLSRT
QVTYLAFPGEMLLRMLRMIIILPLVVC SLVSGAASLDASCLGRGGIAVAYFGLTTLASALAVAFI IK
PGSGAQT LQSSDLGLED SGPPVPKETVDSFLDLARNLFPSNLVVA AFRTYATDYKVVTQNSSSGNVTHE
KIPIGTEIEGMNIGLVLFALVLGVALKKLGSEGEDLIRFFNSLNEATMVLVSWIMWYVPVGIMFLVGSK
IVEMKDIIVLVTS LGKYIFASILGHVHGGIVLPLIYFVFTRKNPFRLLGLLAPFATAFATC SSSATLP
SMMKCIEENNGVDKRISR FILPIGATVNMDGAAIFQCVA AVFIAQLNNVELNAGQIFTILVTATASSVGA
AGVPAGGVLTIAIILEAIGLPTHDLPLILAVDWIVDRTTTVNVEGDALGAGILHHLNQKATKKGEQELA
EVKVEAIPNCKSEETSPLVTHQNPA GPVASAPELESKESVL

>gi|4827012|ref|NP_005062.1| solute carrier family 1 (high affinity
aspartate/glutamate transporter), member 6; excitatory amino acid
transporter 4 [Homo sapiens]
MSSHGNSLFLRESGQRLGRVGWLQRLQESLQQRALRTRLRLQTMTEHVLRFLRRNAFILLTVSAVVIGV
SLAFALRPYQLTYRQIKYFSFPGELLMRMLQMLVLP LIVSSLV TGMASLDNKATGRMGMAAVVYVMVTI
IAVFIGILMVTI IHPGKGSKEGLHREGRIETIPTADAFMDLIRNMFPPNLVEACFKQFKTQYSTRVVRT
MVRTENGSEPGASMPPPFSVENGT SFLENVTRALGTLQEMLSFEETVPVPGSANGINALGLVVF SVAFG
VIGGMKHKGRVLRDFDSLNEAIMRLVGII IYAPVGILFLIAGKILEMEDMAVLGGQLGMYTLTVIVGL
FLHAGIVLPLIYFLVTHRNPFF IGGMLQALITAMGTSSSSATLPITFRCLEEGLGVDRRITRFVLPVGA
TVNMDGTALYEALAAIFIAQVNNYELNLGQIT TISITATAASVGAAGIPQAGLVTMVIVLTSVGLPTEDI
TLIIAVDWFLDRLRTMTNVLGDSIGAAVIEHLSQRELELQEAELTLP SLGPKPYKSLMAQEKGASRGRGGN
ESAM

>gi|18252049|ref|NP_004161.2| solute carrier family 1, member 1;
excitatory amino acid transporter-3; excitatory amino acid carrier 1
[Homo sapiens]
MGKPAKGC EWKRFLKNWVLLSTVA AVVLGITTGVLVREHSNLSTLEKFYFAFPGEILMRMLKLIILPL
IISMITGVAALDSNVSGKIGLRAVYYYFCTTLIAVILGIVLVVSIKPGVTQKVGEIARTGSTPEVSTVD
AMLDLIRNMFENLVQACFQYKTKREEVKPPSDPEMNMTESFTAVMTT AISKNKTKEYKIVGMYSDGI
NVGLIVFCLVFLVIGKMGEGQILVDFFNALSDATMKIVQIIMCYMPLGILFLIAGKIEVEDWEIFR
KLGlyMATVLTGLAIHSIVILPLIYFIVVRKNPFRFAMGMAQALLTALMISSSSATLPVTFRC AEENNQV
DKRITRFVLPVGATINMDGTALYEAVAAV FIAQLNDLDLGIGQIITISITATSASIGAAGVPQAGLVTMV
IVLSAVGLPAEDVTLIIAVDWLDRFRMTNVLGD AFGTGIVEKLSKKELEQMDVSSEVNIVNPFAL EST
ILDNEDSDTKKSYVNGGFAVDKSDTISFTQTSQF

>gi|5032093|ref|NP_005619.1| solute carrier family 1 (neutral amino
acid transporter), member 5; neutral amino acid transporter B; RD114
virus receptor; baboon M7 virus receptor [Homo sapiens]
MVADPPRDSKGLAAA EPTANGGLALASIEDQGAAGGYCGSRDQVRRCLRANLLVLLTVVAVVAGVALGL
GVSGAGGALALGPERLSAFVFP GELLRLRLRMIIILPLVVC SLIGGAASLDPGALGR LGAWALLFFLVTTL
LASALGVGLALALQPGAASAAINASVGAAGSAENAPSKEVLDSFLDLARNIFPSNLVSAAFRSYSTTYEE
RNITGTRVKVPVGQEV EGMNIGLVVFAIVFGVALRKLGP EGELLIRFFNSFNEATMVLVSWIMWYAPVG
IMFLVAGKIVEMEDVGLLFARLGKYILCCLLGHAHGLLVLP LIYFLFTRKNPYRFLWGI VTPLATAFGT
SSSSATLP LMMKCVEENNGVAKHISR FILPIGATVNMDGAALFQCVA AVFIAQLSQQSLDFVKIITILVT
ATASSVGAAGIPAGGVLTIAIILEAVNLPVDHISLILAVDWLVDRSCTVLNVEGDALGAGLLQNYVDRTE
SRSTEPELIQVKSELPLDPLPVPTEEGNPLLKHYRGPAGDATVASEKESVM

>gi|4507047|ref|NP_003036.1| solute carrier family 7 (cationic amino
acid transporter, y+ system), member 1; amino acid transporter,
cationic 1; ecotropic retroviral receptor [Homo sapiens]

MGCKVLLNIGQQMLRRKVDCSREETRLSRCINTFDLVALGVGSTLGAGVYVLAVARENAGPAIVISF
LIAALASVLAGLCYGEFGARVPKTGSAYLYSYVTGELWAFITGWNILISYIIGTSSVARAWSATFDELI
GRPIGEFSRTHMTLNA PGVLAENPDIFAVIIILITGLLT LGVKESAMVNKIFTCINVLVLGFIMVSGFV
KGSVKWNQLTEEDFGNTSGRLCLNNDTKEGKPGVGGFMPFGFSGVLSGAATCFYAFVGFDCIATTGEEVK
NPQKAIPVGIVASLLICFIAYFGVSAALTLMMPYFCLDNNSPLPDAFKHVGWEGAKYAVAVGSLCALSAS
LLGSMFPMPRVIYAMAEDGLLFKFLANVNDRTKTPIIATLASGAVAAMAFLEFLLKDLVDLMSIGTLLAY
SLVAACVLVRLRYQPEQPNLVYQMASTSDLELDPADQNELASTNDSQLGFLPEAEMFSLKTILSPKNMEPSK
ISGLIVNISTSLIAVLIITFCIVTVLGREALTKGALWAVFLLAGSALLCAVVTGVIWRQPESKTKLSFKV
PFLPVLPLSIFVNVYLMQLDQGTWVRFAVWMLIGFIIYFGYGLWHSEEASLDADQARTPDGNLDQCK

>gi|4507049|ref|NP_003037.1| solute carrier family 7 (cationic amino acid transporter, y+ system), member 2; amino acid transporter, cationic 2 [Homo sapiens]

MIPCRAALT FARCLIRRKIVTLDSEDTKLCRCLSTMDLIALGVGSTLGAGVYVLAGEVAKADSGPSIVV
SFLIAALASVMAGLCYAEFGARVPKTGSAYLYTYVTGELWAFITGWNILISYVIGTSSVARAWSGTDFDE
LLSKQIQGFLRTYFRMNYTGLAEYPDFFAVCLILLLAGLLSFGVKESAWVNKVF TAVNILLVLLFVMVAGF
VKGNVANWKISSEFLKNISASAREPPSENGTSIYGAGGFMPYGF TGTLAGAATCFYAFVGFDCIATTGEE
VRNPQKAIPIGIVTSLVCFMAYFGVSAALTLMMPYLLDEKSPLPVAFEYVGWGPakyVVAAGSLCALS
TSLGSMFPLPRILFAMARDGLLFRFLARVSKRQSPVAATLTAGVISALMAFLFDLKLALVDMMSIGTLMA
YSLVAACVLILRYQPGLSYDQPKCSPEKDGLSSPRVTSKSESQVTMLQRQGF SMRTLFCPSLLPTQQSA
SLVSFLVGFLAFLVLGLSVLT TYGVHAITRLEAWSLALLTLFLVLFAI VLT IWRQPQNQQKVA FMVPFL
PFLPAFSILVNIYLMVQLSADTWVRF SIWMAIGFLIYFSYGI RSHSLEGLHLDENNEEDAYPDNVHAAAE
KSAIQANDHHPNRLSSPFIFHEKTSEF

>gi|17939406|ref|NP_116192.2| solute carrier family 7 (cationic amino acid transporter, y+ system), member 3 [Homo sapiens]

MPWQAFRRFGQKLVRRTLESGMAETRLARCLSTDLVALGVGSTLGAGVYVLAGEVAKDKAGPSIVICF
LVAALSSVLAGLCYAEFGARVPRSGSAYLYSYVTGELWAFITGWNILISYVIGTASVARAWSSAFDNLI
GNHISKTLQGSIALHVPVHLAEYPDFFALGLVLLLTGLLALGASESALVTKVFTGVNLLVLGVFMISGFV
KGDVHNWKLTEEDYELAMAE LNDTYSGLPLGSGGFVPFGFEGILRGAATCFYAFVGFDCIATTGEEAQN
QRSIPMGIVISLSVCFLAYFAVSSALTLMMPYQLQPE SPLPEAFLYIGWAPARYVVAAGSLCALSTSL
GSMFPMPRVIYAMAEDGLLFRVLARIHTGTRTPIIATVVSIIAFAFLFKLTDLVDLMSIGTLLAYSL
VSICVLILRYQPDQETKTGEEVELQEEAITTESEKLTWGLFFPLNSIPTPLSGQIVYVCSSLLAVLLTA
LCLVLAQWSVPLLSGDLWTAVVVL LLLLIIGIIVVIWRQPQSSTPLHFKVPALPLPLMSIFVNIYLM
QMTAGTWARFGVWMLIGFAIYFGYGIQHSLEEIKSNQPSRKSRAKTVDLDPGTLYVHSV

>gi|29789100|ref|NP_061325.1| similar to glucosamine-6-sulfatases; sulfatase 2 [Homo sapiens]

MGPPSLVLCLLSATVFSLLGGSSAFLSHHRLKGRFQRDRRNIRPNIIILVLTDDQDVELGSMQVMNKTRRI
MEQGGAHFINAFVTPMCCPSRSSILT GKYVHNHNTYTNNENCSSPSWQAQHESTRFAVYLNSTGYRTAF
FGKYLNEYNGSYVPPGWEKVGLLKNSRFYNYTLCRNGVKEKHGSDYSKDYLTDLITNDSVSFFRTSKM
YPHRPVLMVISHAAPHGPEDSAPQYSRLFPNASQHITPSYNYAPNPDKHWIMRYTGPMKPIHMEFTNMLQ
RKRLQTLMSVDDSMETIYNMLVETGELDNTYIVYTADHG YHIGQFGLVKGKSMPYEFDIRVPFYVRGPNV
EAGCLNPHIVLNIDLAPTILDIAGLDIPADM DGKSILKLLDTERPVNRFHLK KKMVRWRDSFLVERGKLL
HKRDNDKVDAQEENFLPKYQVRKDLQRAEYQTACEQLGQKWQCVEDATGKLKLHKCKGPMRLGGSRLS
NLVPKYYGQGEACTCDSGDYKLSLAGRRKKLFKKKYKASYVRSRSIRSVAIEVDGRVYHVLGDAAQPR
NLTKRHWP GAPEDQDDKDGGDFSGTGGLPDYSAANPIKVTHRCYILENDTVQCDLDLYKSLQAWKDHKLH
IDHEIETLQNKIKNLEVRGHLKKRPEECDCHKISYHTQHKGRLKHRGSSLHPRKGLQEKDKVWLLRE
QKRKKKLRKLLKRLQNNDTCSMPGLTCFTHDNQHWQTAPFWTLGPFCACTSANNTYWC MRTINETHNFL
FCEFATGFLEYFDLNTDPYQLMNAVNTLDRDVLNQLHVQLMELRSCKGYKQCNPRTRNMDLGLKDGGSYE
QYRQFQRRKWPEMKRPSSKSLGQLWEGWEG

>gi|29789064|ref|NP_055985.1| sulfatase FP [Homo sapiens]

MKYSCCALVLAVLTELGLSCLSTVRSRFRGRIGQERKNIRPNIIILVPTDDQDVELGSLQVMNKTRKIM
EHGGATFINAFVTPMCCPSRSSMLTGKYVHNHNTYTNNENCSSPSWQAMHEPRTFAYVYLNNTGYRTAFF
GKYLNEYNGSYIPPGWREWLGLIKNSRFYNYTVCRNGIKEKHGFDYAKDYFTDLITNESINYFKMSKRM
YPHRPVMMVISHAAPHGPEDSAPQFSKLYPNASQHITPSYNYAPNMDKHWIMQYTGPMPLPIHMEFTNILQR

```
>gi|4507839|ref|NP_003355.1| uridine phosphorylase [Homo sapiens]
MAATGANA-EAESHND-CPVRL-LNPNI-AKMKEDILYHFN-LTTSRH-NFPALFGDVKFVCVGGSPSRMKAFIR
CVGAELGLDC-PGRDYPN-ICAGTD-RYAMYKVG-PVLSVSHGMG-IPSI-SIMLHELKLLYYARC-SNVTIIRIG
TSGGIGLE-PGTVVITE-QAVDTCFKA-EFEQIVL-GKRVIRK-TDLNKKL-VQELL-LSAELSEFTTVVGNTMCT
LDFYEGQ-RLDGLALCSY-TEKDKQAYLEA-AAGVRNI-EMESSVFA-AMCSACGLQA-AVVCV-TLLNRL-EGDQ
ISSPNVRLSEYORPORLYTEYSFIKKL-SKA
```

>gi|11641237|ref|NP_071732.1| RAB38; Rab-related GTP-binding protein
[Homo sapiens]
MQAPHKEHLYKLLVIGDLGVGKTSIIKRYVHQNFSSHYRATIGVDFALKVLHWDPETVVRQLWLDIAGQE
RFGNMTRVYYREAMGAFIVFDVTRPATFEAVAKWKNLDLSKLSLPPNGKPVSVVLLANKCDQGKDVLMMNG
LKMDQFCKEHGFVGWFETSAKENINIDEASRCLVKHILANECDLMESIEPDVVKPHLTSTKVASCSCGCAK
L

```
>gi|8923042|ref|NP_060103.1| chromosome 6 open reading frame 37;  
retinal expressed gene C6orf37 [Homo sapiens]  
MAEGEGYFAMSEDELACSPYIPLGGDFGGGDFGGGDFGGGDFGGGGSFGGHCLDYCESPTAHCNV  
LNWEQVQRLDGI LSETIPIHGRGNFPTLELQPSLIVKVVRRLAEKRIGVRDVLNGSAAASHVLHQDSSL  
GYKDLDLIFCADLRGEGEFQTVKDVLDCLLDFLPEGVNKEKITPLTLKEAYVKMVKVCNDSRWSLIS  
LSNNSGKNVELKFVDSLRRQFEFSVDSFQIKLDSLLLFYECSENPMTETFHPTIIGESVYGDFQEAFDHL  
CNKIIATRNP EPIRGGLLKSTCLLVGRFPASDEIKALQRYMCSRFFIDFSDIGEQQRKLESYLNHFV  
GLEDRKYEYLMTLHGVSNNVNCVLCMGHERRQTLNLITMLAIRVLADQNVIPNVANVTCYYQPAPYVADAN  
FSNYIIAOVQPVFTCCOONTYSTWLPNCN
```

103

MAEESSTRDCMSFSVLNWDQVSRLEHVLTEVVIHGRGNFPTLEITLKDIVQTVRSRLEEAGIKVHDVR
 LNGSAAGHVLVKDNLGCKDLDLIFHVALPTEAEFQLVRDVVLCSSLNFLPEGVNKLKISPVTLKEAYVQ
 KLVKVCCTDTRWLSLISLNKNGKNVELKFVDSIRRQFEFSVDSFQIILDSLLFFYDCSNNPISEHFHPTV
 IGESMYGDFEEAFDHLQNRLIATKNPEEIRGGGLLKYSNLLVRDFRPTDQEEIKTLERYMCSRFFIDFPD
 ILEQQRKLETYLQNHFAEEERSKYDYLMILRRVVNESTVCLMGHERRQTLNLI SLLALRVLAEQNI IPSA
 TNVTCYYQPAPYVSDGNFSNYYVAHPVPTYSQPYPTWLPNCN

>gi|16418427|ref|NP_443175.1| hypothetical protein MGC16491 [Homo sapiens]

MPESGAERRDAAAQVGTAATAVATAAPAGGGPDPEALSAFPGRHLSGLSWPQVKRLDALLSEPIPIH
 GRGNFPTLSVQPRQIVQVVRSTLEEQGLHVHVSRLHGSAAASHVLHPESGLGYKDLDLVFRVDRSEASFQ
 LTKAVVLACLLDFLPAGVSRKITPLTLKEAYVQKLVKVCCTSDRWSLISLNKSGKNVELKFVDSVRRQ
 FEFSIDSFQIILDSLLLFGQCSSTPMSEAFHPTVTGESLYGDFTEALEHLRHRVIATRSPEEIRGGGLLK
 YCHLLVRGFRPRPSTDVRALQRYMCSRFFIDFPDLVEQRRTLERYLEAHFGGADAARRYACLVTLHRVVN
 ESTVCLMNHERRQTLDLIAALALQALAEQGPATAALAWRPPGTDGVVPATVNYVTPVQPLLAHAYPTW
 LPCN

>gi|22749287|ref|NP_689843.1| hypothetical protein MGC26999 [Homo sapiens]

MSEIRFTNLTWDQVITLDQVLDEVIPIHGKGNFPTMEVKPKDI IHVVKDQLIGQGIIIVKDARLNGSVASY
 ILASHNGISYKDLDFVIFGVLPNEEFQVVKDAVLDCLLDFLPKDVKEKLSPDIMKDAYVQKLVKVCNG
 HDCWSLISLNNTGKNLELKFVSSLRRQFEFSVDSFQIVLDPMLDFYSDKNAKLTKESYPVVVAESMYGD
 FQEAMTHLQHKLICTRKPEEIRGGGLLKYSLLVHGFKPACMSEIKNLERYMCSRFFIDFPHIEEQKKI
 ESYLHNHFIGGMTKYDYLMTLHGVVNESTVCLMSYERRQILHLITMMALKVLGELNLPNTQKVTCFYQ
 PAPYFAAEARYPIYVPEPPPVSFQPYHPLHFRGSNGMS

WHAT IS CLAIMED IS:

1. A method of identifying a candidate beta-catenin pathway modulating agent, said method comprising the steps of:
 - (a) providing an assay system comprising a MBCAT polypeptide or nucleic acid;
 - (b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
 - (c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as a candidate beta-catenin pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that express the MBCAT polypeptide.
3. The method of Claim 2 wherein the cultured cells additionally have defective beta-catenin function.
4. The method of Claim 1 wherein the assay system includes a screening assay comprising a MBCAT polypeptide, and the candidate test agent is a small molecule modulator.
5. The method of Claim 4 wherein the assay is a binding assay.
6. The method of Claim 1 wherein the assay system is selected from the group consisting of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay comprising a MBCAT polypeptide and the candidate test agent is an antibody.

8. The method of Claim 1 wherein the assay system includes an expression assay comprising a MBCAT nucleic acid and the candidate test agent is a nucleic acid modulator.
9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.
10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.
11. The method of Claim 1 additionally comprising:
 - (d) administering the candidate beta-catenin pathway modulating agent identified in (c) to a model system comprising cells defective in beta-catenin function and, detecting a phenotypic change in the model system that indicates that the beta-catenin function is restored.
12. The method of Claim 11 wherein the model system is a mouse model with defective beta-catenin function.
13. A method for modulating a beta-catenin pathway of a cell comprising contacting a cell defective in beta-catenin function with a candidate modulator that specifically binds to a MBCAT polypeptide, whereby beta-catenin function is restored.
14. The method of claim 13 wherein the candidate modulator is administered to a vertebrate animal predetermined to have a disease or disorder resulting from a defect in beta-catenin function.
15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.
16. The method of Claim 1, comprising the additional steps of:

(d) providing a secondary assay system comprising cultured cells or a non-human animal expressing MBCAT ,

(e) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and

(f) detecting an agent-biased activity of the second assay system,
wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate beta-catenin pathway modulating agent,
and wherein the second assay detects an agent-biased change in the beta-catenin pathway.

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

19. The method of Claim 18 wherein the non-human animal mis-expresses a beta-catenin pathway gene.

20. A method of modulating beta-catenin pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a MBCAT polypeptide or nucleic acid.

21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the beta-catenin pathway.

22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

23. A method for diagnosing a disease in a patient comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the sample with a probe for MBCAT expression;
- (c) comparing results from step (b) with a control;
- (d) determining whether step (c) indicates a likelihood of disease.

24. The method of claim 23 wherein said disease is cancer.

ABSTRACT OF THE DISCLOSURE

Human MBCAT genes are identified as modulators of the beta-catenin pathway, and thus are therapeutic targets for disorders associated with defective beta-catenin function. Methods for identifying modulators of beta-catenin, comprising screening for agents that modulate the activity of MBCAT are provided.